Candidate Name:



millennia institute

H2 CHEMISTRY	9729/02
Paper 2 Structured Questions	14 Sep 2020
	2 hours
Candidates answer on the Question paper.	
Additional materials: Data Booklet	

READ THESE INSTRUCTIONS FIRST

	Do not tu	urn over this	question	paper until	you are	told to do so
--	-----------	---------------	----------	-------------	---------	---------------

Write your name, class and admission number on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer all questions.

The use of an approved scientific calculator is expected, where appropriate.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

Question	1	2	3	4	5	Total
Marks	11	23	15	15	11	75



Class Adm No

Answer all questions in the spaces provided.

1	(a)	De	scribe the structure of a ³¹ P atom, in terms of the numb	er and type of subatomic particles.	For Examiner use
	(b)	Sta	ate and explain how the atomic radius and anionic radi	us of phosphorus differ.	
				[1]	
	(c)	(i)	Solid PCI ₅ can be prepared by the chlorination of lice state symbols, to represent this chlorination.	quid PC <i>I</i> ₃ . Write an equation, with	
		(ii)	By means of an energy cycle and the information gi	ven below, calculate the standard	
			enthalpy change of chlorination of PCl ₅ .		
			$P(s) + \frac{s}{2}Cl_2(g) \longrightarrow PCl_3(l)$	$\Delta H = -339 \text{ kJ mol}^{-1}$	
			$2P(s) + 5Cl_2(g) \longrightarrow 2PCl_5(s)$	$\Delta H = -926 \text{ kJ mol}^{-1}$	

(iii) Write an equation, including state symbols to illustrate the reaction of PCl_5 with water. For *Examiner's* use Suggest the approximate pH of the solution formed.[2] (d) PCl_5 reacts with hydrazine, N₂H₄, to give a molecular compound **X** with composition by mass shown below and a molecular formula of 303.0. P, 20.5 %; N, 9.2%; Cl, 70.3% (i) Calculate the empirical formula of X and hence, state its molecular formula. [2] (ii) Given that X contains a N–N single bond, suggest the structure of X. [1] [Total: 11]

[Turn over

2 Volcanoes can contribute to air pollution through the release of toxic gases when volcanoes erupt. The most abundant volcanic gas is water vapour. However, significant amounts of carbon dioxide, sulfur dioxide, hydrogen sulfide and hydrogen halides are also emitted.

For Examiner's use

Gas	Volume Percentage
Water vapour, H ₂ O	87.1
Carbon dioxide, CO ₂	unknown
Sulfur dioxide, SO ₂	0.5
Hydrogen, H ₂	0.7
Carbon monoxide, CO	0.01
Hydrogen sulfide, H ₂ S	0.23
Hydrogen halides	unknown

Table 1: Volcanic gas composition in town A

The composition of the volcanic gases are expressed in terms of volume percentage, which can be calculated using the formula below.

volume percentage =
$$\frac{\text{volume of gas}}{\text{total volume}} \times 100\%$$

- (a) When carbon dioxide is emitted from volcanoes, it becomes diluted to low concentrations quickly and is not a harmful gas. However, the carbon dioxide gas when cooled, can flow into low-lying areas where it can reach higher concentrations.
 - (i) Suggest why at low temperatures, carbon dioxide would accumulate to high concentrations. Explain your answer.

.....[2]

For Examiner's use

People living in town **A** had to be evacuated as the levels of carbon dioxide gas was increasing rapidly. Breathing air with more than 3% CO₂ can lead to headaches, increased heart rate and difficulty breathing. At about 15%, unconsciousness and death can result.

(ii) Given that 0.35 g of CO₂ was present in 1 dm³ of gas mixture at r.t.p, determine the volume percentage of carbon dioxide present.

[2]

- (iii) Hence, comment on the possible danger if people remained in the town.[1]
- (b) Further analysis of the hydrogen halides composition of the volcanic gases revealed that the hydrogen halides were mostly made up of HC*l*(g) and HBr(g).

(i)

Table 2: Bond energies of hydrogen halides

Bond energy / kJ mol ⁻¹
432
366

State the trends of **two** chemical properties of hydrogen halides that can be explained by the data given in T**able 2**.

.....[2]

(ii) Explain why HBr(g) deviates more from ideality as compared to HC*l*(g).

.....[1]

(iii) Hydrogen halides can be used in the reaction with alkenes to synthesise halogenoalkanes.
 Draw the displayed formula of both the major and minor products of the reaction of HC/(g) with 2-methylbut-2-ene. Label the major and minor products clearly.

- [2]
- (c) Another volcanic gas, sulfur dioxide, also has harmful effects on human health.
 - (i) Draw the 'dot-and-cross' diagram of one molecule of SO₂ and state its bond angle.

[2]

(ii) SO₂ can be converted SO₃ in the presence of atmospheric oxygen. By using appropriate values from the data booklet, write an equation for the conversion of SO₂ to SO₃ and calculate the enthalpy change of this reaction.

use

(iii) SO₂ can be used to produce H_2SO_4 in a multi-step reaction. In one such reaction, $\begin{vmatrix} For \\ Examiner's \end{vmatrix}$ 100 cm³ of H₂SO₄ was produced. The H₂SO₄ produced was tested for its concentration where 25.0 cm³ of the resultant solution required 23.50 cm³ of 2.00 mol dm⁻³ NaOH for complete titration.

Calculate the amount of H₂SO₄ produced in the method above.

[2]

- (d) Student A suggested that one possible method to remove harmful SO₂ from volcanic gases is to react the gaseous SO_2 with magnesium oxide.
 - (i) Explain the Chemistry behind why this method is possible.

.....[1]

Student **B** suggested that besides magnesium oxide, other metal oxides like sodium oxide and calcium oxide can also be used to remove sulfur dioxide.

(ii) The pH of the resulting solutions when one mole of MgO and Na₂O are added to 1 dm³ of water separately are shown in **Table 3** below.

Metal Oxide	pH of resulting solution
MgO	9
Na ₂ O	13
CaO	?

T	ab	le	3:	pН	of	metal	oxides

Predict and suggest an explanation for the pH of a solution containing For Examiner's use



Student **C** suggested that seawater can also be used to absorb and remove SO₂. The main substance in seawater that is responsible for removing SO₂ is the HCO_3^- ions. When SO₂ is absorbed in water, the following two equilibria reactions take place:

 $SO_2(g) + H_2O(l) + \frac{1}{2}O_2(g) \rightleftharpoons SO_4^{2-}(aq) + 2H^+(aq) - (1)$ $HCO_3^-(aq) + H^+(aq) \rightleftharpoons H_2O(l) + CO_2(g) - (2)$ (from seawater)

(iii) Explain how the use of seawater would allow the removal of sulfur dioxide.

[Total: 23]

The Kolbe electrolysis is an electrochemical method used to synthesise alkanes. Carbon 3 (a) dioxide and ethane was produced during the Kolbe electrolysis of an aqueous solution of potassium ethanoate.

The reaction mechanism involves a three-stage process.

In step I, the ethanoate ion CH₃COO⁻ is first converted into an ethoxy radical intermediate CH₃COO•.

In step II, this intermediate then undergoes decarboxylation to form a methyl radical intermediate.

step II $CH_3COO \rightarrow \bullet CH_3 + CO_2$

In step III, two methyl radical intermediates dimerises to form the alkane product. step III 2 \bullet CH₃ \rightarrow CH₃-CH₃

(i) Given that the reaction in step I takes place at the anode, write a half-equation for the reaction.

.....[1]

(ii) Write another half-equation to illustrate the formation of a gaseous product at the cathode.

.....[1]

(iii) Calculate the volume of gas produced at the cathode at room temperature and pressure when a current of 500 mA is passed for 20 minutes through a solution of potassium ethanoate.

[3]

use

(iv) Use the information given above to draw out the full mechanism for steps II and III. Use For Examiner's appropriate curly arrows to indicate which bonds are broken and which bonds are formed.

use

[2]

(v) In a separate Kolbe electrolysis experiment, an alkane X was produced when a mixture of an ethanoate salt and the salt of a carboxylic acid was electrolysed.

A gaseous sample of 0.30 g of X occupied 168 cm³ at 300 K and 1 atm. Calculate the M_r of **X** and hence deduce its molecular formula.

(b) Galvanic cells are electrochemical cells that contain a spontaneous reaction. An example of a galvanic cell include the zinc-carbon battery. Ammonium chloride is used as electrolyte in zinc–carbon batteries, where the outer zinc casing is the negatively charged terminal and is oxidised during discharge.

The following reaction also takes place in the cell.

 $2MnO_2(s) + 2e^- + 2NH_4Cl(aq) \rightarrow Mn_2O_3(s) + 2NH_3(aq) + H_2O(l) + 2Cl^-(aq) --- E^{\Theta} = +0.5 V$

(i) Write a half-equation for the reaction occurring at the negatively charged terminal. Hence, write an overall equation for the reaction that occurs in the cell.

......[2]

(ii) Calculate the voltage that is generated by this cell.

[1]

(iii) Calculate a value of ΔG^{Θ} for the cell reaction, and explain the significance of its sign.

[2]

[Total: 15]



A reaction scheme is shown below. 4 (a)

For

use

(b) Compare the acidicity between phenol and 2,4,6-trinitrophenol. State and explain which of the two compounds is more acidic.

.....[2]

(c) The amino acid, alanine, has the structure shown below.

alanine

(i) By drawing an appropriate structure, explain why alanine has a surprisingly high melting point considering its relatively small molecular mass.

(ii) Solutions of amino acids are buffers important in maintaining optimal pH in our body.

With the aid of an equation, show how alanine behaves as a buffer when small amount of base is added.

For Examiner's use

[2]

[Total: 15]

- 5 (a) Aqueous magnesium chloride can act as a weak monobasic acid. In an experiment, 50.0 cm³ of an aqueous solution of magnesium chloride at an initial pH of 6 was titrated with 1.00 mol dm⁻³ sodium hydroxide. 20 cm³ of NaOH(aq) was required for complete neutralisation.
 - (i) By writing suitable equations, illustrate how aqueous magnesium chloride can act as a weak monobasic acid.

.....[2]

(ii) Calculate the initial concentration of magnesium ions.

(iii) Calculate the K_a value for a solution of aqueous magnesium chloride.

[2]

[1]

- In a different experiment, a solution containing 0.100 mol dm⁻³ magnesium chloride and (b) Examiner's 0.100 mol dm⁻³ barium chloride was prepared. Solid sodium carbonate was added slowly to 1 dm³ of this solution in an attempt to separate the two metal cations. [Given, K_{sp} of MgCO₃ = 3.5×10^{-8} mol² dm⁻⁶; K_{sp} of BaCO₃ = 5.1×10^{-9} mol² dm⁻⁶]
 - (i) Calculate the concentration of carbonate in the solution, needed for the first trace of precipitate to be seen, stating which metal ion is precipitated first.

[2]

For

use

(ii) Determine the concentration of the metal ion in (b)(i) remaining in the solution when the other metal ion just starts to precipitate.

- [2]
- (iii) Given that an effective separation means that less than 1% of a metal ion should remain in solution, deduce if the separation of the metal ion in (b)(i) has been effective.

[Total: 11]

BLANK PAGE