

NAME _____

CLASS 1T _____



Catholic Junior College
JC1 Promotional Examinations
Higher 2

PHYSICS

9749/1

Paper 1: Multiple Choice Questions

30 September 2022

30 minutes

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write your name and tutorial group on this cover page.

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write and shade your name, NRIC / FIN number and HT group on the Answer Sheet (OMR sheet), unless this has been done for you.

There are **fifteen** questions on this paper. Answer **all** questions. For each question, there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet (OMR sheet).

Read the instructions on the Answer Sheet carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

The use of an approved scientific calculator is expected, where appropriate.

PHYSICS DATA:

| | | |
|------------------------------|--------------|--|
| speed of light in free space | c | $= 3.00 \times 10^8 \text{ m s}^{-1}$ |
| permeability of free space | μ_0 | $= 4\pi \times 10^{-7} \text{ H m}^{-1}$ |
| permittivity of free space | ϵ_0 | $= 8.85 \times 10^{-12} \text{ F m}^{-1}$ $\approx (1/(36\pi)) \times 10^{-9} \text{ F m}^{-1}$ |
| elementary charge | e | $= 1.60 \times 10^{-19} \text{ C}$ |
| the Planck constant | h | $= 6.63 \times 10^{-34} \text{ J s}$ |
| unified atomic mass constant | u | $= 1.66 \times 10^{-27} \text{ kg}$ |
| rest mass of electron | m_e | $= 9.11 \times 10^{-31} \text{ kg}$ |
| rest mass of proton | m_p | $= 1.67 \times 10^{-27} \text{ kg}$ |
| molar gas constant | R | $= 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ |
| the Avogadro constant | N_A | $= 6.02 \times 10^{23} \text{ mol}^{-1}$ |
| the Boltzmann constant | k | $= 1.38 \times 10^{-23} \text{ mol}^{-1}$ |
| gravitational constant | G | $= 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$ |
| acceleration of free fall | g | $= 9.81 \text{ m s}^{-2}$ |

PHYSICS FORMULAE:

| | |
|--|--|
| uniformly accelerated motion | $s = ut + \frac{1}{2}at^2$ |
| | $v^2 = u^2 + 2as$ |
| work done on / by a gas | $W = p\Delta V$ |
| hydrostatic pressure | $P = \rho gh$ |
| gravitational potential | $\phi = -\frac{Gm}{r}$ |
| temperature | $T/K = T/^{\circ}\text{C} + 273.15$ |
| pressure of an ideal gas | $p = \frac{1}{3} \frac{Nm}{V} \langle c^2 \rangle$ |
| mean translational kinetic energy of an ideal gas molecule | $E = \frac{3}{2}kT$ |
| displacement of particle in s.h.m. | $x = x_0 \sin \omega t$ |
| velocity of particle in s.h.m. | $v = v_0 \cos \omega t$ $= \pm \omega \sqrt{x_0^2 - x^2}$ |
| electric current | $I = Anvq$ |
| resistors in series | $R = R_1 + R_2 + \dots$ |
| resistors in parallel | $1/R = 1/R_1 + 1/R_2 + \dots$ |
| electric potential | $V = \frac{Q}{4\pi\epsilon_0 r}$ |
| alternating current / voltage | $x = x_0 \sin \omega t$ |
| magnetic flux density due to a long straight wire | $B = \frac{\mu_0 I}{2\pi d}$ |
| magnetic flux density due to a flat circular coil | $B = \frac{\mu_0 NI}{2r}$ |
| magnetic flux density due to a long solenoid | $B = \mu_0 nI$ |
| radioactive decay | $x = x_0 \exp(-\lambda t)$ |
| decay constant | $\lambda = \frac{\ln 2}{t_{\frac{1}{2}}}$ |


- 1 The resistance of a resistor is measured using an ohmmeter and its reading is $47.12\ \Omega$. The manufacturer of the ohmmeter specifies that the ohmmeter reading has an uncertainty of 10%.

How should the resistance of the resistor be recorded, along with its uncertainty?

- A $(47 \pm 5)\ \Omega$
 B $(47.1 \pm 0.1)\ \Omega$
 C $(47.1 \pm 4.7)\ \Omega$
 D $(47.12 \pm 5)\ \Omega$
- 2 Two trains, Train A and Train B, are initially at rest next to each other at a train station. Train A leaves the station with a uniform acceleration of $0.30\ \text{m s}^{-2}$. 2.0 minutes later, Train B leaves the station in the same direction as Train A with a uniform acceleration of $0.50\ \text{m s}^{-2}$.

How long does Train B take to catch up with Train A?

- A 6.9 s B 180 s C 210 s D 410 s
- 3 A stone is thrown from X and moves under gravity to Y, at the top of its path.



Which row correctly describes its motion horizontally and vertically between X and Y? Neglect air resistance.

| | horizontally | vertically |
|---|--------------------------------|------------------------------|
| A | travels at decreasing velocity | velocity decreases uniformly |
| B | travels at increasing velocity | velocity increases uniformly |
| C | travels at constant velocity | velocity increases uniformly |
| D | travels at constant velocity | velocity decreases uniformly |

- 4 Two blocks with masses of 500 g and 400 g are moving towards each other at speeds of 2.0 m s^{-1} and 3.0 m s^{-1} as shown.



The two blocks stick together after the collision and move off with a common speed.


What is the speed and the direction of motion of the blocks after the collision?

- A** 0.22 m s^{-1} to the left
B 0.22 m s^{-1} to the right
C 2.4 m s^{-1} to the left
D 2.4 m s^{-1} to the right
- 5 A tennis ball of mass 60 g is dropped vertically from a height. It hits the ground with a speed of 21 m s^{-1} downwards and then bounces upwards with an initial speed of 14 m s^{-1} . The time of contact between the ball and the ground is 0.30 s.

What is the magnitude of the impulse provided by the ground?

- A** 0.42 N s **B** 2.1 N s **C** 7.0 N s **D** 2100 N s

- 6 The unstretched length of a spring is 30 cm. It is stretched vertically, with one end fixed on a ceiling, to a length of 40 cm by a weight of 2 N attached to the other end as shown. The weight is at rest.




What is the potential energy stored in the spring?

- A** 0.1 J **B** 0.2 J **C** 0.6 J **D** 0.8 J
- 7 A wooden block of density 700 kg m^{-3} is completely submerged below the surface of water by a weight placed on top of it. The block floats to the surface when the weight is removed. The density of water is 1000 kg m^{-3} .

What is the ratio $\frac{\text{upthrust on the wooden block when it is completely submerged}}{\text{upthrust on the wooden block when it is floating}}$?

- A** 0.143 **B** 0.700 **C** 1.43 **D** 7.00

- 8 An object of mass m is hung at one end of a 3.0 m long uniform rod of mass M and placed on a pivot. For the system to be in equilibrium, the pivot has to be placed 1.2 m from the end where the object is hung. When the object's mass is doubled to $2m$, the pivot has to be moved a distance x to the right of its original position for the system to remain in equilibrium as shown.




What is the distance x ?

- A** 0.20 m **B** 0.40 m **C** 0.50 m **D** 0.80 m
- 9 A car with a power output of 100 kW is travelling with a constant speed of 30 m s^{-1} along a straight road. The frictional force on the car is proportional to the square of its speed.





What constant speed will it be travelling at if its power output is 200 kW?

- A** 37.8 m s^{-1} **B** 42.4 m s^{-1} **C** 51.6 m s^{-1} **D** 60.0 m s^{-1}

- 10 The variation with displacement s of the total work done on an object W is shown below.



Which of the following graphs best represents the variation with displacement s of the resultant force F acting on the object?



- 11 An object of mass m is attached to a rigid rod and spun in a uniform vertical circular motion with radius r and a linear velocity of v .

What is the variation of the magnitude of the centripetal acceleration of the object over one revolution?

- A** zero **B** $2g$ **C** $\frac{v^2}{r}$ **D** $2\frac{v^2}{r}$

[Turn over]

12 Which quantity is **not necessarily** the same for satellites in geostationary orbits around the Earth?

- A** angular velocity
- B** centripetal acceleration
- C** kinetic energy
- D** orbital period

13 A mass on a frictionless table is attached to a fixed horizontal spring and displaced a certain horizontal distance from its equilibrium position. When released, it oscillates in a simple harmonic motion with an angular frequency of 3.14 rad s^{-1} .

At what times after its release does the object pass through the equilibrium position?

- A** 0.25 s, 0.75 s, 1.25 s...
- B** 0.25 s, 1.25 s, 2.25 s...
- C** 0.50 s, 1.00 s, 1.50 s...
- D** 0.50 s, 1.50 s, 2.50 s...


14 A sound wave has a frequency of 2500 Hz and a speed of 1500 m s^{-1} .

What is the shortest distance from a point of maximum pressure in the wave to a point of minimum pressure?

- | | | | | | | | |
|----------|--------|----------|--------|----------|--------|----------|--------|
| A | 0.15 m | B | 0.30 m | C | 0.60 m | D | 1.20 m |
|----------|--------|----------|--------|----------|--------|----------|--------|

- 15** A narrow, parallel beam of unpolarised light is directed towards three ideal polarising filters.

The beam meets the first filter with a vertical axis of polarisation and emerges with an intensity I_0 . The axis of polarisation of the second filter is at an angle of 10° to the first filter. The third filter has its axis of polarization parallel to the second filter as shown.



The third filter is now turned an angle of 34° to the second filter. What is the intensity of the transmitted light after passing through the third filter?

- A** $0.25 I_0$ **B** $0.33 I_0$ **C** $0.5 I_0$ **D** $0.67 I_0$

BLANK PAGE