2019 H2 Math Prelim - ASRJC

1

1 (a) Show that
$$\sqrt{\frac{1-2x}{1+2x}}$$
 can be written in the form $\frac{f(x)}{\sqrt{1-4x^2}}$ where $f(x)$ is a polynomial to be determined. Hence, find $\int \sqrt{\frac{1-2x}{1+2x}} dx$ [3]

(b) Show that
$$\int_{e}^{e^2} \frac{1}{x \ln x} dx = \ln 2$$
. [3]

2 It is given that
$$\tan \frac{1}{2}y = \sqrt{2}x$$
, where $-\frac{\pi}{2} < y < \frac{\pi}{2}$.

(i) Show that
$$(1+2x^2)\frac{dy}{dx} = 2\sqrt{2}$$
 . [2]

(iii) Hence, using standards series from the List of Formulae (MF26), find the expansion of
$$\frac{2 \tan^{-1} \sqrt{2x}}{\cos 2x}$$
 in ascending powers of x, up to and including the term in x^3 , giving the coefficients in exact form. [3]

- **3** The position vectors of *A*, *B* and *C* referred to a point *O* are **a**, **b** and **c** respectively. The point *N* is on *AB* such that AN:NB = 2:1.
 - (i) If O is the midpoint of CN, prove that $\mathbf{a} + 2\mathbf{b} + 3\mathbf{c} = \mathbf{0}$. [2]
 - (ii) Show that A, O and M are collinear, and find the ratio AO:OM [3]
 - (iii) If the point *P* is such that $\overline{NP} = \overline{AM}$, show that the ratio of the area of *PNAM* : area of *PNOM* = 12:7 [3]



The diagram shows the graph of y = f(x). The curve crosses the x-axis at x = 0 and x = 3. It has a turning point at (4, 5) and asymptotes with equations y = 2 and x = 2.

Showing clearly the coordinates of turning points, axial intercepts and equations of asymptotes where possible, sketch the graphs of

(i)
$$y = f(|x|);$$
 [2]

(ii)
$$y = f'(x)$$
 [3]

(b) The curve with equation y = f(x) is transformed by a stretch with scale factor 2 parallel to the *x*-axis, followed by a translation of 2 units in the negative *x*-direction, followed by a translation of 3 units in the positive *y*-direction. The equation of the resulting curve is $y = \ln(e^3x)$.

Find the equation of the curve y = f(x). [3]

5 A curve C has parametric equations

$$x = a \sin^2 t$$
, $y = a \cos t$,

where $0 \le t \le \frac{\pi}{2}$ and a > 0.

- (i) Find the cartesian equation of *C*, stating clearly any restrictions on the values of *x* and *y*. [2]
- (ii) Sketch *C*, showing clearly the axial intercepts. [1]
- (iii) The region bounded by *C*, the line $y + x = \frac{5}{4}a$ and the *y*-axis is rotated through 2π radians about the *y*-axis. Show that the exact volume of the solid obtained is $k\pi a^3$ where *k* is a constant to be determined. [5]
- 6 A function f is defined by

$$f(x) = \begin{cases} \frac{6}{x-4} & \text{for } x < -2, \\ -|1+x| & \text{for } -2 \le x \le 1, \\ x^2 - x - 2 & \text{for } x > 1. \end{cases}$$

(ii) Evaluate exactly $\int_{-3}^{4} |f(x)| dx$. [4]

7 Do not use a calculator in answering this question.

The roots of the equation $z^2 + (2-2i)z = -3 - 2i$ are z_1 and z_2 .

- (i) Find z_1 and z_2 in cartesian form x + iy, showing clearly your working. [5]
- (ii) The complex numbers z_1 and z_2 are also roots of the equation

$$z^4 + 4z^3 + 14z^2 + 4z + 13 = 0.$$

Find the other roots of the equation, explaining clearly how the answers are obtained. [2]

(iii) Using your answer in part (i), solve $z^2 + (2+2i)z = 3+2i$. [2]

8 (i) Show that $\frac{3r+4}{(r+2)(r+1)r} = \frac{A}{r+2} + \frac{B}{r+1} + \frac{C}{r}$ where A, B and C are constants to be determined. [1]

(ii) Find the sum to *n* terms of

9

$$\frac{7}{3\times2\times1} + \frac{10}{4\times3\times2} + \frac{13}{5\times4\times3} + \dots$$

(There is no need to express your answer as a single algebraic fraction). [4]

(iii) Hence show that
$$\sum_{r=5}^{n+3} \frac{3r-5}{(r-1)(r-2)(r-3)} < \frac{4}{3}$$
. [3]



The diagram above shows an object with O at the centre of its rectangular base ABCD where AB = 8 cm and BC = 4 cm. The top side of the object, EFGH is a square with side 2 cm long and is parallel to the base. The centre of the top side is vertically above O at a height of h cm.

(i) Show that the equation of the line *BG* may be expressed as $\mathbf{r} = \begin{pmatrix} 4 \\ -2 \\ 0 \end{pmatrix} + t \begin{pmatrix} -3 \\ 1 \\ h \end{pmatrix}$,

where *t* is a parameter.

(ii) Find the sine of the angle between the line BG and the rectangular base ABCD in terms of h. [2]

[1]

It is given that h = 6.

- (iii) Find the cartesian equation of the plane BCFG. [3]
- (iv) Find the shortest distance from the point A to the plane BCFG. [2]

- (v) The line l, which passes through the point A, is parallel to the normal of plane BCFG. Given that, the line l intersects the plane BCFG at a point M, use your answer in part (iv) to find the shortest distance from point M to the rectangular base ABCD.
- 10 Commonly used in building materials, sand is the second largest world resource used by humans after water. To reduce the environmental impacts of sand mining, an alternative approach is to make "sand" by crushing rock.

A machine designed for this purpose produces sand in large quantities. Sand falling from the output chute of the machine forms a pile in the shape of a right circular cone such that the height of the cone is always equal to $\frac{4}{3}$ of the radius of its base.

[Volume of a cone = $\frac{1}{3}\pi r^2 h$ and curved surface area of a cone = $\pi r l$ where *r* is the radius of the base area, *h* is the height of the cone and *l* is the slant length of the cone]



A machine operator starts the machine.

- (i) Given that V and A denote the volume and the curved surface area of the conical pile respectively, write down V and A in terms of r, the radius of its base.
 - (ii) Hence show that the rate of change of A with respect to V is inversely proportional to the radius of the conical pile. [3]

An architect is tasked to design sand-lined walking paths in a large park. He decides to base his design of the paths on the shape of astroids, which are shapes with equations

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = k^{\frac{2}{3}} \ (k \ge 0).$$

On a piece of graph paper, he sketches an astroid with the equation $x^{\frac{2}{3}} + y^{\frac{2}{3}} = k^{\frac{2}{3}}$.

(b) The tangent at a point $P(x_1, y_1)$ on the curve meets the x-axis at Q and the y-axis at R. Show that the length of QR is independent of where P lies on the curve. [7] 11 (a) Find the sum of all integers between 200 and 1000 (both inclusive) that are not divisible by 7. [4]



Snowflakes can be constructed by starting with an equilateral triangle (Fig. 1), then repeatedly altering each line segment of the resulting polygon as follows:

- 1. Divide each outer line segments into three segments of equal length.
- 2. Add an equilateral triangle that has the middle segment from step 1 as its base.
- 3. Remove the line segment that is the base of the triangle from step 2.
- 4. Repeat the above steps for a number of iterations, *n*.

The 1st, 2nd and 3rd iteration produces the snowflakes in Fig. 2, Fig. 3 and Fig. 4 respectively.

(i) If a_0 denotes the area of the original triangle and the area of each new triangle added in the n^{th} iteration is denoted by a_n , show that

$$a_n = \frac{1}{9}a_{n-1}$$
 for all positive integers *n*. [2]

- (ii) Write down the number of sides in the polygons in Fig.1, Fig. 2 and Fig. 3, and deduce, with clear explanations, that the number of new triangles added in the n^{th} iteration is $T_n = k(4^n)$ where k is a constant to be determined. [2]
- (iii) Find the total area of triangles added in the n^{th} iteration, A_n in terms of a_0 and n. [2]
- (iv) Show that the total area of the snowflake produced after the n^{th}

iteration is
$$a_0 \left[\frac{8}{5} - \frac{3}{5} \left(\frac{4}{9} \right)^n \right]$$
 units². [3]

The snowflake formed when the above steps are followed indefinitely is called the Koch snowflake.

(v) Determine the least number of iterations needed for the area of the snowflake to exceed 99% of the area of a Koch snowflake. [3]

Section A: Pure Mathematics [40 marks]

1 Given that the curve
$$y = \frac{ax^2 + bx + c}{x - 1} = f(x)$$
 passes through the points (-1, -3),

$$\left(\frac{1}{2}, -\frac{3}{2}\right)$$
 and $\left(5, \frac{33}{2}\right)$, find the values of *a*, *b* and *c*. [3]

Hence find the exact range of values of x for which the gradient of y = f(x) is positive. [3]

2 The functions f and g are defined by:

f:
$$x \mapsto |ax - x^2|$$
, $x \in \mathbb{R}$
g: $x \mapsto ae^{-x} + 1$, $x \ge 0$

where *a* is a positive constant and $a \ge 5$.

(i) Sketch the graphs of y = g(x) and y = f(x) on separate diagrams.

Hence find the range of the composite function fg, leaving your answer in exact form in terms of a. [5]

- (ii) Given that the domain of f is restricted to the subset of R for which x ≥ k, find the smallest value of k, in terms of a, for which f⁻¹ exists. Hence, without finding the expression for f⁻¹(x), sketch the graphs of y = f(x) and y = f⁻¹(x) on a single diagram, stating the exact coordinates of any points of intersection with the axes. [3]
- (iii) With the restricted domain of f in part (ii), find with clear working, the set of values of x such that $f f^{-1}(x) = f^{-1} f(x)$. [2]

3 A student is investigating a chemical reaction between two chemicals A and B. In his experiments, he found that the reaction produces a new product C which does not react with A and B after being formed.

His experiments also suggest that the rate at which the product, C is formed is given by the differential equation

$$\frac{\mathrm{d}x}{\mathrm{d}t} = (a - x)\sqrt{b - x},$$

where a and b are the initial concentrations (gram/litre) of A and B just before the reaction started and x is the concentration (gram/litre) of C at time t (mins).

(i) If a = b = 9, solve the differential equation to show that

$$x = 9 - \frac{36}{(3t+2)^2}.$$
 [4]

(ii) Sketch the graph of x and describe the behaviour of x as t increases. Find the amount of time needed for the concentration of C to reach 4.5 gram/litre. [3]

It is now given that a > b instead.

- (iii) Solve the differential equation to find t in terms of x, a and b by using the substitution $u = \sqrt{b-x}$ for $0 \le x < b$. [6]
- 4 The illumination of an object by a light source is directly proportional to the strength of the source and inversely proportional to the square of the distance from the source.

A small object, P, is placed on the straight line passing through two light sources A and B that are eight metres apart. It is known that the light sources are similar except that S, the strength of light source A, is a positive constant and is three times that of light source B. It can be assumed that I, the total illumination of the object P by the two light source is the sum of the illumination due to each light source.

The distance between the object P and light source A is denoted by x.

(i) Show that when *P* is between *A* and *B*,

$$I = kS \left[\frac{1}{x^2} + \frac{1}{\alpha (\beta - x)^2} \right]$$
 where k is a positive constant and,
nts to be determined. [1]

 α and β are constants to be determined.

(ii) Show that there is only one value of x that will give a stationary value of I. Determine this value and the nature of the stationary value without the use of a graphic calculator. [5]

It is now given that
$$I = \frac{4x^2 - 48x + 192}{x^2 (x^2 - 16x + 64)}$$
 for $0 < x < 8$.

www.KiasuExamPaper.com 52

- (iii) Sketch the graph of *I*, indicating clearly any equations of asymptote and coordinates of the stationary point. [3]
- (iv) It is desired to place P such that $I = \sqrt{\lambda (x x_1)^2}$ where x_1 is the value of x found in part (ii) and λ is a positive constant. Find the value of λ if there is only one possible position to place P. [2]

Section B: Statistics [60 marks]

5 Greenhouse gases generated by Singapore comes mainly from the burning of fossil fuels to generate energy for industries, buildings, households and transportation.

The following table* shows the total greenhouse gas emissions, X (kilotonnes of CO₂), and the total deaths in Singapore, Y from 2007 to 2016.

Year		2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Total 4261		42613	41606	43100	47063	49930	48712	50299	50908	51896	51519
Greenhouse											
Gas Emissions											
(X)											
Total	Deaths	17140	17222	17101	17610	18027	18481	18938	19393	19862	20017
(Y)											

*Extracted from www.singstat.gov.sg

- (i) Draw a scatter diagram for these values, labelling the axes and state the product moment correlation between *X* and *Y*. Use your diagram to explain why the best model for the relationship between *X* and *Y* may not be given by Y = aX + b, where *a* and *b* are constants. [3]
- (ii) Comment on the correctness of the following statement.

"Since there exists a high positive linear correlation between total greenhouse gas emission and total deaths, the deaths are a result of the greenhouse gas emissions." [1]

(iii) Explain clearly which of the following equations, where *a* and *b* are constants, provides the more accurate model of the relationship between *X* and *Y*.

(A)
$$Y = aX^4 + b$$

(B) $\lg Y = aX + b$ [3]

The Government has pledged that Singapore's greenhouse gas emissions will peak around 2030 at the equivalent of 65 million tonnes of carbon dioxide.

(iv) Using the model you have chosen in part (iii), write down the equation for the relationship, giving the regression coefficients correct to 5 significant figures. Hence find the estimated total deaths in the year 2030 if Singapore's pledge is fulfilled and comment on the reliability of the estimate. [3]

6 Birth weight can be used to predict short and long-term health complications for babies. Studies show that the birth weight of babies born to mothers who do not smoke in a certain hospital can be assumed to follow a normal distribution with mean 3.05 kg and variance $\sigma^2 \text{ kg}^2$.

Birth weight	Classification
Less than 1.5 kg	Very low birth weight
1.5 kg to 2.5 kg	Low birth weight
2.5 kg to 4.0 kg	Normal birth weight
More than 4.0 kg	High birth weight

The hospital classifies babies based on their birth weight as shown in the table below.

(i) A sample showed that 20.2% of the babies born to mothers who do not smoke have low birth weight. If this is true for the entire population, find two possible values of σ , corrected to 2 decimal places. Explain clearly why one of the values of σ found should be rejected. [4]

Studies also show that babies born to mothers who smoke have a lower mean birth weight of 2.80 kg.

For the remaining of the question, you may assume the birth weight of babies born to mothers who smoke is also normally distributed and that the standard deviations for the birth weight of babies born to mothers who do not smoke and mothers who smoke are both 0.86 kg.

- (ii) Three pregnant mothers-to-be who smoke are randomly chosen. Find the probability that all their babies will not be of normal birth weight. State an assumption that is needed for your working.
 [3]
- (iii) Find the probability that the average birth weight of two babies from mothers who do not smoke differs from twice the birth weight of a baby from a mother who smokes by less than 2 kg.[3]

Babies whose birth weight not classified as normal will have to remain in hospital for further observation until their condition stabilises. Depending on the treatment received and length of stay, the mean hospitalisation cost per baby is \$2800 and standard deviation is \$500.

(iv) Find the probability that the average hospitalisation cost from a random sample of 50 babies, whose birth weight not classified as normal, exceeds \$3000. [2]

- Three cards are to be drawn at random without replacement from a pack of six cards numbered 0, 0, 1, 1, 1, 1. The random variables X_1, X_2 and X_3 denotes the numbers on the first, second and third card respectively.
 - (i) If $Y = X_1 X_2 + X_3$, show that $P(Y = 1) = \frac{1}{3}$ and find the probability distribution of *Y*. [3]

[2]

[Turn Over

(ii) Find the values of E(Y) and Var(Y).

7

- (iii) Find the probability that the difference between Y and E(Y) is less than one standard deviation of Y. [2]
- 8 A factory makes a certain type of muesli bar which are packed in boxes of 20. The factory claims that each muesli bar weigh at least 200 grams. Muesli bars that weigh less than 200 grams are sub-standard. It is known that the probability that a muesli bar is sub-standard is 0.02 and the weight of a muesli bar is independent of other muesli bars.
 - (i) Find the probability that, in a randomly chosen box of muesli bars, there are at least one but no more than five sub-standard muesli bars. [2]

The boxes of muesli bars are sold in cartons of 12 boxes each.

(ii) Find the probability that, in a randomly chosen carton, at least 75% of the boxes will have at most one sub-standard muesli bar. [2]

As part of quality control for each batch of muesli bars produced daily, one box of muesli bars is randomly selected to be tested. If all the muesli bars in the box weigh at least 200 grams, the batch will be released for sales. If there are more than one substandard muesli bar, the batch will not be released. If there is exactly one sub-standard muesli bar, a second box of muesli bars will be tested. If all the muesli bars in the second box weigh at least 200 grams, the batch will be released. Otherwise, the batch will not be released.

(iii)Find the probability that the batch is released on a randomly selected day. [2](iv) If the factory operates 365 days in a year, state the expected number of days in the year when the batch will not be released. [1]

To entice more people to buy their muesli bars, a scratch card is inserted into each box. The probability that a scratch card will win a prize is *p*. It may be assumed that whether a scratch card will win a prize or not, is independent of the outcomes of the other scratch cards.

Jane buys a carton of muesli bars.

(v) Write in terms of *p*, the probability that Jane win 2 prizes. [1]

The probability that Jane win 2 prizes is more than twice the probability that she win 3 prizes.

(vi) Find in exact terms, the range of values that *p* can take. [2]

9 In preparation for an upcoming event, a student management team is considering having meetings on this Friday, Saturday and Sunday. The probability that the meeting is conducted on Friday is $\frac{1}{6}$. On each of the other days, the probability that a meeting is conducted when a meeting has already been conducted on the previous day, is $\frac{2}{5}$ and the probability that a meeting is conducted, when a meeting has not been conducted on the previous day, is $\frac{1}{3}$.

Find the probability that

(i) a meeting is conducted on Sunday, [3]
 (ii) a meeting is not conducted on Friday given that there is a meeting on Sunday. [3]

The student management team consists of 7 girls and 3 boys. At one of the meetings, they sit at a round table with 10 chairs.

(iii) Find the probability that the girls are seated together. [2]

Two particular boys are absent from the meeting.

- (iv) Find the probability that two particular girls do not sit next to each other. [3]
- 10 Along a 3km stretch of a road, the speed in km/h of a vehicle is a normally distributed random variable *T*. Over a long period of time, it is known that the mean speed of vehicles traveling along that stretch of the road is 90.0 km/h. To deter speeding, the traffic governing body introduced a speed monitoring camera. Subsequently, the speeds of a random sample of 60 vehicles are recorded. The results are summarised as follows.

$$\sum t = 5325, \qquad \sum \left(t - \bar{t}\right)^2 = 2000.$$

- (i) Find unbiased estimates of the population mean and variance, giving your answers to 2 decimal places. [2]
- (ii) Test, at the 5% significance level, whether the speed-monitoring camera is effective in deterring the speeding of vehicles on the stretch of road. [4]
- (iii) In another sample of size n (n > 30) that was collected independently, it is given that $\overline{t} = 89.0$. The result of the subsequent test using this information and the unbiased estimate of the population variance in **part (i)** is that the null hypothesis is not rejected. Obtain an inequality involving n, and hence find the largest possible value n can take. [4]

www.KiasuExamPaper.com 57



ANDERSON SERANGOON JUNIOR COLLEGE

MATHEMATICS

9758

H2 Mathematics Paper 1 (100 marks)

30 August 2019

3 hours

Additional Material(s): List of Formulae (MF26)

CANDIDATE NAME	
CLASS	

READ THESE INSTRUCTIONS FIRST

Write your name and class in the boxes above.

Please write clearly and use capital letters.

Write in dark blue or black pen. HB pencil may be used for graphs and diagrams only.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions and write your answers in this booklet. Do not tear out any part of this booklet.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

You are expected to use an approved graphing calculator.

Where unsupported answers from a graphing calculator are not allowed in a question, you are required to present the mathematical steps using mathematical notations and not calculator commands.

All work must be handed in at the end of the examination. If you have used any additional paper, please insert them inside this booklet.

The number of marks is given in brackets [] at the end of each question or part question.



Question number	Marks
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
Total	

This document consists of 23 printed pages and 1 blank page.

1 (a) Show that
$$\sqrt{\frac{1-2x}{1+2x}}$$
 can be written in the form $\frac{f(x)}{\sqrt{1-4x^2}}$ where $f(x)$ is a polynomial to be determined. Hence, find $\int \sqrt{\frac{1-2x}{1+2x}} dx$ [3]

<u>Solution</u>

$$\int \sqrt{\frac{1-2x}{1+2x}} \, dx = \int \frac{\sqrt{1-2x}}{\sqrt{1+2x}} \times \frac{\sqrt{1-2x}}{\sqrt{1-2x}} \, dx$$
$$= \int \frac{1-2x}{\sqrt{1-4x^2}} \, dx$$
$$= \int \frac{1}{\sqrt{1-4x^2}} - \frac{2x}{\sqrt{1-4x^2}} \, dx$$
$$= \frac{1}{2} \int \frac{2}{\sqrt{1-(2x)^2}} \, dx + \frac{1}{2} \int \left(\frac{-8x}{2}\right) (1-4x^2)^{\frac{-1}{2}} \, dx$$
$$= \frac{1}{2} \sin^{-1} 2x + \frac{1}{2} \sqrt{1-4x^2} + c \text{, where } c \text{ is an arbitrary constant.}$$

(b) Show that
$$\int_{e}^{e^2} \frac{1}{x \ln x} dx = \ln 2$$
. [3]

$$\int_{e}^{e^{2}} \frac{1}{x \ln x} dx = \int_{e}^{e^{2}} \frac{1}{x \ln x} dx$$
$$= \left[\ln \left| \ln x \right| \right]_{e}^{e^{2}}$$
$$= \ln \left| \ln e^{2} \right| - \ln \left| \ln e \right|$$
$$= \ln 2 - \ln 1$$
$$= \frac{\ln 2}{\left| \text{ASource Paper} \right|}$$

2 It is given that
$$\tan \frac{1}{2} y = \sqrt{2}x$$
, where $-\frac{\pi}{2} < y < \frac{\pi}{2}$.

(i) Show that
$$(1+2x^2)\frac{dy}{dx} = 2\sqrt{2}$$
 . [2]

Solution

 $\tan\frac{1}{2}y = \sqrt{2}x \qquad \qquad \text{---}(1)$

Differentiate w.r.t x, $\frac{1}{2}\sec^2\left(\frac{1}{2}y\right)\frac{dy}{dx} = \sqrt{2}$

- $\left[1 + \tan^2\left(\frac{1}{2}y\right)\right]\frac{dy}{dx} = 2\sqrt{2}$ $\left(1 + 2x^2\right)\frac{dy}{dx} = 2\sqrt{2} \qquad \qquad \text{---}(2)$
 - (ii) Use the result from part (i) to find the first two non-zero terms in the Maclaurin series for *y*, giving the coefficients in exact form. [3]

Solution

Differentiate Eq(2) w.r.t x,
$$(1+2x^2)\frac{d^2y}{dx^2} + 4x\frac{dy}{dx} = 0$$
 ---(3)
 $(1+2x^2)\frac{d^3y}{dx^3} + 4x\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4x\frac{d^2y}{dx^2} = 0$ ---(4)

When x = 0, from (1), (2), (3) & (4),

 $y = 0, \frac{dy}{dx} = 2\sqrt{2}, \frac{d^2y}{dx^2} = 0, \frac{d^3y}{dx^3} = -8\sqrt{2}$ $\therefore \quad y = 0 + 2\sqrt{2}x + 0x^2 + \frac{-8\sqrt{2}}{3!}x^3 + \dots$ $= 2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots$ $I = 2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots$ (iii) Hence, using standards series from the List of Formulae (MF26), find the expansion of $\frac{2 \tan^{-1} \sqrt{2x}}{\cos 2x}$ in ascending powers of x, up to and including the term in x^3 , giving the coefficients in exact form. [3]

$$\frac{2\tan^{-1}\sqrt{2}x}{\cos 2x} = \frac{y}{\cos 2x}$$

$$= \left(2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots\right) \left[1 - \frac{1}{2!}(2x)^2 + \frac{1}{4!}(2x)^4 + \dots\right]^{-1}$$

$$= \left(2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots\right) \left[1 - 2x^2 + \frac{2}{3}x^4 + \dots\right]^{-1}$$

$$= \left(2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots\right) \left[1 - \left(-2x^2 + \frac{2}{3}x^4 + \dots\right)\right]$$

$$= \left(2\sqrt{2}x - \frac{4\sqrt{2}}{3}x^3 + \dots\right) \left(1 + 2x^2 + \dots\right)$$

$$= 2\sqrt{2}x + 4\sqrt{2}x^3 - \frac{4\sqrt{2}}{3}x^3 + \dots$$

$$= 2\sqrt{2}x + \frac{8\sqrt{2}}{3}x^3 \text{ (up to the } x^3 \text{ term)}$$



3 The position vectors of A, B and C referred to a point O are **a**, **b** and **c** respectively. The point N is on AB such that AN:NB = 2:1.

(i) If O is the midpoint of CN, prove that $\mathbf{a} + 2\mathbf{b} + 3\mathbf{c} = \mathbf{0}$. [2]

Solution

By ratio theorem,
$$\overrightarrow{ON} = \frac{1}{3}(\mathbf{a} + 2\mathbf{b})$$

Since *O* is the midpoint of *CN*, $\overline{ON} = -\mathbf{c}$

$$\therefore \overrightarrow{ON} = \frac{1}{3} (\mathbf{a} + 2\mathbf{b}) = -\mathbf{c}$$
$$\Rightarrow \mathbf{a} + 2\mathbf{b} = -3\mathbf{c}$$
$$\Rightarrow \mathbf{a} + 2\mathbf{b} + 3\mathbf{c} = \mathbf{0}$$



The point *M* is on *CB* such that CM:MB = 2:3.

(ii) Show that *A*, *O* and *M* are collinear, and find the ratio *AO*:*OM* [3] <u>Solution</u>

By the ratio theorem, $\overrightarrow{OM} = \frac{2\mathbf{b} + 3\mathbf{c}}{2+3}$

$$=\frac{-\mathbf{a}}{5}=-\frac{1}{5}\overrightarrow{OA}$$

Hence, A, O and M are collinear (Shown)

Since $\left| \overrightarrow{OM} \right| = \frac{1}{5} \left| \overrightarrow{OA} \right|$, the ratio AO:OM = 5:1



(iii) If the point *P* is such that $\overline{NP} = \overline{AM}$, show that the ratio of the area of *PNAM* : area of *PNOM* = 12:7 [3]

Solution

Method 1

Area of //gram PNAM

 $=h\left|\overrightarrow{AM}\right|$ where *h* is the height of the //gram





 \therefore the ratio of the area of *PNAM* : area of *PNOM* = 12:7

Method 2

Area of //gram PNAM

$$= \left| \overrightarrow{PN} \times \overrightarrow{PM} \right|$$
$$= \left| \frac{6}{5} \underbrace{a}_{\infty} \times \left(\frac{2}{3} \underbrace{a}_{\infty} - \frac{2}{3} \underbrace{b}_{\infty} \right) \right| = \frac{4}{5} \left| \underbrace{a}_{\infty} \times \underbrace{b}_{\infty} \right|$$

Area of trapezium PNOM

$$= \left| \overrightarrow{PN} \times \overrightarrow{PM} \right| - \frac{1}{2} \left| \overrightarrow{ON} \times \overrightarrow{OA} \right|$$
$$= \frac{4}{5} \left| \overrightarrow{a} \times \cancel{b} \right| - \frac{1}{2} \left| \left(\frac{1}{3} \overrightarrow{a} + \frac{2}{3} \cancel{b} \right) \times \overrightarrow{a} \right|$$
$$= \frac{4}{5} \left| \overrightarrow{a} \times \cancel{b} \right| - \frac{1}{2} \left| -\frac{2}{3} \overrightarrow{a} \times \cancel{b} \right| = \frac{7}{12} \left| \overrightarrow{a} \times \cancel{b} \right|$$

 \therefore the ratio of the area of *PNAM* : area of *PNOM* = 12:7



4 (a)



The diagram shows the graph of y = f(x). The curve crosses the x-axis at x = 0 and x = 3. It has a turning point at (4, 5) and asymptotes with equations y = 2 and x = 2.

Showing clearly the coordinates of turning points, axial intercepts and equations of asymptotes where possible, sketch the graphs of

(i)
$$y = f(|x|);$$
 [2]

<u>Solution</u>



(ii)
$$y = f'(x)$$
 [3]

Solution



(b) The curve with equation y = f(x) is transformed by a stretch with scale factor 2 parallel to the *x*-axis, followed by a translation of 2 units in the negative *x*-direction, followed by a translation of 3 units in the positive *y*-direction. The equation of the resulting curve is $y = \ln(e^3x)$.

Find the equation of the curve y = f(x).

Solution

Translation of 3 units in the negative y-direction

$$y = \ln\left(e^{3}x\right) \xrightarrow{\text{replace } y \text{ with } y+3} y = \ln\left(e^{3}x\right) - 3 = 3 + \ln x - 3 = \ln x$$

Translation of 2 units in the positive x-direction

$$y = \ln x \xrightarrow{\text{replace } x \text{ with } x - 2} y = \ln (x - 2)$$

Stretch with scale factor $\frac{1}{2}$ parallel to the x – axis $y = \ln(x-2)$ Islandwide Delivery | Whatsapp Only 88660031 A curve C has parametric equations

$$x = a\sin^2 t, \qquad y = a\cos t,$$

where $0 \le t \le \frac{\pi}{2}$ and a > 0.

(i) Find the cartesian equation of C, stating clearly any restrictions on the values of x and y. [2]

Solution

5

$$\sin^{2} t + \cos^{2} t = 1 \qquad \Rightarrow \qquad \frac{x}{a} + \left(\frac{y}{a}\right)^{2} = 1$$
$$ax + y^{2} = a^{2} \text{ where } 0 \le x \le a \text{ and } 0 \le y \le a$$

(ii) Sketch *C*, showing clearly the axial intercepts. [1]



(iii) The region bounded by *C*, the line $y + x = \frac{5}{4}a$ and the *y*-axis is rotated through 2π radians about the *y*-axis. Show that the exact volume of the solid obtained is $k\pi a^3$ where *k* is a constant to be determined. [5]

$$y + x = \frac{5}{4}a \implies x = \frac{5}{4}a - y$$

$$\therefore a\left(\frac{5}{4}a - y\right) + y^{2} = a^{2}$$

$$y^{2} - ay + \frac{1}{4}a^{2} = 0$$

$$\left(y - \frac{1}{2}a\right)^{2} = 0$$

$$\therefore y = \frac{1}{2}a$$

$$\Rightarrow \text{Volume required} = \frac{1}{3}\pi \left(\frac{5}{4}a - \frac{1}{2}a\right)^{2} \left(\frac{5}{4}a - \frac{1}{2}a\right) - \pi \int_{\frac{1}{2}a}^{a} \left(\frac{a^{2} - y^{2}}{a}\right)^{2} dy$$

$$= \frac{1}{3}\pi \left(\frac{3}{4}a\right)^{3} - \frac{\pi}{a^{2}} \int_{\frac{1}{2}a}^{a} a^{4} - 2a^{2}y^{2} + y^{4} dy$$

$$= \frac{9}{64}\pi a^{3} - \frac{\pi}{a^{2}} \left[a^{4}y - \frac{2}{3}a^{2}y^{3} + \frac{1}{5}y^{5}\right]_{\frac{1}{2}a}^{a}$$

$$= \frac{9}{64}\pi a^{3} - \frac{\pi}{a^{2}} \left[a^{5} - \frac{2}{3}a^{5} + \frac{1}{5}a^{5} - \frac{1}{12}a^{5} - \frac{1}{160}a^{5}\right]$$

$$=\frac{29}{960}\pi a^3$$
 units³; $k=\frac{29}{960}$ (Shown)



A function f is defined by

Sketch the graph of f.

6

(i)

<u>Solution</u>

(i)

$$f(x) = \begin{cases} \frac{6}{x-4} & \text{for } x < -2, \\ -|1+x| & \text{for } -2 \le x \le 1, \\ x^2 - x - 2 & \text{for } x > 1. \end{cases}$$

(i) Evaluate exactly
$$\int_{-3}^{4} |f(x)| dx$$
. [4]

$$\int_{-3}^{4} \left| f(x) \right| dx = \int_{-3}^{-2} \frac{-6}{x-4} dx + \frac{1}{2} \left(1 \right) \left(1 \right) + \frac{1}{2} \left(2 \right) \left(2 \right) - \int_{1}^{2} x^{2} - x - 2 dx + \int_{2}^{4} x^{2} - x - 2 dx$$

$$= -6 \left[\ln \left| x - 4 \right| \right]_{-3}^{-2} + \frac{5}{2} - \left[\frac{1}{3} x^3 - \frac{1}{2} x^2 - 2x \right]_{1}^{2} + \left[\frac{1}{3} x^3 - \frac{1}{2} x^2 - 2x \right]_{2}^{4}$$

Islandwide Delivery | Whatsapp Only BB660031

$$= -6 \ln \frac{6}{7} + \frac{5}{2} - \left[\frac{8}{3} - \frac{4}{2} - 4 - \frac{1}{3} + \frac{1}{2} + 2 \right] + \left[\frac{64}{3} - \frac{16}{2} - 8 - \frac{8}{3} + \frac{4}{2} + 4 \right]$$

$$= -6 \ln \frac{6}{7} + \frac{37}{3}$$

[3]

12

7 Do not use a calculator in answering this question.

The roots of the equation $z^2 + (2-2i)z = -3 - 2i$ are z_1 and z_2 .

(i) Find
$$z_1$$
 and z_2 in cartesian form $x + iy$, showing clearly your working. [5]

Solution

(i) Let
$$z = x + iy$$

 $\therefore (x + iy)^2 + (2 - 2i)(x + iy) = -3 - 2i$
 $x^2 + 2xyi - y^2 + 2x + 2yi - 2xi + 2y = -3 - 2i$
 $(x^2 + 2x - y^2 + 2y) + i(2xy + 2y - 2x) = -3 - 2i$
By comparing real and imaginary parts,
 $x^2 + 2x - y^2 + 2y = -3$...(1)

$$2xy + 2y - 2x = -2$$

y(x + 1) = x - 1
$$y = \frac{x - 1}{x + 1}$$
...(2)

Subt (2) into (1):

$$x^{2} + 2x - \left(\frac{x-1}{x+1}\right)^{2} + 2\left(\frac{x-1}{x+1}\right) = -3$$

$$x^{2} (x+1)^{2} + 2x (x+1)^{2} - (x-1)^{2} + 2(x-1)(x+1) = -3(x+1)^{2}$$

$$x^{2} (x^{2} + 2x+1) + 2x (x^{2} + 2x+1) - (x^{2} - 2x+1) + 2(x^{2} - 1) = -3(x^{2} + 2x+1)$$

$$x^{4} + 4x^{3} + 9x^{2} + 10x = 0$$
Let $f(x) = x(x^{3} + 4x^{2} + 9x + 10)$

$$f(0) = 0 \text{ and } f(-2) = -8 + 16 - 18 + 10 = 0$$

$$\Rightarrow x = 0, -2 \text{ are roots of } x^{4} + 4x^{3} + 9x^{2} + 10x = 0$$
When $x = 0, y = -1$ and when $x = -2, y = 3$

$$\therefore z_{1} = -i \text{ and } z_{2} = 2 + 3i$$

(ii) The complex numbers z_1 and z_2 are also roots of the equation

 $z^4 + 4z^3 + 14z^2 + 4z + 13 = 0.$

Find the other roots of the equation, explaining clearly how the answers are obtained. [2]

Solution

Since all the coefficients of $z^4 + 4z^3 + 14z^2 + 4z + 13 = 0$ are real, the other roots of the equation must be complex conjugates of -i and -2 + 3i.

Hence the other two roots are i and -2 - 3i.

(iii) Using your answer in part (i), solve
$$z^2 + (2+2i)z = 3+2i$$
. [2]

$$z^{2} + (2 - 2i)z = -3 - 2i$$

-z² + (-2 + 2i)z = 3 + 2i
-z² + (2i² + 2i)z = 3 + 2i
(iz)² + (2i + 2)iz = 3 + 2i
Hence from part (i), -iz = -i or -2 + 3i

$$\Rightarrow$$
 z = 1 or $\frac{-2+3i}{-i} \times \frac{i}{i} = -3-2i$



8 (i) Show that
$$\frac{3r+4}{(r+2)(r+1)r} = \frac{A}{r+2} + \frac{B}{r+1} + \frac{C}{r}$$
 where A, B and C are constants to be determined. [1]

Solution

$$\frac{3r+4}{(r+2)(r+1)r} = \frac{A}{r+2} + \frac{B}{r+1} + \frac{C}{r}$$

By cover-up rule, $A = \frac{-2}{(-1)(-2)} = -1, B = \frac{1}{(1)(-1)} = -1, C = \frac{4}{(2)(1)} = 2$

(ii) Find the sum to *n* terms of

$$\frac{7}{3\times2\times1} + \frac{10}{4\times3\times2} + \frac{13}{5\times4\times3} + \dots$$

(There is no need to express your answer as a single algebraic fraction). [4]

The sum to *n*th terms
$$= \sum_{r=1}^{n} \frac{3r+4}{(r+2)(r+1)r}$$
$$= \sum_{r=1}^{n} \left[-\frac{1}{r+2} - \frac{1}{r+1} + \frac{2}{r} \right]$$
$$= \left[-\frac{1}{3} - \frac{1}{2} + 2 - \frac{1}{4} - \frac{1}{3} + 1 - \frac{1}{5} - \frac{1}{4} + \frac{2}{3} - \frac{1}{5} - \frac{1}{4} + \frac{2}{3} - \frac{1}{7} + \frac{1}{7} - \frac{1}{7}$$

(iii) Hence show that
$$\sum_{r=5}^{n+3} \frac{3r-5}{(r-1)(r-2)(r-3)} < \frac{4}{3}$$
. [3]

$$\sum_{r=5}^{n+3} \frac{3r-5}{(r-1)(r-2)(r-3)}$$

$$= \sum_{r+3=5}^{r+3=n+3} \frac{3(r+3)-5}{(r+3-1)(r+3-2)(r+3-3)}$$

$$= \sum_{r=2}^{r=n} \frac{3r+4}{(r+2)(r+1)(r)}$$

$$= \sum_{r=1}^{n} \frac{3r+4}{(r+2)(r+1)(r)} - \frac{3(1)+4}{(3)(2)(1)}$$

$$= \left[\frac{5}{2} - \frac{1}{n+2} - \frac{2}{n+1}\right] - \frac{7}{6}$$

$$= \frac{4}{3} - \left[\frac{1}{n+2} + \frac{2}{n+1}\right] < \frac{4}{3} \text{ since } \frac{1}{n+2} + \frac{2}{n+1} > 0 \text{ for all } n.$$





The diagram above shows an object with O at the centre of its rectangular base ABCD where AB = 8 cm and BC = 4 cm. The top side of the object, EFGH is a square with side 2 cm long and is parallel to the base. The centre of the top side is vertically above O at a height of h cm.

(i) Show that the equation of the line *BG* may be expressed as
$$\mathbf{r} = \begin{pmatrix} 4 \\ -2 \\ 0 \end{pmatrix} + t \begin{pmatrix} -3 \\ 1 \\ h \end{pmatrix}$$
,
where *t* is a parameter. [1]

Solution

$$B(4,-2,0) \text{ and } G(1,-1,h)$$
$$\overrightarrow{BG} = \begin{pmatrix} 1\\-1\\h \end{pmatrix} - \begin{pmatrix} 4\\-2\\0 \end{pmatrix} = \begin{pmatrix} -3\\1\\h \end{pmatrix}$$
$$l_{BG} : \overrightarrow{r} = \begin{pmatrix} 4\\-2\\0 \end{pmatrix} + t \begin{pmatrix} -3\\1\\h \end{pmatrix}, t \in \mathbb{R}$$



www.KiasuExamPaper.com 73 (ii) Find the sine of the angle between the line BG and the rectangular base ABCD in terms of *h*. [2]

Solution

Let θ be the angle between the line *BG* and the rectangular base *ABCD*.

$$\sin\theta = \frac{1}{\sqrt{10 + h^2}} \begin{pmatrix} -3\\1\\h \end{pmatrix} \cdot \begin{pmatrix} 0\\0\\1 \end{pmatrix}$$

 $\sin\theta = \frac{h}{\sqrt{10 + h^2}}$

It is given that h = 6.

/

(iii) Find the cartesian equation of the plane BCFG. [3]

Solution

A normal perpendicular to plane BCFG is

$$\underline{n} = \overrightarrow{BG} \times \overrightarrow{GF} = \begin{pmatrix} -3\\1\\6 \end{pmatrix} \times \begin{pmatrix} 0\\2\\0 \end{pmatrix} = \begin{pmatrix} -12\\0\\-6 \end{pmatrix} = -6 \begin{pmatrix} 2\\0\\1 \end{pmatrix}$$

Equation of plane *BCFG* is $\underline{r} \cdot \begin{pmatrix} 2\\0\\1 \end{pmatrix} = \begin{pmatrix} 2\\0\\1 \end{pmatrix} \cdot \begin{pmatrix} 4\\-2\\0 \end{pmatrix}$
 $\underline{r} \cdot \begin{pmatrix} 2\\0\\1 \end{pmatrix} = 8$

Cartesian equation of plane is 2x + z - 8 = 0

Find the shortest distance from the point *A* to the plane *BCFG*. (iv)

[2]

Shortest distance from point A to plane
$$BCFG = \frac{|\underline{a} \cdot \underline{n} - d|}{|\underline{n}|}$$

Islandwide Delivery | Whatsapp Only 88660031

$$= \frac{1}{\sqrt{5}} \begin{vmatrix} -4 \\ -2 \\ 0 \end{vmatrix} \cdot \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} - 8 = \frac{16}{\sqrt{5}} = \frac{16\sqrt{5}}{5} \text{ units}$$

(v) The line *l*, which passes through the point *A*, is parallel to the normal of plane *BCFG*. Given that, the line *l* intersects the plane *BCFG* at a point *M*, use your answer in part (iv) to find the shortest distance from point *M* to the rectangular base *ABCD*.

Solution

$$\overline{AM} = \frac{16}{\sqrt{5}}\,\hat{n} = \frac{16}{\sqrt{5}} \left(\frac{1}{\sqrt{5}} \begin{pmatrix} 2\\0\\1 \end{pmatrix} \right) = \frac{16}{5} \begin{pmatrix} 2\\0\\1 \end{pmatrix}$$

Shortest distance from *M* to the base = z-coordinate of \overrightarrow{AM}

$$=\frac{16}{5}$$



10 Commonly used in building materials, sand is the second largest world resource used by humans after water. To reduce the environmental impacts of sand mining, an alternative approach is to make "sand" by crushing rock.

A machine designed for this purpose produces sand in large quantities. Sand falling from the output chute of the machine forms a pile in the shape of a right circular cone such that the height of the cone is always equal to $\frac{4}{3}$ of the radius of its base.

[Volume of a cone = $\frac{1}{3}\pi r^2 h$ and curved surface area of a cone = $\pi r l$ where *r* is the radius of the base area, *h* is the height of the cone and *l* is the slant length of the cone]



A machine operator starts the machine.

(a) (i) Given that V and A denote the volume and the curved surface area of the conical pile respectively, write down V and A in terms of r, the radius of its base.

Solution

$$V = \frac{1}{3}\pi r^2 \left(\frac{4}{3}r\right) = \frac{4}{9}\pi r^3$$

 $A = \pi r l$

$$= \pi r \sqrt{r^2 + h^2}$$
$$= \pi r \sqrt{r^2 + \frac{16}{9}r^2} = \frac{5}{3}\pi r^2$$

(ii) Hence show that the rate of change of A with respect to V is inversely proportional to the radius of the conical pile. [3] Islandwide Delivery | Whatsapp Only 88660031

$$\frac{dV}{dr} = \frac{4}{9}\pi(3r^2) = \frac{4}{3}\pi r^2 \quad \text{and} \quad \frac{dA}{dr} = \frac{5}{3}\pi(2r) = \frac{10}{3}\pi r$$

$$\frac{dA}{dV} = \frac{dA}{dr} \times \frac{dr}{dV}$$
$$= \frac{10}{3}\pi r \times \frac{1}{\frac{4}{3}\pi r^2}$$
$$= \frac{5}{2r} \qquad \text{(Shown)}$$

An architect is tasked to design sand-lined walking paths in a large park. He decides to base his design of the paths on the shape of astroids, which are shapes with equations

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = k^{\frac{2}{3}} \ (k > 0).$$

On a piece of graph paper, he sketches an astroid with the equation $x^{\frac{2}{3}} + y^{\frac{2}{3}} = k^{\frac{2}{3}}$.

(b) The tangent at a point $P(x_1, y_1)$ on the curve meets the x-axis at Q and the y-axis at R. Show that the length of QR is independent of where P lies on the curve. [7]

Solution

(ii) $x^{\frac{2}{3}} + y^{\frac{2}{3}} = k^{\frac{2}{3}}$

Differentiating wrt x, $\frac{2}{3}x^{-\frac{1}{3}} + \frac{2}{3}y^{-\frac{1}{3}}\frac{dy}{dx} = 0$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\left(\frac{y}{x}\right)^{\frac{1}{3}}$$

: the equation of the tangent at $P(x_1, y_1)$ is $y - y_1 = \left(-\frac{y_1}{x_1}\right)^{\frac{1}{3}} (x - x_1)$

When
$$x$$
 Example y and y

When
$$y = 0$$
, $-y_1 = \left(-\frac{y_1}{x_1}\right)^{\frac{1}{3}} (x - x_1)$
 $x = v_1^{\frac{2}{3}} x_1^{\frac{1}{3}} + x_1$

 $x = y_1^{3} x_1^{3} + x_1$ The length of $QR = \sqrt{x^2 + y^2}$

$$= \sqrt{\left(y_{1}^{\frac{2}{3}}x_{1}^{\frac{1}{3}} + x_{1}\right)^{2} + \left(y_{1}^{\frac{1}{3}}x_{1}^{\frac{2}{3}} + y_{1}\right)^{2}}$$
$$= \sqrt{x_{1}^{\frac{2}{3}}\left(y_{1}^{\frac{2}{3}} + x_{1}^{\frac{2}{3}}\right)^{2} + y_{1}^{\frac{2}{3}}\left(x_{1}^{\frac{2}{3}} + y_{1}^{\frac{2}{3}}\right)^{2}}$$
$$= \sqrt{x_{1}^{\frac{2}{3}}\left(k^{\frac{2}{3}}\right)^{2} + y_{1}^{\frac{2}{3}}\left(k^{\frac{2}{3}}\right)^{2}}$$
$$= k^{\frac{2}{3}}\sqrt{x_{1}^{\frac{2}{3}} + y_{1}^{\frac{2}{3}}}$$
$$= k^{\frac{2}{3}}\sqrt{k^{\frac{2}{3}}}$$
$$= k \text{ which is a constant.}$$

Hence, the length of QR is independent of where P lies on the curve. (Shown)



11 (a) Find the sum of all integers between 200 and 1000 (both inclusive) that are not divisible by 7. [4]

Solution

Sum required = (200 + 201 + 202 + ... + 1000) - (203 + 210 + 217 + ... + 994)

$$= \frac{1000 - 200 + 1}{2} (200 + 1000) - 7(29 + 30 + 31 + ... + 142)$$
$$= 480600 - 7 \left(\frac{142 - 29 + 1}{2}\right) (29 + 142)$$
$$= 412371$$



Snowflakes can be constructed by starting with an equilateral triangle (Fig. 1), then repeatedly altering each line segment of the resulting polygon as follows:

- 1. Divide each outer line segments into three segments of equal length.
- 2. Add an equilateral triangle that has the middle segment from step 1 as its base.
- 3. Remove the line segment that is the base of the triangle from step 2.
- 4. Repeat the above steps for a number of iterations, *n*.



(i) If a_0 denotes the area of the original triangle and the area of each new triangle added in the n^{th} iteration is denoted by a_n , show that

$$a_n = \frac{1}{9}a_{n-1}$$
 for all positive integers *n*. [2]

Solution

Each new triangle added in the n^{th} iteration is similar to the triangle added in the previous iteration (AAA).

Since the length of a side of the new triangle is $\frac{1}{3}$ of the length of a side of the triangle in the previous iteration, $a_n = \left(\frac{1}{3}\right)^2 a_{n-1} = \frac{1}{9}a_{n-1}$ (Shown)

(ii) Write down the number of sides in the polygons in Fig.1, Fig. 2 and Fig. 3, and deduce, with clear explanations, that the number of new triangles added in the n^{th} iteration is $T_n = k(4^n)$ where k is a constant to be determined. [2]

Solution

No of sides in the polygon (Fig 1) = 3, No of sides in the polygon (Fig 2) = 12

No of sides in the polygon (Fig 3) = 48

GP with first term 3 and common ratio 4

Since there is 1 new triangle for every side in the next iteration, the number of new triangles added in the n^{th} iteration = the number of sides in the polygon after the $(n - 1)^{\text{th}}$ iteration

i.e.
$$T_1 = 3$$
, $T_2 = 12$, $T_3 = 48$, ...
 $\therefore T_n = 3(4)^{n-1} = \frac{3}{4}(4)^n$ and $k = \frac{3}{4}$

(iii) Find the total area of triangles added in the n^{th} iteration, A_n in terms of a_0 and n. [2]

iteration is
$$a_0 \left[\frac{8}{5} - \frac{3}{5} \left(\frac{4}{9} \right)^n \right]$$
 units². [3]

Solution

Total area required =
$$a_0 + \sum_{r=1}^n \frac{3}{4} \left(\frac{4}{9}\right)^r a_0$$

= $a_0 + \frac{3}{4} a_0 \frac{\frac{4}{9} \left(1 - \frac{4^n}{9}\right)}{1 - \frac{4}{9}}$
= $a_0 \left[1 + \frac{3}{5} \left(1 - \frac{4^n}{9}\right)\right]$
= $a_0 \left[\frac{8}{5} - \frac{3}{5} \left(\frac{4}{9}\right)^n\right]$ (Shown)

The snowflake formed when the above steps are followed indefinitely is called the Koch snowflake.

(v) Determine the least number of iterations needed for the area of the snowflake to exceed 99% of the area of a Koch snowflake. [3]

Solution

Area of a Koch snowflake =
$$\lim_{n \to \infty} a_0 \left[\frac{8}{5} - \frac{3}{5} \left(\frac{4}{9} \right)^n \right] = \frac{8}{5} a_0$$

Consider
$$a_0 \left[\frac{8}{5} - \frac{3}{5} \left(\frac{4}{9} \right)^n \right] > 0.99 \times \frac{8}{5} a_0$$

 $0.016 - \frac{3}{5} \left(\frac{4}{9} \right)^n > 0$

From GC,

	n
	4
	5
	6
-	5 6

Hence the least value of *n* is 5.



ANDERSON SERANGOON JUNIOR COLLEGE

MATHEMATICS

9758

3 hours

H2 Mathematics Paper 2 (100 marks)

18 September 2019

Additional Material(s): List of Formulae (MF26)

CANDIDATE NAME			 	
CLASS		/		

READ THESE INSTRUCTIONS FIRST

Write your name and class in the boxes above.

Please write clearly and use capital letters.

Write in dark blue or black pen. HB pencil may be used for graphs and diagrams only.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions and write your answers in this booklet. Do not tear out any part of this booklet.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

You are expected to use an approved graphing calculator.

Where unsupported answers from a graphing calculator are not allowed in a question, you are required to present the mathematical steps using mathematical notations and not calculator commands.

All work must be handed in at the end of the examination. If you have used any additional paper, please insert them inside this booklet.

The number of marks is given in brackets [] at the end of each question or part question.



Question number	Marks
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
Total	

This document consists of 20 printed pages and 4 blank pages.

Section A: Pure Mathematics [40 marks]

1 Given that the curve
$$y = \frac{ax^2 + bx + c}{x - 1} = f(x)$$
 passes through the points (-1, -3),
 $\left(\frac{1}{2}, -\frac{3}{2}\right)$ and $\left(5, \frac{33}{2}\right)$, find the values of a, b and c . [3]

Solution

$$\frac{a(-1)^2 - b + c}{-1 - 1} = -3 \qquad \Rightarrow \qquad a - b + c = 6 \qquad ---(1)$$

$$\frac{a\left(\frac{1}{2}\right)^2 + \frac{1}{2}b + c}{\frac{1}{2} - 1} = -\frac{3}{2} \qquad \Rightarrow \qquad \frac{1}{4}a + \frac{1}{2}b + c = \frac{3}{4} \qquad ---(2)$$

$$\frac{a(5)^2 + 5b + c}{5 - 1} = \frac{33}{2} \qquad \Rightarrow \qquad 25a + 5b + c = 66 \qquad ---(3)$$

Using GC, a = 3, b = -2, c = 1.

Hence find the exact range of values of x for which the gradient of y = f(x) is positive. [3]

$$f'(x) > 0 \qquad \Rightarrow \frac{d}{dx} \left(\frac{3x^2 - 2x + 1}{x - 1} \right) > 0$$

$$\Rightarrow \frac{(x - 1)(6x - 2) - (3x^2 - 2x + 1)(1)}{(x - 1)^2} > 0$$

$$\Rightarrow \frac{3x^2 - 6x + 1}{(x - 1)^2} > 0$$

Since $(x - 1)^2 > 0$, $3x^2 - 6x + 1 > 0$
$$= \frac{3x^2 - 6x + 1 > 0}{(\sqrt{3}x - \sqrt{3} - \sqrt{2})(\sqrt{3}x - \sqrt{3} + \sqrt{2}) > 0}$$

$$= \frac{1 - \sqrt{\frac{2}{3}}}{1 - \sqrt{\frac{2}{3}}}$$

2 The functions f and g are defined by:

$$f: x \mapsto |ax - x^2|, \qquad x \in \mathbb{R}$$

 $g: x \mapsto ae^{-x} + 1, \qquad x \ge 0$

where *a* is a positive constant and $a \ge 5$.

(i) Sketch the graphs of y = g(x) and y = f(x) on separate diagrams. Hence find the range of the composite function fg, leaving your answer in exact form in terms of *a*. [5]



From the graphs,

$$R_g = (1, a + 1]$$

Since $a \ge 5$, $\frac{a}{2} \ge \frac{5}{2} > 1 \implies 0 < 1 < \frac{a}{2} \implies \frac{a^2}{4} \ge a - 1$
and $a^2 \ge 5a \implies \frac{a^2}{4} \ge \frac{5a}{4} = a + \frac{1}{4}a \ge a + 1$

$$[0, \infty) \xrightarrow{g} (1, \alpha + 1] \xrightarrow{f} R_{fg}$$

From the graphs, $R_{fg} = \left[0, \frac{a^2}{4}\right]$



(ii) Given that the domain of f is restricted to the subset of R for which x ≥ k, find the smallest value of k, in terms of a, for which f⁻¹ exists. Hence, without finding the expression for f⁻¹(x), sketch the graphs of y = f(x) and y = f⁻¹(x) on a single diagram, stating the exact coordinates of any points of intersection with the axes. [3]

Solution

The smallest value of *k* for which f^{-1} exists = *a*.



(iii) With the restricted domain of f in part (ii), find with clear working, the set of values of x such that $f f^{-1}(x) = f^{-1} f(x)$. [2]

Solution

 $f^{-1}f(x) = x \text{ for } x \in D_{f}, \text{ i.e } x \in [a, \infty)$ $ff^{-1}(x) = x \text{ for } x \in D_{f^{-1}}, \text{ i.e } x \in [0, \infty)$ For f f⁻¹(x) = f⁻¹ f(x), [0, \omega) \cap [a, \omega) = [a, \omega) Set of values of x required = [a, \omega)



3 A student is investigating a chemical reaction between two chemicals A and B. In his experiments, he found that the reaction produces a new product C which does not react with A and B after being formed.

His experiments also suggest that the rate at which the product, C is formed is given by the differential equation

$$\frac{\mathrm{d}x}{\mathrm{d}t} = (a - x)\sqrt{b - x},$$

where a and b are the initial concentrations (gram/litre) of A and B just before the reaction started and x is the concentration (gram/litre) of C at time t (mins).

(i) If a = b = 9, solve the differential equation to show that

$$x = 9 - \frac{36}{(3t+2)^2}.$$
 [4]

(i)
$$\frac{dx}{dt} = (9-x)(9-x)^{\frac{1}{2}}$$
$$\int (9-x)^{-\frac{3}{2}} dx = \int 1 dt$$
$$-2(-1)(9-x)^{-\frac{1}{2}} = t + c$$
When $t = 0, x = 0, 2(9-0)^{-\frac{1}{2}} = 0 + c$
$$\therefore c = \frac{2}{3}$$
$$\Rightarrow \frac{2}{\sqrt{9-x}} = t + \frac{2}{3} = \frac{3t+2}{3}$$
$$\frac{4}{9-x} = \frac{(3t+2)^2}{9}$$
$$9 - x = \frac{36}{(3t+2)^2}$$
$$\therefore x = 9 \frac{36}{(3t+2)^2}$$
EVALUATE: The second seco

(ii) Sketch the graph of x and describe the behaviour of x as t increases. Find the amount of time needed for the concentration of C to reach 4.5 gram/litre. [3]



[G1 – shape with asymptote (no need for correct value of limit]

As *t* increase, *x* increases and approaches the value of 9 gram/litre.

From GC or otherwise,
$$t = 0.276$$
 mins or 16.6 s (3sf) or $\frac{2(\sqrt{2}-1)}{3}$

It is now given that a > b instead.

(iii) Solve the differential equation to find t in terms of x, a and b by using the substitution $u = \sqrt{b-x}$ for $0 \le x < b$. [6]

Solution

$$\frac{du}{dx} = \frac{-1}{2\sqrt{b-x}} = -\frac{1}{2u}$$

$$u = \sqrt{b-x} \implies x = b - u^2, \quad \therefore \quad \frac{du}{dt} \times \frac{dx}{du} = (a - b + u^2)u$$

$$\frac{du}{dt} \times (-2u) = (a - b + u^2)u$$

$$-2\int \frac{1}{(a - b) + u^2} du = \int 1 dt$$

$$-\frac{2}{\sqrt{a-b}} \tan^{-1} \frac{u}{\sqrt{a-b}} + C = t$$
When $t = 0, x = 0$, $u = \sqrt{b}$, $\therefore 0 = \frac{2}{\sqrt{a-b}} \tan^{-1} \sqrt{\frac{b}{a-b}}$

$$\therefore C = \frac{2}{\sqrt{a-b}} \tan^{-1} \sqrt{\frac{b}{a-b}} - \frac{2}{\sqrt{a-b}} \tan^{-1} \sqrt{\frac{b-x}{a-b}}$$

$$\Rightarrow t = \frac{2}{\sqrt{a-b}} \tan^{-1} \sqrt{\frac{b}{a-b}} - \frac{2}{\sqrt{a-b}} \tan^{-1} \sqrt{\frac{b-x}{a-b}}$$

www.KiasuExamPaper.com 87

4 The illumination of an object by a light source is directly proportional to the strength of the source and inversely proportional to the square of the distance from the source.

A small object, P, is placed on the straight line passing through two light sources A and B that are eight metres apart. It is known that the light sources are similar except that S, the strength of light source A, is a positive constant and is three times that of light source B. It can be assumed that I, the total illumination of the object P by the two light source is the sum of the illumination due to each light source.

The distance between the object P and light source A is denoted by x.

Show that when *P* is between *A* and *B*, **(i)**

$$I = kS \left[\frac{1}{x^2} + \frac{1}{\alpha (\beta - x)^2} \right]$$
 where k is a positive constant and,
nts to be determined. [1]

 α and β are constants to be determined.

Solution

$$I = \frac{kS}{x^2} + \frac{k\left(\frac{1}{3}S\right)}{\left(8-x\right)^2} \text{ where } k > 0 \qquad A \qquad P \qquad B$$
$$= kS\left[\frac{1}{x^2} + \frac{1}{3\left(8-x\right)^2}\right] \text{ so } \alpha = 3 \text{ and } \beta = 8$$

(ii) Show that there is only one value of x that will give a stationary value of I. Determine this value and the nature of the stationary value without the use of a graphic calculator. [5]

$$\frac{dI}{dx} = kS \left[-\frac{2}{x^3} - 2(-1)\frac{1}{3(8-x)^3} \right] = 2kS \left[-\frac{1}{x^3} + \frac{1}{3(8-x)^3} \right]$$
Let $\frac{dI}{dx} = 0 \implies -\frac{1}{x^3} + \frac{1}{3(8-x)^3} = 0$
 $x^3 = 3(8-x)^3$
 $x^3 = 3(8-x)^3$
Listandwide $x^3 = 3\sqrt[3]{8-x}$
 $x^3 =$

Hence *I* will have a stationary value only when $x = \frac{8\sqrt[3]{3}}{1+\sqrt[3]{3}}$. (Shown)

$$\frac{d^2 I}{dx^2} = 2kS\left[\frac{3}{x^4} + \frac{(-3)(-1)}{3(8-x)^4}\right]$$
$$= 2kS\left[\frac{3}{x^4} + \frac{1}{(8-x)^4}\right] > 0 \text{ for all } x.$$

 $\Rightarrow I \text{ is minimum when } x = \frac{8\sqrt[3]{3}}{1+\sqrt[3]{3}}.$

It is now given that
$$I = \frac{4x^2 - 48x + 192}{x^2 (x^2 - 16x + 64)}$$
 for $0 < x < 8$.

(iii) Sketch the graph of *I*, indicating clearly any equations of asymptote and coordinates of the stationary point. [3]

Solution



(iv) It is desired to place P such that $I = \sqrt{\lambda - (x - x_1)^2}$ where x_1 is the value of x found in part (ii) and λ is a positive constant. Find the value of λ if there is only one possible position to place P. [2]

Solution

$$I = \sqrt{\lambda - (x - x_1)^2}$$
 is the equation of a semi-circle centred at $(x_1, 0)$ and radius $\sqrt{\lambda}$.
Example 7
Hence when there is only 1 position to place *P*, it must happen when $\sqrt{\lambda} = 0.2276094$

i.e. $\lambda = 0.0518$ (3sf)

Section B: Statistics [60 marks]

5 Greenhouse gases generated by Singapore comes mainly from the burning of fossil fuels to generate energy for industries, buildings, households and transportation.

The following table* shows the total greenhouse gas emissions, X (kilotonnes of CO₂), and the total deaths in Singapore, Y from 2007 to 2016.

Year		2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Total 42613		41606	43100	47063	49930	48712	50299	50908	51896	51519	
Greenhouse											
Gas Emissions											
(X)											
Total	Deaths	17140	17222	17101	17610	18027	18481	18938	19393	19862	20017
(Y)											

*Extracted from www.singstat.gov.sg

(i) Draw a scatter diagram for these values, labelling the axes and state the product moment correlation between *X* and *Y*. Use your diagram to explain why the best model for the relationship between *X* and *Y* may not be given by Y = aX + b, where *a* and *b* are constants. [3]

Solution



From the GC, r = 0.90527 = 0.905 (3 sf)

From the scatter diagram, the points follow a curvilinear trend rather than a linear trend, hence Y = aX + b may not be the best model.

(ii) Comment on the correctness of the following statement.

"Since there exists a high positive linear correlation between total greenhouse gas emission and total deaths, the deaths are a result of the greenhouse gas emissions." [1]

Solution

High positive linear correlation does not indicate causation, hence it is incorrect to state that the deaths are a result of the greenhouse gas emission. Moreover, there are other factors affecting the total deaths in Singapore.

(iii) Explain clearly which of the following equations, where a and b are constants, provides the more accurate model of the relationship between X and Y.

(A)
$$Y = aX^4 + b$$

(B) $\lg Y = aX + b$ [3]

Solution

From GC, r = 0.928 for $Y = aX^4 + b$ r = 0.912 for $\lg Y = aX + b$

Since the product moment correlation coefficient for $Y = aX^4 + b$ is closer to 1 as compared to the one for $\lg Y = aX + b$, $Y = aX^4 + b$ is the better model.

The Government has pledged that Singapore's greenhouse gas emissions will peak around 2030 at the equivalent of 65 million tonnes of carbon dioxide.

(iv) Using the model you have chosen in part (iii), write down the equation for the relationship, giving the regression coefficients correct to 5 significant figures. Hence find the estimated total deaths in the year 2030 if Singapore's pledge is fulfilled and comment on the reliability of the estimate. [3]

Solution

The required equation is $Y = 6.4326 \times 10^{-16} X^4 + 14911$

When
$$X = 65000$$
, $Y = 6.4326 \times 10^{-16} (65000)^4 + 14911$

= 26393.6

KIASE 26400 (Asf)

The estimate is not reliable as extrapolation is carried out where the linear correlation outside the range of X, i.e. [41606, 51896] may not be valid.

6 Birth weight can be used to predict short and long-term health complications for babies. Studies show that the birth weight of babies born to mothers who do not smoke in a certain hospital can be assumed to follow a normal distribution with mean 3.05 kg and variance $\sigma^2 \text{ kg}^2$.

Birth weight	Classification
Less than 1.5 kg	Very low birth weight
1.5 kg to 2.5 kg	Low birth weight
2.5 kg to 4.0 kg	Normal birth weight
More than 4.0 kg	High birth weight

The hospital classifies babies based on their birth weight as shown in the table below.

(i) A sample showed that 20.2% of the babies born to mothers who do not smoke have low birth weight. If this is true for the entire population, find two possible values of σ , corrected to 2 decimal places. Explain clearly why one of the values of σ found should be rejected. [4]

Solution

Let X be random variable representing the birth weight of a baby in kg.

:. $X \sim N(3.05, \sigma^2)$ P(1.5 < X < 2.5) = 0.202 By plotting the graph of y = P(1.5 < X < 2.5) - 0.202using GC, we have $\sigma = 0.70$ (2 dp) or $\sigma = 1.56$ (2 dp)

If $\sigma = 0.70$, P(2.5 < X < 4) = 0.70 If $\sigma = 1.56$, P(2.5 < X < 4) = 0.37



Reject $\sigma = 1.56$ as this would mean that it is very much more likely that a baby will be born with "abnormal" birth weight than normal birth weight.

(Or when $\sigma = 1.56$, P(X < 0) = 0.03 compared with P(X < 0) = 6.6 \times 10^{-6} for $\sigma = 0.7$)

Studies also show that babies born to mothers who smoke have a lower mean birth weight of 2.80 kg.

For the remaining of the question, you may assume the birth weight of babies born to mothers who smoke is also normally distributed and that the standard deviations for the birth weight of babies born to mothers who do not smoke and mothers who smoke are both 0.86 kg.

(ii) Three pregnant mothers-to-be who smoke are randomly chosen. Find the probability that all their babies will not be of normal birth weight. State an assumption that is needed for your working. [3]

Solution

Let *Y* be r.v. denoting the birth weight of a baby in kg from a mother who smokes. $Y \sim N(2.8, 0.86^2)$

Required probability =
$$[1 - P(2.5 < Y < 4)]^3$$

= $(1 - 0.55493)^3$
= 0.088157
= 0.0882 (3 sf)

Assumption: The birth weights of babies are independent of one another.

(iii) Find the probability that the average birth weight of two babies from mothers who do not smoke differs from twice the birth weight of a baby from a mother who smokes by less than 2 kg.[3]

Solution

Let
$$A = \frac{X_1 + X_2}{2} - 2Y$$

 $E(A) = 3.05 - 2(2.80) = -2.55$
 $Var(A) = 0.5^2 (0.86^2 \times 2) + 2^2 (0.86^2) = 3.3282$
 $\therefore A \sim N(-2.55, 3.3282)$
Required probability = $P(|A| < 2) = P(-2 < A < 2) = 0.37521$
 $= 0.375 (3 \text{ s.f.})$

Babies whose birth weight not classified as normal will have to remain in hospital for further observation until their condition stabilises. Depending on the treatment received and length of stay, the mean hospitalisation cost per baby is \$2800 and standard deviation is \$500.

(iv) Find the probability that the average hospitalisation cost from a random sample of 50 babies, whose birth weight not classified as normal, exceeds \$3000. [2]

Solution

Let W be random variable representing the hospitalisation charge of a baby. $E(W) = 2800 \text{ and } Var(W) = 500^{2} = 250000$ Since n = 50 is large, by Central Limit Theorem, $\overline{W} = \frac{W_{1} + W_{2} + \ldots + W_{50}}{50} \text{ N} 2800, \frac{500^{2}}{50} \text{ approximately}$ i.e. $\overline{W} \sim N(2800, 5000)$ approximately = N(2800, 5000) approximately[M1] $\Rightarrow P(\overline{W} > 3000) = 0.00234 \text{ (3.s.f)}$ [A1] 7 Three cards are to be drawn at random without replacement from a pack of six cards numbered 0, 0, 1, 1, 1, 1.

13

The random variables X_1 , X_2 and X_3 denotes the numbers on the first, second and third card respectively.

(i) If
$$Y = X_1 - X_2 + X_3$$
, show that $P(Y = 1) = \frac{1}{3}$ and find the probability distribution of *Y*. [3]

Solution

$$P(Y=1) = P(0, 0, 1) + P(1, 0, 0) + P(1, 1, 1)$$
$$= \frac{2}{6} \times \frac{1}{5} \times \frac{4}{4} + \frac{4}{6} \times \frac{2}{5} \times \frac{1}{4} + \frac{4}{6} \times \frac{3}{5} \times \frac{2}{4}$$
$$= \frac{1}{3} \quad (Shown)$$

У	-1	0	1	2
P(Y=y)	$P(0, 1, 0) = \frac{2}{6} \times \frac{4}{5} \times \frac{1}{4}$ $= \frac{1}{15}$	P(1, 1, 0) + P(0, 1, 1) = $\frac{4}{6} \times \frac{3}{5} \times \frac{2}{4} + \frac{2}{6} \times \frac{4}{5} \times \frac{3}{4}$ = $\frac{2}{5}$	$\frac{1}{3}$	$P(1, 0, 1) = \frac{4}{6} \times \frac{2}{5} \times \frac{3}{4}$ $= \frac{1}{5}$

(ii) Find the values of E(Y) and Var(Y).

Solution

$$E(Y) = -\frac{1}{15} + 0 + \frac{1}{3} + \frac{2}{5} = \frac{2}{3} = 0.667 \text{ (3sf)}$$
$$Var(Y) = \frac{1}{15} + 0 + \frac{1}{3} + \frac{4}{5} - \left(\frac{2}{3}\right)^2 = \frac{34}{45} = 0.756 \text{ (3sf)}$$

(iii) Find the probability that the difference between
$$Y$$
 and $E(Y)$ is less than one standard deviation of Y . [2]

Solution

$$P\left(\left|Y - \frac{2}{3}\right| < \sqrt{\frac{34}{45}} = P(Y = 0) + P(Y = 0)$$

= $\frac{2}{5} + \frac{1}{3} = \frac{11}{15} = 0.733 (3 \text{ sf})$

[2]

- 8
- A factory makes a certain type of muesli bar which are packed in boxes of 20. The factory claims that each muesli bar weigh at least 200 grams. Muesli bars that weigh less than 200 grams are sub-standard. It is known that the probability that a muesli bar is sub-standard is 0.02 and the weight of a muesli bar is independent of other muesli bars.
 - (i) Find the probability that, in a randomly chosen box of muesli bars, there are at least one but no more than five sub-standard muesli bars. [2]

Solution

Let *X* be the random variable denoting the number of sub-standard muesli bars, out of 20. \therefore *X* ~ B(20, 0.02)

 \Rightarrow Probability required = P(1 \le X \le 5)

$$= P(X \le 5) - P(X = 0)$$

= 0.33239
= 0.332 (3 sf)

The boxes of muesli bars are sold in cartons of 12 boxes each.

(ii) Find the probability that, in a randomly chosen carton, at least 75% of the boxes will have at most one sub-standard muesli bar. [2]

Solution

Let *Y* be the random variable denoting the number of boxes of muesli bars that have at most one sub-standard muesli bars, out of 12 boxes.

$$P(X \le 1) = 0.94010$$

 $\therefore Y \sim B(12, 0.94010)$

Probability required = $P(Y \ge 0.75 \times 12)$

$$= P(Y \ge 9)$$

= 1 - P(Y \le 8)
= 0.99568
= 0.996
KIASU



As part of quality control for each batch of muesli bars produced daily, one box of muesli bars is randomly selected to be tested. If all the muesli bars in the box weigh at least 200 grams, the batch will be released for sales. If there are more than one sub-standard muesli bar, the batch will not be released. If there is exactly one sub-standard muesli bar, a second box of muesli bars will be tested. If all the muesli bars in the second box weigh at least 200 grams, the batch will be released. Otherwise, the batch will not be released.

(iii)Find the probability that the batch is released on a randomly selected day. [2]

Solution

Probability required = $P(X_1 = 0) + P(X_1 = 1) P(X_2 = 0)$ = 0.84953 = 0.850 (3 sf)

(iv) If the factory operates 365 days in a year, state the expected number of days in the year when the batch will not be released.

Solution

Expected number of days when the batch will not be released

$$= 365 - 365 \times 0.84953$$

= 54.9 (3 sf)

To entice more people to buy their muesli bars, a scratch card is inserted into each box. The probability that a scratch card will win a prize is *p*. It may be assumed that whether a scratch card will win a prize or not, is independent of the outcomes of the other scratch cards.

Jane buys a carton of muesli bars.

(v) Write in terms of *p*, the probability that Jane win 2 prizes.

Solution

Let W be the random variable denoting the number of prizes Jane wins, out of 12.



[1]

The probability that Jane win 2 prizes is more than twice the probability that she win 3 prizes.

(vi) Find in exact terms, the range of values that p can take. [2]

Solution

$$P(W = 2) > 2 P(W = 3)$$

$$66p^{2} (1-p)^{10} > 2 {\binom{12}{3}} p^{3} (1-p)^{9}$$

$$66p^{2} (1-p)^{10} - 440p^{3} (1-p)^{9} > 0$$

$$p^{2} (1-p)^{9} [3(1-p) - 20p] > 0$$

$$p^{2} (1-p)^{9} [3-23p] > 0$$
Since $0 \le p \le 1, 0$

9 In preparation for an upcoming event, a student management team is considering having meetings on this Friday, Saturday and Sunday. The probability that the meeting is conducted on Friday is $\frac{1}{6}$. On each of the other days, the probability that a meeting is conducted when a meeting has already been conducted on the previous day, is $\frac{2}{5}$ and the probability that a meeting is conducted, when a meeting has not been conducted on the previous day, is $\frac{1}{3}$.

97

Find the probability that

a meeting is conducted on Sunday, (i)



Solution

P(meeting is conducted on Sunday)

 $=\frac{1}{6}\left[\frac{2}{5}\cdot\frac{2}{5}+\frac{3}{5}\cdot\frac{1}{3}\right]+\frac{5}{6}\left[\frac{1}{3}\cdot\frac{2}{5}+\frac{2}{3}\cdot\frac{1}{3}\right]$

(ii) a meeting is not conducted on Friday given that there is a meeting on Sunday.

[3]

Solution

P(no meeting on Friday | meeting on Sunday)

 $= \frac{P(\text{no meeting on Friday and meeting on Sunday})}{P(\text{no meeting on Friday})}$ $= \frac{\frac{5}{6} \left[\frac{1}{3} \cdot \frac{2}{5} + \frac{2}{3} \cdot \frac{1}{3}\right]}{\left(\frac{481}{1350}\right)}$ $= \frac{400}{481}$ = 0.83160= 0.832 (correct to 3 s.f.)

The student management team consists of 7 girls and 3 boys. At one of the meetings, they sit at a round table with 10 chairs.

(iii) Find the probability that the girls are seated together. [2]

Solution

P(girls are seated together) =
$$\left\lfloor \frac{(4-1)!7!}{(10-1)!} \right\rfloor$$

= $\frac{1}{12}$

Two particular boys are absent from the meeting.

(iv) Find the probability that two particular girls do not sit next to each other. [3]

Probability required paper
Islandwide Delivery (Marsall)
$$\frac{1}{2!}$$

 $=\frac{7}{9}$

10 Along a 3km stretch of a road, the speed in km/h of a vehicle is a normally distributed random variable *T*. Over a long period of time, it is known that the mean speed of vehicles traveling along that stretch of the road is 90.0 km/h. To deter speeding, the traffic governing body introduced a speed monitoring camera. Subsequently, the speeds of a random sample of 60 vehicles are recorded. The results are summarised as follows.

$$\sum t = 5325, \qquad \sum \left(t - \bar{t}\right)^2 = 2000.$$

(i) Find unbiased estimates of the population mean and variance, giving your answers to 2 decimal places. [2]

Solution

Unbiased estimate of the population mean = $\bar{t} = \frac{5325}{60} = 88.75$

Unbiased estimate of the population variance = $s^2 = \frac{\sum (t - \bar{t})^2}{59} = \frac{2000}{59}$ = 33.89830508 = 33.90

(ii) Test, at the 5% significance level, whether the speed-monitoring camera is effective in deterring the speeding of vehicles on the stretch of road. [4]

Solution

Let μ denote the population mean speed of the vehicles traveling along the stretch of the road.

To test $H_0: \mu = 90.0$

Against H₁: $\mu < 90.0$

Conduct a one-tail test at 5% level of significance, i.e., $\alpha = 0.05$

Under H₀, since n = 60 and is sufficiently large, by Central Limit Theorem, $\overline{T} \sim N\left(90.0, \frac{33.89830508}{60}\right)$ approximately.

Using GC, p-value = 0.0481545117

Since p-value < 0.05, we reject H₀. There is sufficient evidence at 5% level of significance to conclude that the mean speed along the 3km stretch of road has been reduced.

ExamPaper

(iii) In another sample of size n (n > 30) that was collected independently, it is given that $\overline{t} = 89.0$. The result of the subsequent test using this information and the unbiased estimate of the population variance in **part** (i) is that the null hypothesis is not rejected. Obtain an inequality involving n, and hence find the largest possible value n can take. [4]

Solution

If the null hypothesis is not rejected, z_{calc} must lie outside the critical region.

Critical Region:
$$z \le -1.644853626$$

Test Statistics, $Z = \frac{\overline{T} - 90.0}{\sqrt{\frac{33.89830508}{n}}} \sim N(0, 1)$
 $\therefore z_{calc} = \frac{89 - 90.0}{\sqrt{\frac{33.89830508}{n}}} > -1.644853626$
 $\frac{-\sqrt{n}}{\sqrt{33.89830508}} > -1.644853626$
 $\sqrt{n} < 9.576708062$
 $n < 91.713$

Since n is an integer, the largest possible value n can take is 91.



www.KiasuExamPaper.com 101