



NEW TOWN SECONDARY SCHOOL
Preliminary Examination
Secondary 4 Express

NAME

--

CLASS

--	--	--

INDEX
NUMBER

--	--

Additional Mathematics

Paper 2

4049/02

2 August 2021

09:05 – 11:20

2 hours 15 min

READ THESE INSTRUCTIONS FIRST

Write your name, register number and class in the spaces provided above and on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.

For Examiner's Use

This document consists of **16** printed pages.

Setter: Mr Alan Cheng

*Mathematical Formulae***1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}.$$

Binomial Expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \cdots + \binom{n}{r}a^{n-r}b^r + \cdots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\cdots(n-r+1)}{r!}$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$

- 1 Express $\frac{x^2 + 2x - 19}{(x-1)(x+3)^2}$ as a sum of three partial fractions. [5]

- 2 The table below shows experimental values of two variables x and y .


x	2	4	6	8
y	8.48	5.99	4.90	4.24

It is known that x and y are connected by the equation $yx^n = k$, where k and n are constants.


- (a) Plot $\ln y$ against $\ln x$, using a scale of 4 cm for 1 unit on both axes, for the given data and draw a straight line graph on the grid on page 5. [2]
- (b) Use your graph to estimate the value of n and of k . [3]

- (c) Use your graph to **estimate** the value of x when $y = e^2$. [2]

- (d) On the same diagram, draw the straight line representing the equation $y = x^3$ and hence find the value of x for which $x^{3+n} = k$ [3]



- 3 The diagram shows part of the curve $y = \frac{9}{(3-x)^2} - 1$ which intersects the x -axis at the origin and at the point P . The normal at the point Q on the curve cuts the x -axis at R . The gradient of the curve at Q is 18.



- (a) Find the coordinates of Q .

[3]

(b) Find the coordinates of R . [3]

(c) Find the area of the shaded region. [4]



- 4 (a) The equation of a curve is $y = (b^2 - 2ac)x^2 + 4(a + c)x - 8$.
- (i) Show that the roots of the equation are real if a , b and c are real. [3]
- (ii) State the conditions that the roots are equal. [2]
- (b) The equation of a curve is $mx^2 - 4x + m - 3$, where m is a constant.
Find the range of values of m for which $mx^2 - 4x + m - 3$ is **not** negative for all real values of x . [5]

5 (a) Show that $\frac{3x}{1-3x} = \frac{1}{1-3x} - 1$ and hence find $\int \frac{3x}{1-3x} dx$. [4]

(b) Given that $y = x \ln(1-3x)$, find an expression for $\frac{dy}{dx}$. [2]

(c) Using the results from (a) and (b), find $\int_{-1}^0 \ln(1-3x) dx$. [4]

- 6 In Figure 1, $AE = 3$ cm and angle $BAE = x$. Triangle ABE is cut and rotated clockwise about E until AE rests on ED . The resulting shape is shown in Figure 2 below.



- (a) Show that the perimeter, P cm of the resulting shape $BEBADC B$ is given by $P = 14 \sin x + 8 \cos x + 2$

[4]

- (b) Express $14\sin x + 8\cos x + 2$ in the form of $R\sin(x + \alpha) + 2$, where $R > 0$ and $0^\circ \leq \alpha \leq 90^\circ$. [4]
Hence find the maximum value of $14\sin x + 8\cos x + 2$.

- (c) Find the value of x for which $P = 15$ cm. [2]

7 The equation of a curve is $y = 4x^2e^{-3x}$.

- (a) Find an expression for $\frac{dy}{dx}$ and obtain the exact value of the coordinates of the stationary points of the curve. [6]

- (b) Find an expression for $\frac{d^2y}{dx^2}$ and hence the nature of these stationary points. [5]

- 8** $D(-6, 4)$, $E(2, 4)$ and F are three points on a circle with radius 5 units.

The normal to the circle at F passes through E .

- (a) Explain why EF is a diameter of the circle. [1]

- (b) The centre of the circle is above DE . Find the equation of the circle. [4]

- (c) Find the coordinates of F . [2]

(d) Find the equation of the tangent at F . [3]

(e) Find the coordinates of the two points on the circle which has zero gradient. [2]

- 9 (a) By using the substitutions $2^x = a$ and $5^x = b$, show that the equation $\frac{4^x - 25^x}{10^x + 4^x} = \frac{1}{2}$ can be simplified to $a = 2b$, where $a \neq -b$. [4]

- (b) Solve the equation $2\log_2 x - 2\log_x 8 = -1$. [5]

- (c) Show that the equation $\log_2(2x-6) - \log_2(x-2) = 2$ has no real solutions. [3]