

## 2015 VJC JC2 Prelim Paper 1 Solutions

**Q1)** Let  $P_n$  be the statement:

$$w_n = an + (n-1), \quad n \in \mathbb{N}^+.$$

LHS of  $P_1 = w_1 = a$  (given)

RHS of  $P_1 = a(1) + (1-1) = a$

$\therefore P_1$  is true.

Assume  $P_k$  is true for some  $k \in \mathbb{N}^+$  i.e.  $w_k = ak + (k-1)$

We want to show  $P_{k+1}$  is true i.e.  $w_{k+1} = a(k+1) + k$

LHS of  $P_{k+1} = w_{k+1}$

$$\begin{aligned} &= \frac{1}{k}[(k+1)w_k + 1] \\ &= \frac{1}{k}\{(k+1)[ak + (k-1)] + 1\} \\ &= a(k+1) + \frac{(k+1)(k-1) + 1}{k} \\ &= a(k+1) + \frac{k^2}{k} \\ &= a(k+1) + k \\ &= \text{RHS of } P_{k+1} \end{aligned}$$

$\therefore P_k$  is true  $\Rightarrow P_{k+1}$  is true

Since we have shown that

(1)  $P_1$  is true and

(2)  $P_k$  is true  $\Rightarrow P_{k+1}$  is true.

$\therefore$  By mathematical induction,  $P_n$  is true for all positive integers  $n$ .

**Q2)**

Sub (1,1) and (2,2) into  $y = h(x)$ .

$$a+b+c+d=1 \quad \dots \quad (1)$$

$$8a+4b+2c+d=2 \quad \dots \quad (2)$$

Since (2,2) is also the stationary point,  $h'(2)=0$ . i.e.

$$12a+4b+c=0 \quad \dots \quad (3)$$

Using the GC,


$$a = -\frac{1}{2} - \frac{1}{4}d$$

$$b = \frac{3}{2} + \frac{5}{4}d$$

$$c = -2d$$

$$\frac{ab}{c} \leq 0$$

$$\frac{\left(-\frac{1}{2} - \frac{1}{4}d\right)\left(\frac{3}{2} + \frac{5}{4}d\right)}{-2d} \leq 0$$



$$\{d \in \mathbb{R} : d \leq -2 \text{ or } -\frac{6}{5} \leq d < 0\}$$

$$\text{Q3)} \quad y = \frac{ax^2 + bx + d}{x-2} = 2x + 3 + \frac{k}{x-2}$$

By observation,  $a = 2$

$$\Rightarrow 2x^2 + bx + d = (2x+3)(x-2) + k$$

Compare coefficients of  $x$ :  $b = 3 - 4 \Rightarrow b = -1$

$$y = \frac{2x^2 - x + d}{x-2}$$

Given:  $d < -6$

Asymptotes:  $y = 2x + 3, x = 2$


Axial intercepts: when  $x = 0, y = -\frac{d}{2}$

$$\text{When } y = 0, 2x^2 - x + d = 0$$

$$x = \frac{1 \pm \sqrt{1-4(2)(d)}}{4} = \frac{1 \pm \sqrt{1-8d}}{4}$$

The coordinates are  $\left(0, \frac{d}{2}\right), \left(\frac{1-\sqrt{1-8d}}{4}, 0\right), \left(\frac{1+\sqrt{1-8d}}{4}, 0\right)$ .

Let  $\alpha = \frac{1-\sqrt{1-8d}}{4}$  and  $\beta = \frac{1+\sqrt{1-8d}}{4}$




**Q4(i)**

Let  $y = e^x - 7x$ . So,  $\frac{dy}{dx} = e^x - 7$ .

$$\frac{dy}{dx} = 0 \Rightarrow e^x - 7 = 0$$

$$x = \ln 7$$

$$\therefore \min \lambda = \ln 7$$




**Q4(ii)** Let  $x = g^{-1}(1)$


$$\Rightarrow g(x) = 1$$

$$\Rightarrow e^x - 7x = 1$$



From the GC,

$$x = 3.13$$



**Q4(iii)**

**Q5(i)**  $y = \frac{1}{f(x)}$

**Q5(ii)**

$$\begin{aligned}
 \int_{-a}^0 [2 - f'(x)] dx &= [2x - f(x)] \Big|_{-a}^0 \\
 &= [2(0) - f(0)] - [2(-a) - f(-a)] \\
 &= -(-5a) + 2a + (-4a) \\
 &= 3a
 \end{aligned}$$

**Q6(i)**

$n$	Amount at end of year $n$
1	$1.08(1000)$
2	$1.08[1000 + 1.08(1000)] = 1000(1.08 + 1.08^2)$
3 ⋮	$1.08[1000 + 1000(1.08) + 1000(1.08)^2]$ $= 1000(1.08 + 1.08^2 + 1.08^3)$
$n$	$1000(1.08 + 1.08^2 + \dots + 1.08^n)$

Amount at the end of year 2040

$$= 1000 \left[ (1.08) + (1.08)^2 + \dots + (1.08)^{26} \right]$$

$$= 1000 \left\{ \frac{1.08 [1 - (1.08)^{26}]}{1 - 1.08} \right\}$$

= 86351 (to nearest dollar)

**Q6(ii)**  $S_n = \underbrace{1000 + 1080 + 1160 + \dots}_{n \text{ terms}} > 86351$

$$S_n = \frac{n}{2} [2(1000) + (n-1)(80)] > 86351$$

$$\Rightarrow 40n^2 + 960n - 86351 > 0$$

$$\Rightarrow n < -59.987 \text{ (N.A.)} \quad \text{or} \quad n > 35.987$$

∴ Least number of years that he still needs to save = 36

The year at which Mr Woo's savings in this savings plan will first exceed \$86351 = 2015 + 36 - 1 = 2050

### Q7

Rate of salt flowing into tank per minute is  $12 \times (0.125) = 1.5 \text{ kg}$

Rate of salt flowing out per minute is  $\frac{12}{400} \times q = 0.03q$

Therefore,  $\frac{dq}{dt} = 1.5 - 0.03q$ .

$$\frac{dq}{dt} = 1.5 - 0.03q$$

$$\int \frac{1}{1.5 - 0.03q} dq = \int 1 dt$$

$$-\frac{1}{0.03} \ln |1.5 - 0.03q| = t + C$$

$$|1.5 - 0.03q| = Ae^{-0.03t}$$

$$1.5 - 0.03q = Be^{-0.03t}$$

When  $t = 0, q = 100, 1.5 - 0.03(100) = B$

$$B = -1.5$$

$$1.5 - 0.03q = -1.5e^{-0.03t}$$

1.6 kg per litre =  $0.16 \times 400 = 64 \text{ kg of salt in the tank}$

Thus  $1.5 - 0.03(64) = -1.5e^{-0.03t}$


$$t = 42.4 \text{ min (3 s.f)}$$

$$0.03q = 1.5(1 + e^{-0.03t})$$

$$q = 50(1 + e^{-0.03t})$$

When  $t$  is large,  $e^{-0.03t} \rightarrow 0$

Thus, the amount of salt in the tank decreases to 50kg.



**Q8(i)**

$$\text{LHS} = ((x + iy)^2)^*$$

$$= (x^2 + (iy)^2 + 2xyi)^*$$

$$= (x^2 - y^2 + 2xyi)^*$$

$$= x^2 - y^2 - 2xyi$$

$$\text{RHS} = ((x + iy)^*)^2$$

$$= (x - iy)^2$$

$$= x^2 + (iy)^2 - 2xyi$$

$$= x^2 - y^2 - 2xyi$$

**Q8(ii)** Let  $z = x + iy$ , where  $x, y \in \mathbb{C}$

$$(x + iy)^2 = 1 - 4\sqrt{3}i$$

$$\Rightarrow \begin{cases} x^2 - y^2 = 1 & \text{--- (1)} \\ 2xy = -4\sqrt{3} & \text{--- (2)} \end{cases}$$

$$(2) \Rightarrow y = \frac{-2\sqrt{3}}{x}$$

$$(1) \Rightarrow x^2 - \frac{12}{x^2} = 1 \Rightarrow x^4 - x^2 - 12 = 0 \Rightarrow (x^2 - 4)(x^2 + 3) = 0$$

$$x \in \mathbb{C} \Rightarrow x^2 \neq 0 \Rightarrow x^2 = 4 \Rightarrow x = \pm 2, \quad y = \mp \sqrt{3}$$

$$\therefore z = 2 - \sqrt{3}i \quad \text{or} \quad -2 + \sqrt{3}i$$

**Q8(iii)**  $w^2 = 4 + 16\sqrt{3}i = 4(1 + 4\sqrt{3}i)$

$$\text{* both sides: } (w^2)^* = 4(1 - 4\sqrt{3}i)$$

$$\text{using (i): } (w^*)^2 = 4(1 - 4\sqrt{3}i)$$

$$\text{using (ii): } w^* = \sqrt{4}(2 - \sqrt{3}i) \quad \text{or} \quad \sqrt{4}(-2 + \sqrt{3}i)$$

$$= 4 - 2\sqrt{3}i \quad \text{or} \quad -4 + 2\sqrt{3}i$$

$$\therefore w = 4 + 2\sqrt{3}i \quad \text{or} \quad -4 - 2\sqrt{3}i$$

**Q8(iv)**  $z_1 = 2 - \sqrt{3}i$  and  $z_2 = -2 + \sqrt{3}i$

Given:  $\arg(z^2) = \theta$ .


$$\begin{aligned}\arg(z_1 z_2) &= \arg[(2 + \sqrt{3}i)(2 - \sqrt{3}i)] \\ &= \arg(-4 + 4\sqrt{3}i) \\ &= \arg(-1 + 4\sqrt{3}i) \\ &= \arg(-z^2) \\ &= \arg(-1) + \arg(-z^2) \\ &= \theta + \pi\end{aligned}$$

**Alternative**

Given:  $\arg(z^2) = \theta$  (where  $\theta < 0$ )

$$\begin{aligned}\arg(z_1 z_2) &= \arg(z_1) + \arg(z_2) \\ &= \frac{\theta}{2} + \left(\frac{\theta}{2} + \pi\right) \\ &= \theta + \pi\end{aligned}$$

**Q9(i)**



**Q9(ii)**  $\frac{dx}{dt} = 0.1, \frac{dy}{dt} = ?$  at  $x = \frac{1}{2}e^{\frac{\pi}{6}}$

$$e^\theta \sin \theta = \frac{1}{2}e^{\frac{\pi}{6}} \Rightarrow \theta = \frac{\pi}{6}$$

$$\frac{dy}{dx} = \frac{e^\theta \cos \theta - e^\theta \sin \theta}{e^\theta \cos \theta + e^\theta \sin \theta} = \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta}$$

$$\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$$

$$\frac{dy}{dt} = \frac{\cos\left(\frac{\pi}{6}\right) - \sin\left(\frac{\pi}{6}\right)}{\cos\left(\frac{\pi}{6}\right) + \sin\left(\frac{\pi}{6}\right)} \times (0.1)$$

$$= \frac{\sqrt{3}/2 - 1/2}{\sqrt{3}/2 + 1/2} (0.1)$$

$$= \frac{\sqrt{3}-1}{10(\sqrt{3}+1)}$$

$$\text{Q9(iii)} \quad \frac{dy}{dx} = \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta}$$

At point  $P$ , tangent // y-axis

$$\Rightarrow \frac{dy}{dx} \text{ is undefined}$$

$$\Rightarrow \cos \theta + \sin \theta = 0$$

$$\Rightarrow \tan \theta = -1$$

$$\Rightarrow \theta = -\frac{\pi}{4} \quad \left( \because -\frac{\pi}{2} < \theta \leq \frac{\pi}{2} \right)$$

$$x = e^{-\frac{\pi}{4}} \sin\left(-\frac{\pi}{4}\right) = -\frac{1}{\sqrt{2}} e^{-\frac{\pi}{4}}$$

Equation of tangent at point  $P$  is

$$x = -\frac{1}{\sqrt{2}} e^{-\frac{\pi}{4}}$$

#### **Q9(iv)**

$$\text{Coordinates of } P: \left( -\frac{1}{\sqrt{2}} e^{-\frac{\pi}{4}}, \frac{1}{\sqrt{2}} e^{-\frac{\pi}{4}} \right)$$

$$OP = \sqrt{\frac{1}{2} e^{-\frac{\pi}{2}} + \frac{1}{2} e^{-\frac{\pi}{2}}} = e^{-\frac{\pi}{4}}$$

Note that  $P$  lies on the line  $y = -x$  and  $OP \perp OQ$ ,

then  $Q$  lies on the line of  $y = x$ .

$$e^\theta \sin \theta = e^\theta \cos \theta$$

$$\tan \theta = 1$$

$$\theta = \frac{\pi}{4}$$

$$x = e^{\frac{\pi}{4}} \sin\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} e^{\frac{\pi}{4}}$$

$$y = e^{\frac{\pi}{4}} \cos\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} e^{\frac{\pi}{4}}$$

$$OQ = \sqrt{\frac{1}{2} e^{\frac{\pi}{2}} + \frac{1}{2} e^{\frac{\pi}{2}}} = e^{\frac{\pi}{4}}$$

$$\text{Area of } \Delta POQ = \frac{1}{2}(OP)(OQ) = \frac{1}{2} \left( e^{-\frac{\pi}{4}} \right) \left( e^{\frac{\pi}{4}} \right) = \frac{1}{2} \text{ units}^2$$

$$\mathbf{Q10(i)} \quad v_n = u_1 + u_2 + u_3 + \dots + u_n$$

$$\begin{aligned}
&= \frac{A}{1!} - \frac{2A^2}{2!} + \frac{A^3}{3!} \\
&+ \frac{A^2}{2!} - \frac{2A^3}{3!} + \frac{A^4}{4!} \\
&+ \frac{A^3}{3!} - \frac{2A^4}{4!} + \frac{A^5}{5!} \\
&+ \dots \\
&+ \frac{A^{n-2}}{(n-2)!} - \frac{2A^{n-1}}{(n-1)!} + \frac{A^n}{n!} \\
&+ \frac{A^{n-1}}{(n-1)!} - \frac{2A^n}{n!} + \frac{A^{n+1}}{(n+1)!} \\
&+ \frac{A^n}{n!} - \frac{2A^{n+1}}{(n+1)!} + \frac{A^{n+2}}{(n+2)!} \\
&= A - \frac{2A^2}{2!} + \frac{A^2}{2!} + \frac{A^{n+1}}{(n+1)!} - \frac{2A^{n+1}}{(n+1)!} + \frac{A^{n+2}}{(n+2)!} \\
&= A - \frac{A^2}{2} - \frac{A^{n+1}}{(n+1)!} + \frac{A^{n+2}}{(n+2)!} \quad (\text{shown})
\end{aligned}$$

$$\mathbf{Q10(ii)} \quad \sum_{n=2}^N \left\{ \frac{1}{N} \left( v_n + \frac{A^{n+1}}{(n+1)!} - \frac{A^{n+2}}{(n+2)!} \right) + 7^{n-N} \right\}$$

$$\begin{aligned}
&= \sum_{n=2}^N \left\{ \frac{1}{N} \left( A - \frac{A^2}{2} \right) + 7^{n-N} \right\} \\
&= \frac{1}{N} \left( A - \frac{A^2}{2} \right) (N-1) + \frac{1}{7^N} \sum_{n=2}^N 7^n \\
&= \left( \frac{N-1}{N} \right) \left( A - \frac{A^2}{2} \right) + \frac{1}{7^N} \left[ \frac{7^2(1-7^{N-1})}{1-7} \right] \\
&= \left( \frac{N-1}{N} \right) \left( A - \frac{A^2}{2} \right) + \frac{1}{7^N} \cdot \frac{7^2}{6} \cdot (7^{N-1} - 1) \\
&= \left( \frac{N-1}{N} \right) \left( A - \frac{A^2}{2} \right) + \frac{49}{6} \left( \frac{1}{7} - \frac{1}{7^N} \right)
\end{aligned}$$

$$\frac{N-1}{N} = 1 - \frac{1}{N}$$

As  $N \rightarrow \infty$ ,  $\frac{1}{N} \rightarrow 0$ , so  $1 - \frac{1}{N} \rightarrow 1$ . Also,  $\frac{1}{7^N} \rightarrow 0$ .

Hence as  $N \rightarrow \infty$ , Series  $\rightarrow A - \frac{A^2}{2} + \frac{7}{6}$

$\sum_{n=2}^N \left\{ \frac{1}{N} \left( v_n + \frac{A^{n+1}}{(n+1)!} - \frac{A^{n+2}}{(n+2)!} \right) + 7^{n-N} \right\}$  converges.

Limit  $= A - \frac{A^2}{2} + \frac{7}{6}$ .

**Q11)** Let  $M$  be a point on plane  $\pi$ .

$$\overrightarrow{OM} = \begin{pmatrix} 0 \\ 0 \\ 13 \end{pmatrix}, \overrightarrow{MP} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix},$$

$$\frac{\left| \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} a \\ -2 \\ 1 \end{pmatrix} \right|}{\sqrt{a^2 + 5}} = 1$$

$$\sqrt{a^2 + 5} = |a - 4 - 1|$$

$$(a^2 + 5) = a^2 - 10a + 25$$

$$a = 2$$

$$\textbf{Q11ii)} l_1 : r = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 3 \\ -2 \end{pmatrix}, \lambda \in \mathbb{R}$$

$$l_2 : r = \begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \mu \in \mathbb{R}$$

$$\overrightarrow{OA} = \begin{pmatrix} 2+4\lambda \\ 1+3\lambda \\ 3-2\lambda \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 4+\mu \\ \mu \\ 5 \end{pmatrix}$$

$$\overrightarrow{AB} = \begin{pmatrix} 2+\mu-4\lambda \\ \mu-3\lambda-1 \\ 2+2\lambda \end{pmatrix}$$

$$\overrightarrow{AB} \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = 0, \quad \overrightarrow{AB} \cdot \begin{pmatrix} 4 \\ 3 \\ -2 \end{pmatrix} = 0$$

$$1 = -2\mu + 7\lambda, \quad 1 = -7\mu + 29\lambda$$

Solving,

$$\therefore \mu = -\frac{22}{9}, \quad \lambda = -\frac{5}{9}$$

$$\overrightarrow{OA} = \begin{pmatrix} -\frac{2}{9} \\ -\frac{2}{3} \\ \frac{37}{9} \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} \frac{14}{9} \\ -\frac{22}{9} \\ 5 \end{pmatrix}$$

**Q11(ii)**

$$\left[ \begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right] \cdot \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} = 8 + 2\mu - 2\mu + 5 = 13$$

Hence  $l_2$  is in plane  $\pi$ .

**Alternative**

$$l_2: \quad r = \begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} = 2 - 2 = 0$$

$\Rightarrow l_2$  is perpendicular to  $\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$ .

$\Rightarrow l_2$  is parallel to  $\pi$ .

$$\begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} = 8 + 5 = 13$$


$\Rightarrow$  a point in  $l_2$  is also in  $\pi$ .

Hence  $l_2$  is in plane  $\pi$ .

**Q11(ii)**

$$\overrightarrow{OB} = \frac{\overrightarrow{OA} + \overrightarrow{OA'}}{2}$$

$$\overrightarrow{OA'} = 2 \begin{pmatrix} \frac{14}{9} \\ -\frac{22}{9} \\ 5 \end{pmatrix} - \begin{pmatrix} -\frac{2}{9} \\ -\frac{2}{3} \\ \frac{37}{9} \end{pmatrix} = \frac{1}{9} \begin{pmatrix} 30 \\ -38 \\ 53 \end{pmatrix}$$



reflection  
of  $l_1$


$$\text{Line of reflection of } l_1 \text{ in } \pi: \quad r = \frac{1}{9} \begin{pmatrix} 30 \\ -38 \\ 53 \end{pmatrix} + \beta \begin{pmatrix} 4 \\ 3 \\ -2 \end{pmatrix}, \beta \in \mathbb{R}$$

**Alternative**

Let  $D$  be the point  $(0, 0, 13)$  on  $\pi$

and  $C$  be the point  $(2, 1, 3)$  on  $l_1$

$$\overrightarrow{CD} = \begin{pmatrix} 0 \\ 0 \\ 13 \end{pmatrix} - \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} -2 \\ -1 \\ 10 \end{pmatrix}$$



reflection  
of  $l_1$

$$\overrightarrow{CF} = \left[ \begin{pmatrix} -2 \\ -1 \\ 10 \end{pmatrix} \cdot \frac{1}{\sqrt{9}} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} \right] \frac{1}{\sqrt{9}} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} = \frac{8}{9} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$$

$$\overrightarrow{OF} = \overrightarrow{CF} + \overrightarrow{OC} = \frac{8}{9} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} + \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \frac{1}{9} \begin{pmatrix} 34 \\ -7 \\ 35 \end{pmatrix}$$

$$\overrightarrow{OF} = \frac{\overrightarrow{OA} + \overrightarrow{OA'}}{2}$$

$$\overrightarrow{OA'} = 2 \begin{pmatrix} \frac{34}{9} \\ -\frac{7}{9} \\ \frac{35}{9} \end{pmatrix} - \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \frac{1}{9} \begin{pmatrix} 50 \\ -23 \\ 43 \end{pmatrix}$$

Line of reflection of  $l_1$  in  $\pi$ :  $r = \frac{1}{9} \begin{pmatrix} 50 \\ -23 \\ 43 \end{pmatrix} + \beta \begin{pmatrix} 4 \\ 3 \\ -2 \end{pmatrix}, \beta \in \mathbb{R}$

**Q11(iii)**

$$n = \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} \times \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 4 \end{pmatrix}$$

$$r \cdot \begin{pmatrix} -1 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 4 \end{pmatrix}$$

Equation of plane  $p$  is

$$-x + y + 4z = 16$$