UisitFREETESTPAPER.com


for more papers



Website: freetestpaper.com
Facebook.com/freetestpaper



Twitter.com/freetestpaper





JURONG SECONDARY SCHOOL 2022 GRADUATION EXAMINATION 2 SECONDARY 4 EXPRESS/ SECONDARY 5 NORMAL (ACADEMIC)

| MATHEMATICS | 4048/0 |
|-------------|-----------------|
| CLASS | INDEX NUMBER |
| NAME | |
| CANDIDATE 1 | |

26 Aug 2022 PAPER 1 2 hours

Candidates answer on the Question paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in. Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer all the questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.


The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place. For π , use either your calculator value or 3.142, unless the question requires the answer in

terms of π .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 80.



This document consists of 23 printed pages, including this page.

4048/1/GE2/22

[Turn Overl

Mathematical Formulae

Compound interest

Total amount =
$$P\left(1 + \frac{r}{100}\right)^n$$

Mensuration

Curved surface area of a cone = πrl

Surface area of a sphere $=4\pi r^2$

Volume of a cone
$$=\frac{1}{3}\pi r^2 h$$

Volume of a sphere
$$=\frac{4}{3}\pi r^3$$

Area of triangle
$$ABC = \frac{1}{2}ab\sin C$$

Arc length = $r\theta$, where θ is in radians

Sector area =
$$\frac{1}{2}r^2\theta$$
, where θ is in radians

Trigonometry

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Statistics

$$Mean = \frac{\sum fx}{\sum f}$$

Standard deviation =
$$\sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2}$$

| 1 | Simplify | -2(| [-3x-4] |)+5. |
|---|----------|-----|---------|------|
|---|----------|-----|---------|------|

| Answer | [1 | [] |
|--------|----|----|
|--------|----|----|

The ratio of interior angle: exterior angle of a regular polygon is 5:1.

Calculate the number of sides that the polygon has.


3 A photocopier can photocopy at a rate of 30 double-sided papers per minute.

Calculate

(a) in minutes, the time taken to print 600 pages on double-sided setting,

(b) the number of double-sided papers that can be photocopied in 1 hour.

A survey was conducted to find out the preferred brand of mobile phone that secondary school students like.




State one aspect of the graph that may be misleading and explain how this may lead to a misinterpretation of the graph.

Solve the inequality $-2+x \le 2-3x < 9+4x$.

Answer [2]

| 6 | Find a possib $ax + by = 1$ | ole set of integer valu | ues of a and b | such that the | lines $x+y=1$ and |
|---|-----------------------------|-------------------------------|--------------------|---------------|-------------------|
| | (a) do not in | ntersect, | | | |
| | • | | Answer | | [1] |
| | (b) intersect | t at exactly one point. | | | |
| | | | Answer | | [1] |
| 7 | Factorise 4ax | $x^2 - 4ay^2 + 2bx^2 - 2by^2$ | completely. | | |
| | | | | | |
| | | | | | |
| | | | Answe | r | [2] |



| Answer | | | | | | | | | | | | | | | | [2] | Ì |
|--------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|-----|---|
|--------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|-----|---|

Given that $25 \times 125^x = 5$, find the value of x.

10 Solve
$$\frac{5}{(2-x)^2} + \frac{1}{x-2} = 4$$
.

Answer
$$x = \dots [3]$$

11 (a) Express $1+x^2-5x$ in the form of $(x+a)^2+b$.

(b) Write down the equation of line of symmetry of the graph of $y = 1 + x^2 - 5x$.


(c) Write down the coordinates of intersection between the line of symmetry and the graph of $y = 1 + x^2 - 5x$.

| 12 | The area of triangle ABC is 7.5 cm ² . Two of the sides are of length 5 cm and 6 cm respectively. Find the length of the third side. |
|----|--|
| | |
| | • |
| | |
| | Answer |
| 13 | The Ideal Gas Law |
| | PV = 8.3145nT |
| | relates pressure P (measured in Pascal), volume V (measured in metres ³), the number of moles of a gas n (measured in moles), and temperature T (measured in Kelvins). |
| | One mole of gas is found to exert a pressure of 101 325 Pascal at room temperature 293 Kelvins. Find the volume of the gas in cm ³ . |
| | |
| | |
| | |
| | Answercm ³ [2] |

| (a) | The length of the Mississippi River is 3766 km. |
|------------|---|
| | Calculate the length, in centimetres, of Mississippi River on the map. |
| (b) | Answer |
| | Answerkm² [2] |
| (a) | n is a positive integer. Show that, for all n , $6n^2 + 25n + 11$ is not a prime number. |
| | |
| (b) | Hence, determine whether 16261 is a prime number. Show your working clearly. |
| •••••• | |
| | [2] |
| - (| a) |

16 The diagram represents a plot of land, ABCD. The police is looking for a robber hiding in the land.

Scale: 1 cm represents 1 km



Here are the accounts from three witnesses:

Witness 1: The robber is located more than 6 km away from A.

Witness 2: The robber is located nearer to B than A.

Witness 3: The robber is located nearer to AD than AB.

Shade the region where the robber is hiding.

[4]

| 17 | ξ={ | $\{\text{integers } x: 1 < x < 21\}$ | |
|----|------------|--|--|
| | A = | (integers that are perfect squares) | |
| | B = | {integers that are not prime numbers} | |
| | <i>C</i> = | {integers that are divisible by 5} | |
| | (a) | List down all the elements in A , B and | 1 <i>C</i> . |
| | | | |
| | | | Answer A = |
| | | | <i>B</i> = |
| | | | <i>C</i> =[2] |
| | A nu | number is chosen randomly from ξ . | |
| | (b) | Find the number of elements in $B \cap C$ | C'. |
| | | | |
| | | | |
| | | | |
| | | | Answer[1] |
| | (c) | Hence, find the probability that the one divisible by 5. | chosen number is not a prime number and is |
| | | | |
| | | | |
| | | | Accessory [13] |
| | | | Answer[1] |

| 18 | (a) | The prime factorisation of 450 is $2 \times 3^2 \times 5^2$. Express 648 as a product of its prime factors. |
|----|-----|---|
| | (b) | Answer 648 =[1] Using your answer to part (a), determine whether 583200 is a perfect square. |
| | | |
| | (c) | Find the smallest integer n such that $450n$ is a multiple of 648 . |

Answer n =[1]

| 19 | (a) | The points $A(0,2)$ and $B(1,0)$ lie on the graph $y = x^2 + ax + b$. |
|----|-----|--|
| | | Find the values of a and b . |

13

| Answer | <i>a</i> = |
|--------|---------------|
| | <i>b</i> =[2] |

(b) C is another point on the graph such that the gradient of line AB is twice the gradient of AC. Find the coordinates of point C.

Answer (.....) [3]

| 20 | It is given that | $\overline{AB} =$ | $\begin{pmatrix} -12 \\ a \end{pmatrix}$ | $\bigg), \ a>0$ | and | $\left \overline{BA} \right = 37$ | unit |
|----|------------------|-------------------|--|-----------------|-----|-------------------------------------|------|
| 20 | It is given that | $\overline{AB} =$ | $\begin{pmatrix} -12 \\ a \end{pmatrix}$ | a>0 | and | $\left \overline{BA} \right = 37$ | uni |


(a) Find the value of a.

| _ | | | | | | | | | | | | | | | | _ | ~ - | |
|--------|---|---|--|------|--|--|--|--|------|--|----|----|--|------|------|---|-----|---|
| Answer | а | = | | | | | | | | | ٠. | ٠. | | | | ľ | 2 | l |

(b) C is a point (0,10) and $\overrightarrow{AB} = \overrightarrow{DC}$. Find the coordinates of D.

(c) What type of quadrilateral is ABCD? Explain your answer.

The cumulative frequency curve shows the marks obtained by the 120 students in a recent Mathematics Test. The maximum mark is 100.



(a) Complete the grouped frequency table for the marks obtained.

| Marks obtained by students, x | Frequency |
|-------------------------------|-----------|
| $0 \le x < 20$ | 10 |
| $20 \le x < 40$ | |
| $40 \le x < 60$ | |
| $60 \le x < 80$ | 28 |
| $80 \le x < 100$ | 8 |

[1]

(b) Calculate an estimate of the mean mark.

$$0 \le x < 20$$

$$20 \le x < 40$$

$$40 \le x < 60$$


$$60 \le x < 80$$

$$80 \le x < 100$$

Answermarks [1]

| (c) | Calculate an estimate of the standard deviation. |
|-----|---|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | Answermarks [2] |
| (d) | The passing mark is 50. Two students are chosen at random. Find the probability that both of them passed. |
| | |
| | |
| | |
| | Answer[1] |
| | |

22 The diagram shows the speed-time graph of a particle.



(a) Find the speed of the particle when t = 12s.


Answerm/s [2]

(b) Find the distance covered before the particle starts to slow down.


Answerm [2]

(c) Complete the distance-time graph of the particle.

[1]



23




In the diagram, ABC and AED are straight lines and BE is parallel to CD.

| (a) | Show that triangle ABE is similar to triangle ACD . Give a reason for each state | | | | | | | | | | |
|-----|--|--|--|--|--|--|--|--|--|--|--|
| | you make. | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |

(b) Show that
$$\frac{BC}{ED} = \frac{AB}{AE}$$
. [2]

| (c) | A circle is drawn with points B, C and D on its circumference. It is further given that angle BCD + angle $BED > 180^{\circ}$. Explain why point E is inside the circle |
|------|---|
| | |
| | |
| | |
| •••• | [1 |

24



OBC, ADC, AEB and OED are straight lines.

$$\overrightarrow{OA} = \mathbf{a}$$
 and $\overrightarrow{OB} = \mathbf{b}$.

$$OB = \frac{1}{3}OC$$
, $AE : EB = 3:2$ and $OE : ED = 1:1$.


(a) Find \overrightarrow{OE} in terms of a and/or b as simply as possible.

Answer: $\overrightarrow{OE} = \dots [2]$

(b) Find the value of $\frac{\text{area of } \triangle OBE}{\text{area of } BEDC}$.

Answer [2]

25



AC is a diameter of a circle with centre O. $AC = \sqrt{2}r$ and AB = BC. COB is a semi-circular arc, centre P. What percentage of the circle is **not** shaded?

Answer% [5]





JURONG SECONDARY SCHOOL 2022 GRADUATION EXAMINATION 2 SECONDARY 4 EXPRESS/ SECONDARY 5 NORMAL ACADEMIC

| CANDIDATE NAME | |
|-------------------|-----------------|
| CLASS | INDEX NUMBER |

MATHEMATICS

4048/02

PAPER 2

24 August 2022 2 hour 30 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all the questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For π , use either your calculator value or 3.142, unless the question requires the answer in terms of π .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 100.

For Examiner's Use

This document consists of 26 printed pages including this page.

Mathematical Formulae

Compound interest

Total amount =
$$P\left(1 + \frac{r}{100}\right)^n$$

Mensuration

Curved surface area of a cone = πrl

Surface area of a sphere = $4\pi r^2$

Volume of a cone =
$$\frac{1}{3}\pi r^2 h$$

Volume of a sphere =
$$\frac{4}{3}\pi r^3$$

Area of triangle
$$ABC = \frac{1}{2}ab\sin C$$

Arc length = $r\theta$, where θ is in radians

Sector area =
$$\frac{1}{2}r^2\theta$$
, where θ is in radians

Trigonometry

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Statistics

$$Mean = \frac{\sum fx}{\sum f}$$

Standard deviation =
$$\sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2}$$

Answer all the questions.

- 1 (a) It is given that $y = \frac{5x-2}{3x+1}$.
 - (i) Find y when x = 7.

Answer
$$y = \dots$$
 [1]

(ii) Express x in terms of y.

Answer
$$x = \dots$$
 [2]

(b) Solve $\frac{x-5}{3} - \frac{x+3}{2} = -4$.

Answer
$$x = \dots [2]$$

| (c) Solve these | simultaneous | equations. |
|-----------------|--------------|------------|
|-----------------|--------------|------------|

$$3x - 2y = 10$$

$$2x + y = 23$$

| Answer | x = | |
|--------|------------|---------|
| | <i>y</i> = | [3] |

(d) Simplify $\frac{4x^2-9}{6x^2-x-12}$.

Answer[3]

2 Mrs Teo is a Mathematics tutor.

She offers tutorial sessions for Lower Secondary and Upper Secondary students on weekdays and on weekends.

Each student attends one session a week for 4 sessions in every month.

The matrix N shows the number of students she tutors every month.

Lower Secondary Upper Secondary

$$\mathbf{N} = \begin{pmatrix} 7 & 6 & \\ 4 & 3 & \end{pmatrix}$$
 Weekday Weekend

(a) Evaluate the matrix M = 4N.

(b) Mrs Teo charges \$70 for each Lower Secondary session and \$80 for each Upper Secondary session.

Represent the session charges in a 2×1 column matrix C.

Answer
$$C = [1]$$


(c) Evaluate the matrix P = MC.

Answer
$$P = [1]$$

| | 6 | |
|---------------------|---|-----|
| (d) | State what the elements of P represent. | |
| | Answer | |
| | | [1] |
| mon The stude | Teo wants to increase her tuition fees for the weekend sessions by 10% for the last 3 ths before examinations start. number of students registered for the weekday sessions are 10 Lower Secondary ents and 8 Upper Secondary students. On weekends she has 3 Lower Secondary ents and 2 Upper Secondary students. | |
| (e) | Calculate the total amount of money she earns during this 3 month period. | |
| | | |
| | | |
| | | |
| | | |
| | Answer | [4] |

7

3



The diagram below shows part of a regular 18-sided polygon, ABCDEF.... The lines BC and ED are extended to meet at the point X such that XC = XD.

(a) Find $\angle XCD$.

| Answer | , | [1] |
|--------|---|-----|
| | | |

(b) Find $\angle CXD$.

(c) Explain why BX = EX.

Answer

(d) Show that ΔXCD and ΔXBE are similar.

Answer

 4 The first four terms in a sequence of numbers are given below.

$$T_1 = 5^2 - 18 = 7$$

$$T_2 = 6^2 - 22 = 14$$

$$T_3 = 7^2 - 26 = 23$$

$$T_4 = 8^2 - 30 = 34$$

(a) Find T_5 .

| Answer | | [1] |
|--------|--|-----|
|--------|--|-----|

(b) Show that the *n*th term of the sequence, T_n , is given by $n^2 + 4n + 2$.

Answer

[2]

(c) Determine and explain if 962 is a term of the sequence.

Answer

[2]

| (d) | Find and si | mplify an | expression, | in terms | of n , | for T_n | $-T_{n-1}$ |
|-----|-------------|-----------|-------------|----------|----------|-----------|-------------|
| () | | 1 2 | , | | • | " | <i>n</i> -1 |

| | Answer | [3] |
|-----|---|-----|
| (e) | Explain why the difference between consecutive terms in the sequence is always odd. | |
| | Answer | |
| | | |
| | | [1] |

5 The variables x and y are connected by the equation

$$y = x + \frac{5}{x^2} - 3$$


Some corresponding values of x and y are given in the table below.

| x | 1 | 1.5 | 2 | 2.5 | 3 | 4 | 5 | 6 |
|---|---|-----|-----|-----|-----|-----|---|-----|
| y | 3 | 0.7 | 0.3 | 0.3 | 0.6 | 1.3 | p | 3.1 |

(a) Find the value of p.

Answer
$$p = \dots$$
 [1]

(b) On the grid, draw the graph of
$$y = x + \frac{5}{x^2} - 3$$
 for $1 \le x \le 6$.




| (c) | The equation $x + \frac{5}{x^2} - 3 = 0$ has no solution. Explain how this can be seen from your graph. | | | | | | |
|-----|--|---|--|--|--|--|--|
| | | | | | | | |
| | | [1] | | | | | |
| | 7.33 | De descripce descript find the small out of the same when the | | | | | |

(d) By drawing a tangent, find the gradient of the curve when x = 1.5.

Answer [2]

| (e) | (i) | On the same axes, draw the line with gradient 2 that passes through the point with coordinates $(0.5, 2)$. | [2] |
|-----|-------|---|-----|
| | (ii) | Write down the equation of this line. | |
| | | | |
| | | Answer | [1] |
| | (iii) | Show, with mathematical justification, that the equation $5-4x^2-x^3=0$ can be solved by finding the points of intersection of the straight line and the curve. | |
| | | Answer | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | [0] |
| | (iv) | Use your graphs to solve the equation $5-4x^2-x^3=0$. | [2] |
| | | | |
| | | | |
| | | | |
| | | Answer | [1] |
| | | Answer | [1] |

6



The diagram shows two concentric circles with centre O. PQ is the diameter of the larger circle and RS is the diameter of the smaller circle. PS and RQ are tangents to the smaller circle.

(a) Show that the triangle *PSO* is congruent to triangle *QRO*. Give a reason for each statement you make.

Answer

| (b) | The | radius of the larger circle is 8 cm and the radius of the smaller circle is 4.36 cm. | |
|-----|------|--|----|
| | (i) | Calculate the area of triangle QRO. | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | Answer | 3] |
| | (ii) | Given that angle QOR is 0.995 radians, calculate the shaded area. | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | Answer cm^2 [| 4] |

7 The seating capacity of an auditorium is 300. The seats are categorised as 1, 2, 3 and 4. The costs of the tickets in the categories are in the table below.

| CATEGORY | TICKET PRICES |
|----------|---------------|
| CAT 1 | S\$ 348 |
| CAT 2 | S\$ 288 |
| CAT 3 | S\$ 228 |
| CAT 4 | S\$ 148 |

| (a) | The number of seats are distributed in the ratio 1:2:4:5 for CAT 1 to 4 respectively. |
|-----|---|
| | If all the seats are sold out, calculate the ticket sales for category 4. |

| Answer | \$ | | | | | | | • | | | | | | | | | | | | | | | | | | | | | | [| 2 |) | |
|--------|----|--|--|--|--|--|--|---|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|---|---|---|--|
|--------|----|--|--|--|--|--|--|---|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|---|---|---|--|

The auditorium has 2 entrances, front entrance F and back entrance B. Entrance F is able to evacuate x people out of the auditorium per second.

(b) (i) Write down an expression, in terms of x, for the time taken in seconds for all 300 people to evacuate if only entrance F is open.

| Answer | s | [1] |
|--------|---|-----|
| | | L-1 |

(ii) Entrance F is able to let approximately 2 more people evacuate per second compared to entrance B.

Write down an expression, in terms of x, for the time taken in seconds for all 300 people to evacuate the auditorium if only entrance B is open.

(iii) Usually, only one entrance is open at any time.

| It takes approximately 3 seconds more for all the people to evacuate the |
|--|
| auditorium using entrance B as compared to entrance F . |

Form an equation in x and show that it reduces to $x^2 - 2x - 200 = 0$.

Answer

[3]

(iv) Solve the equation $x^2 - 2x - 200 = 0$.


Answer x = or [2]

(v) Find the number of people that can be evacuated in 1 minute if only entrance F is opened.

Answer people [2]


8 An equilateral triangle ABC, has sides of 8 cm.

O is the centre of the equilateral triangle.



(a) Show that the length of *OA* is 4.619 cm, correct to 3 decimal places. *Answer*

The triangle ABC forms the base of a pyramid. The vertex, X is vertically above O and AX = 18 cm.



(b) Calculate the total surface area of the pyramid.

(c) Part of the pyramid is cut off at triangle PQR such that P is a point on XC, Q is a point on AC and R is a point at BC.
 PQ = 5 cm, QR = 4 cm and PR = 3.5 cm.
 Find angle PQR.

| A | 0 | 「つつ |
|--------|---|-----|
| Answer | | [4] |

(d) The ratio of AQ : QC = 3 : 1. Calculate the angle of elevation of X from Q.

Answer [4]

| 9 (a) A group of 50 patients had their blood pro |
|--|
|--|

The results are shown in the table below.

| Blood | $100 < x \le 120$ | $120 < x \le 140$ | $140 < x \le 160$ | $160 < x \le 180$ |
|-----------------|-------------------|-------------------|-------------------|-------------------|
| Pressure (mmHg) | | | | |
| Frequency | 27 | 12 | 6 | 5 |


| | Frequency | 27 | 12 | 6 | 5 | |
|-------|-----------------------------------|---------------------------------------|--|---|---------------------|-------|
| (i) | State the medi | an class of blood | pressure. | | | |
| | | | | | | . [1] |
| (ii) | A blood pressu Find the percer | re of 140 mmHg ntage of patients v | or higher indication who may have high | es high blood pres gh blood pressure | ssure. | |
| | | | | | | |
| | | | | Answer | % | [1] |
| (iii) | Calculate the s | tandard deviation | of the blood pres | ssures. | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | Answer | mmHg | g [2] |
| (iv) | The standard d patients was 22 | leviation of the bl 2.1 mmHg. | ood pressures tak | en by a second gr | roup of | |
| | Comment on o information. | ne difference bet | ween the two dist | ributions, making | g reference to this | |
| | Answer | | | | | • |
| | | | | | ., | • |
| | ••••• | | | | | . [1] |

| (b) | Two If th | awer contains 2 blue socks and 6 white socks. socks are taken from the drawer at random without replacement. e two socks are different colours, then a third sock is taken from the drawer. erwise, no third sock is taken. |
|-----|--------------|--|
| | (i) | Draw a tree diagram to show the probabilities of the possible outcomes. |
| | | Answer |

Answer [2]

10 Mr Graham bought a 2-room flat in Sembawang.

The floor plan can be modelled as five rectangles as shown below.



(a) Calculate the total floor area of Mr Graham's kitchen.


Answer m^2 [1]

(b) The optimal kitchen cabinets' height, including the counter top, is calculated based on the concept of ergonomics. Ergonomics is the process of designing products that fit the people who use them.

Using the diagram as reference, kitchen cabinet designers view that optimal cabinet height should be the distance between one's elbow and the ground, with 10 cm allowance for elbow movement.


The diagram below shows a drawing of Mrs Graham whose height is 168 cm.

Calculate the range of optimal kitchen cabinets' heights, including the counter top, for Mrs Graham.



| Answer | | cm | [2] |
|--------|--|----|-----|
|--------|--|----|-----|

(c) Mr Graham plans to install a combination of kitchen cabinets along the L-shaped shaded area in the kitchen.



In the combination, he requires one of the cabinets to have a sink, a corner cabinet and at least one of the other cabinets to have wire basket shelves.

He wants to customise the height of the cabinets to suit Mrs Graham's height. He also wishes to install a counter top for the top of the cabinet, considering to use either granite, ceramic or acrylic as its material.

He has a budget of \$2200 to purchase the kitchen cabinets and L-shaped counter top. Using the information in the tables given on the next page and your answer in (b), propose a possible combination that will best suit Mr Graham's requirements that will optimise storage and space while maximising the use of his budget.

Answer

Table 1: Kitchen Cabinets

| Cabinet | | 2 | 3 | 74 | 5 | 6 |
|-------------|---|-------------------|------------------------------|------------------------------------|---|--|
| Description | Cabinet with sink, 1 drawer and 1 door (no shelves) | Corner cabinet | Cabinet with 4 drawers | Cabinet with 2 shelves and 2 doors | Cabinet with 1 door and wire basket shelves | Cabinet with 1 drawer, 3 shelves and 1 door. |
| Cost | \$156 | \$166 | \$260 | \$290 | \$166 | \$193 |
| Width 50 cm | | 60 cm | 90 cm | 80 cm | 60 cm | 70 cm |
| Depth 60 cm | | 60 cm | 60 cm | 60 cm | 60 cm | 60 cm |

Table 2: Height Customisation for Kitchen Cabinet


| Height | Cost per cabinet |
|--------------|------------------|
| 50 cm | Free |
| up to 75 cm | .\$15 |
| up to 105 cm | \$25 |

Table 3: Materials for L-Shaped Counter Top

| Materials | Overall Cost |
|-----------|--------------|
| Granite | \$1200 |
| Ceramic | \$800 |
| Acrylic | \$600 |

End of Paper

2 hours





JURONG SECONDARY SCHOOL 2022 GRADUATION EXAMINATION 2 SECONDARY 4 EXPRESS/ SECONDARY 5 NORMAL (ACADEMIC)

| CANDIDATE NAME | |
|-------------------|-----------------|
| CLASS | INDEX NUMBER |
| MATHEMATICS | 4048/01 |
| PAPER 1 | 26 Aug 2022 |

Candidates answer on the Question paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in. Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer all the questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.


If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place. For π , use either your calculator value or 3.142, unless the question requires the answer in

terms of π .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.



This document consists of 23 printed pages, including this page.

Mathematical Formulae

Compound interest

Total amount =
$$P\left(1 + \frac{r}{100}\right)^n$$

Mensuration

Curved surface area of a cone = πrl

Surface area of a sphere = $4\pi r^2$

Volume of a cone =
$$\frac{1}{3}\pi r^2 h$$

Volume of a sphere
$$=\frac{4}{3}\pi r^3$$

Area of triangle
$$ABC = \frac{1}{2}ab\sin C$$

Arc length = $r\theta$, where θ is in radians

Sector area =
$$\frac{1}{2}r^2\theta$$
, where θ is in radians

Trigonometry

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Statistics

$$Mean = \frac{\sum fx}{\sum f}$$

Standard deviation =
$$\sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2}$$

| 1 | Simplify | -2(-3x | -4)+ | 5. |
|---|----------|--------|------|----|
|---|----------|--------|------|----|

$$-2(-3x-4)+5=6x+8+5$$

$$=6x+13$$

B1

| Answer | | ľ | ĺ | • |
|--------|--|---|---|---|
|--------|--|---|---|---|

2 The ratio of interior angle: exterior angle of a regular polygon is 5:1. Calculate the number of sides that the polygon has.

Exterior angle =
$$\frac{1}{1+5} \times 180^\circ = 30^\circ$$

Number of sides =
$$\frac{360^{\circ}}{30^{\circ}}$$
 = 12

BI

Answer sides [1]

3 A photocopier can photocopy at a rate of 30 double-sided papers per minute.

Calculate

(a) in minutes, the time taken to print 600 pages on double-sided setting, 600 pages = 300 double sided pages:

Time taken
$$=\frac{300}{30}=10$$
 animises

B,I

Answer _____ minutes [1]

(b) the number or nonble-sided papers that can be photocopied in 1 hour. 1 hour = 60 minutes

No. of pages printed = $30 \times 60 = 1800$


B1

Answerdouble-sided papers [1]

4048/1/GE2/22

[Turn Over]

4 A survey was conducted to find out the preferred brand of mobile phone that secondary school students like.



State one aspect of the graph that may be misleading and explain how this may lead to a misinterpretation of the graph

The y-axis does not start with 0. MI

Hence it looks take there are usee times the immber of students choosing Samsung than Oppo, which in fact is only two times. B1 (14 students for Samsung, 7 students for Oppo)

| | | | | | | [3] |
|------|--|--|------|------|------|------------|
| | | | | | | * * * E ** |

Solve the inequality $-2+x \le 2-3x < 9+4x$.

$$-2+x \le 2-3x < 9+4x^2$$

MI: Split inequality

 $-2+x \le 2-3x$ AND 2-3x < 9+4x

 $4x \le 4$ AND -7 < 7x

 $x \le 1 \text{ AND } x > -1$

 $\therefore -1 < x \le 1$

Al: (Don't give if there is no AND)

Answer [2]

and the same and the same of the same and the

4048/1/GE2/22

| 6 | Find a possible se | et of | integer | values | of a | and | b | such | that | the | lines | x+y=1 | and |
|---|--------------------|-------|---------|--------|------|-----|---|------|------|-----|-------|-------|-----|
| | ax + by = 1 | | | | | | | | | | | | |

| (a) do 1 | not intersect |
|----------|---------------|
|----------|---------------|

a = 2

b = 2

B1: Any integer values of a and b such that a = b but $a \neq 0,1$, $b \neq 0,1$

Answer a =

$$b = \dots [1]$$

(b) intersect at exactly one point.

a = 1

b = 2

B1: Any integer values of a and b such that $a \neq b$

Answer a =

$$b = \dots [1]$$

7 Factorise $4ax^2 - 4ay^2 + 2bx^2 - 2by^2$ completely.

$$4ax^2 - 4ay^2 + 2bx^2 - 2by^2$$

$$=4a(x^{2}-y^{2})+2b(x^{2}-y^{2})$$

$$= \left(4a + 2b\right)\left(x^2 - y^2\right)$$

$$=2(2a+b)(x-y)(x+y)$$

B1:
$$(x+y)(x-y)$$

8 Simplify
$$\left(\frac{a^{16}}{b^{-6}}\right)^{\frac{3}{2}}$$
.

$$\left(\frac{a^{16}}{b^{-6}}\right)^{\frac{3}{2}} = \frac{\left(a^{16}\right)^{\frac{3}{2}}}{\left(b^{-6}\right)^{\frac{3}{2}}}$$
$$= \frac{a^{24}}{b^{-9}}$$
$$= a^{24}b^{9}$$

M1: Correct multiplication of power to the indices respectively

A1: Must be in positive index notation

| Answer | | | | | | | | | | | | | | | | | | | | | | | [2 |) |
|--------|------|------|---|----|-------|----|---|---|---|-----|----|-------|---|---|---|----|---|---|---|---|---|-----|----|----|
| Answei | | | ٠ | ٠. | ٠ | ٠. | ٠ | ٠ | ٠ | • • | ٠, | • | ٠ | ٠ | • | ٠. | • | ٠ | ٠ | ٠ | ٠ | • • | L٤ | ٠. |

Given that $25 \times 125^x = 5$, find the value of x.

$$25 \times 125^x = 5$$

M1: Convert all bases to base 5 correctly

$$5^2 \times \left(5^3\right)^x = 5$$

$$5^2 \times 5^{3x} = 5$$

$$5^{2+3x} = 5^1$$

$$\therefore 2 + 3x = 1$$

$$x = -\frac{1}{3}$$

A1

Answer $x = \dots [2]$

10 Solve
$$\frac{5}{(2-x)^2} + \frac{1}{x-2} = 4$$
.

$$\frac{5}{(2-x)^2} + \frac{1}{x-2} = 4$$

$$\frac{5}{(x-2)^2} + \frac{1}{x-2} = 4$$

$$\frac{5}{(x-2)^2} + \frac{x-2}{(x-2)^2} = 4$$

$$\frac{5+x-2}{\left(x-2\right)^2}=4$$

$$\frac{3+x}{\left(x-2\right)^2}=4$$

$$3+x=4(x-2)^2$$

$$3 + x = 4\left(x^2 - 4x + 4\right)$$

$$3 + x = 4x^2 - 16x + 16$$

$$4x^2 - 17x + 13 = 0$$

$$(x-1)(4x-13)=0$$

$$x = 1 \text{ or } x = 3.25$$

M1: Common denominator (only accept quadratic denominator)

M1: Factorisation or quadratic formula

A1

Answer $x = \dots [3]$

11 (a) Express $1+x^2-5x$ in the form of $(x+a)^2+b$.

$$1 + x^2 - 5x = x^2 - 5x + 1$$

$$= \left(x - \frac{5}{2}\right)^2 - \left(\frac{5}{2}\right)^2 + 1$$

$$= \left(x - 2.5\right)^2 - 5.25$$

$$= \left[x + \left(-2.5\right)\right]^2 + \left(-5.25\right)$$

B1: Must be this form

Answer[1]

(b) Write down the equation of line of symmetry of the graph of $y = 1 + x^2 - 5x$. Eqn of line of symmetry: x = 2.5

Answer[1]

| (c) | Write down the coordinates of intersection between the line of symmetry and the |
|-----|---|
| | graph of $y = 1 + x^2 - 5x$. |

Point of intersection is at the turning point.

$$(2.5, -5.25)$$

B1

Answer (.....) [1]

The area of triangle ABC is 7.5 cm². Two of the sides are of length 5 cm and 6 cm respectively. Find the length of the third side.

Area of $\Delta = 7.5$

M1: Using Sine Rule correctly

$$\frac{1}{2}(5)(6)\sin\theta = 7.5$$

$$\sin\theta = \frac{1}{2}$$

$$\theta = \sin^{-1}\frac{1}{2} \text{ or } 180^{\circ} - \sin^{-1}\frac{1}{2}$$

$$\theta = 30^{\circ} \text{ or } 150^{\circ}$$

Length of third side

M1: Using Cosine Rule correctly (FT)

$$= \sqrt{5^2 + 6^2 - 2(5)(6)\cos 30^\circ}$$

 $\cap R$

$$\sqrt{5^2+6^2-2(5)(6)\cos 150^\circ}$$

:. Length of third side = 3.01 cm or 10.6 cm

A1: Both correct

13 The Ideal Gas Law

$$PV = 8.3145nT$$

relates pressure P (measured in Pascal), volume V (measured in metres³), the number of moles of a gas n (measured in moles), and temperature T (measured in Kelvins).

One mole of gas is found to exert a pressure of 101 325 Pascal at room temperature 293 Kelvins. Find the volume of the gas in cm³.

$$V = \frac{8.3145(1)(293)}{101325}$$
= 0.0240 m³ (3 s.f)
= 24000 cm³
A1

- A map of the United States of America has a scale of 1:8 000 000. 14
 - The length of the Mississippi River is 3766 km. Calculate the length, in centimetres, of Mississippi River on the map.

1 cm: 8000000 cm

1 cm: 80 km

 $80 \text{ km} \rightarrow 1 \text{ cm}$

$$3766 \text{ km} \rightarrow \frac{3766}{80} = 47.075 \text{ cm}$$

B1: 47.075

The area of California on the map is 66.25 cm². Calculate the actual area, in square kilometres, of California.

1 cm: 8000000 cm

M1: Using proportion

1 cm: 80 km

 $\therefore 1 \text{ cm}^2 : 6400 \text{ km}^2$

 $1 \text{ cm}^2 \rightarrow 6400 \text{ km}^2$

$$66.25 \text{ cm}^2 \rightarrow 66.25(6400) = 424000 \text{ km}^2$$

A1

Answerkm² [2]

15 n is a positive integer.


Show that, for all n, $6n^2 + 25n + 11$ is not a prime number.

$$6n^2 + 25n + 11 = (3n+11)(2n+1)$$

B1

For all positive integer n, 3n+11 and 2n+1are also integers more than 1. Hence for all positive n, $6n^2 + 25n + 11$ is a composite

| number since it can be factorised into smaller B1 : With integers greater than 1. | th the idea of factors more than 1 |
|--|--|
| | |
| | [2] |
| (b) Hence, determine whether 16261 is a prime $6n^2 + 25n + 11 = 16261$ $6n^2 + 25n - 16250 = 0$ (n-50)(6n+325) = 0 $n = 50$ or $n = -\frac{325}{6}$ (NA, since <i>n</i> is a positive integer) | M1: Can also use previous part to guess $n = 50$ (No working required) |
| By previous part, 16261 is a composite number. | negative minus 1) |
| | |
| | |
| | [2] |
| The diagram represents a plot of land, ABCD. The in the land. | ne police is looking for a robber hiding |
| Scale: 1 cm represents 1 km | |



Here are the accounts from three witnesses:

Witness 1: The robber is located more than 6 km away from A.

Witness 2: The robber is located nearer to B than A.

Witness 3: The robber is located nearer to AD than AB.

Shade the region where the robber is hiding.

[4]

| 17 $\xi = \{\text{integers } x : 1 < x < 21\}$ $A = \{\text{integers that are perfect squares}$ $B = \{\text{integers that are not prime numb}$ $C = \{\text{integers that are divisible by 5}\}$ | |
|---|---|
| (a) List down all the elements in A , $A = \{4,9,16\}$ $B = \{4,6,8,9,10,12,14,15,16,18,20\}$ $C = \{5,10,15,20\}$ | B2 Minus one mark per mistake Answer $A = \dots$ |
| | <i>B</i> = |
| | <i>C</i> =[2] |
| A number is chosen randomly from ξ | ; ; |
| (b) Find the number of elements in $B = \{4, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20\}$ $C = \{5, 10, 15, 20\}$ $B \cap C' = \{4, 6, 8, 9, 12, 14, 16, 18\}$ \therefore Required number = 8 | B∩C'. B1 |
| | Answer[1] |
| (c) Hence, find the probability that not divisible by 5. Required probability = $\frac{8}{19}$ | B1: Follow through from part (b) answer, but denominator must be 19 |
| | Answer[1] |

The prime factorisation of 450 is $2 \times 3^2 \times 5^2$.

Express 648 as a product of its prime factors.

18

(a)

| $648 = 2^3 \times 3^4$ | B1 | | | | |
|--|--|--|--|--|--|
| e e e | | | | | |
| | | | | | |
| | | | | | |
| · | | | | | |
| | | | | | |
| | | | | | |
| | Answer 648 =[1] | | | | |
| (b) Using your answer to part (a), determined that $450 \times 648 = 291600$ | ermine whether 583200 is a perfect square. | | | | |
| $\therefore 583200 = 2 \times 450 \times 648 = 2^5 \times 3^6 \times 5^2$ | B1 | | | | |
| 583200 is not a perfect square as the power | | | | | |
| (exponent) of base 2 in 583200 is not an even | - | | | | |
| number. | B1 | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | ••••••••••••••••••••••••••••••••••••••• | | | | |
| | [2] | | | | |
| •••••• | [2] | | | | |
| (c) Find the smallest integer n such that 450n is a multiple of 648. | | | | | |
| LCM (450,648) | | | | | |
| $n = \frac{7}{450}$ | | | | | |
| $=\frac{2^3 \times 3^4 \times 5^2}{2 \times 3^2 \times 5^2}$ | | | | | |
| | | | | | |
| $=2^2\times3^2$ | | | | | |
| = 36 | B1 | | | | |
| | Answer $n = \dots [1]$ | | | | |
| | | | | | |

BP~170

19 (a) The points A(0,2) and B(1,0) lie on the graph $y = x^2 + ax + b$. Find the values of a and b.

When x = 0, y = 2:

$$2 = b$$

B1

When
$$x = 1$$
, $y = 0$:

$$0 = 1^2 + a + 2$$

$$a = -3$$

B1

Answer a =

M1: Formulating equation with gradients

$$b = \dots [2]$$

(b) C is another point on the graph such that the gradient of line AB is twice the gradient of AC. Find the coordinates of point C.

Method 1

Let
$$C = (c, c^2 - 3c + 2)$$

Gradient of $AB = 2 \times Gradient$ of AC

B1: Gradient of AB

and solve

$$\frac{2-0}{0-1} = 2\left(\frac{c^2 - 3c + 2 - 2}{c}\right)$$

$$-2=2\left(\frac{c^2-3c}{c}\right)$$

$$-1 = \frac{c^2 - 3c}{c}$$

$$c^2 - 3c = -c$$

$$c^2-2c=0$$

$$c(c-2)=0$$

c = 0 (NA, since will get back point A) OR c = 2 A1

$$\therefore C = (2, 2^2 - 3(2) + 2) = (2, 0)$$

Method 2

Let
$$C = (x, y)$$

Gradient of $AB = 2 \times$ Gradient of AC

$$\frac{2-0}{0-1} = 2\left(\frac{y-2}{x-0}\right)$$

$$\therefore y = -x + 2$$

Solve simultaneously with $y = x^2 - 3x + 2$:

$$x^2 - 3x + 2 = -x + 2$$

$$x^2 - 2x = 0$$

$$x(x-2)=0$$

x = 0 (NA, since will get back point A) OR x = 2

$$\therefore C = (2, 2^2 - 3(2) + 2) = (2, 0)$$

A1

B1: Gradient of AB

simultaneously

M1: Formulating equations and solve

Answer (.....) [3]

20 It is given that
$$AB = \begin{pmatrix} -12 \\ a \end{pmatrix}$$
, $a > 0$ and $|BA| = 37$ units.

(a) Find the value of a.

$$|BA| = |AB| = 37$$

$$\begin{pmatrix} -12 \\ a \end{pmatrix} = 37$$

$$\sqrt{\left(-12\right)^2 + a^2} = 37$$

$$144 + a^2 = 1369$$

$$a^2 = 1225$$

$$a = 35 \text{ or } a = -35 \text{ (NA)}$$

M1: Distance formula

A1

Answer $a = \dots [2]$

(b) C is a point (0,10) and $\overrightarrow{AB} = \overrightarrow{DC}$. Find the coordinates of D.

 $\overrightarrow{AB} = \overrightarrow{DC}$ $\begin{pmatrix} -12 \\ 35 \end{pmatrix} = \overrightarrow{OC} - \overrightarrow{OD}$ $\overrightarrow{OD} = \overrightarrow{OC} - \begin{pmatrix} -12 \\ 35 \end{pmatrix}$

 $= \begin{pmatrix} 0 \\ 10 \end{pmatrix} - \begin{pmatrix} -12 \\ 35 \end{pmatrix}$ $= \begin{pmatrix} 12 \\ 10 \end{pmatrix}$

 \therefore Coordinates of D = (12, -25)

A1

M1: OC-OD

Answer (.....) [2]

(c) What type of quadrilateral is ABCD? Explain your answer.

AB = DC

There is a pair of parallel sides with equal length.


∴ ABCD is a parallelogram.

B1: Parallelogram

.....

.....[1]

The cumulative frequency curve shows the marks obtained by the 120 students in a recent Mathematics Test. The maximum mark is 100.



(a) Complete the grouped frequency table for the marks obtained.

| Marks obtained by students, x | Frequency |
|-------------------------------|-----------|
| $0 \le x < 20$ | 10 |
| $20 \le x < 40$ | 34 |
| $40 \le x < 60$ | 40 |
| $60 \le x < 80$ | 28 |
| $80 \le x < 100$ | 8 |

[1]

(b) Calculate an estimate of the mean mark.

Required estimates x < 20

$$= \frac{(10)(10) + (36)(54) + (50)(40) + (70)(28) + (90)(8)}{40 \le x < 60_{120}}$$

$$= 48\frac{1}{3} \text{ marks } 80 \le x < 100$$

B1: Accept 48.7 marks

Answermarks [1]

(c) Calculate an estimate of the standard deviation. Required estimate

$$= \sqrt{\frac{\sum fx^{2}}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^{2}}$$

$$= \sqrt{\frac{(10)^{2}(10) + (30)^{2}(34) + (50)^{2}(40) + (70)^{2}(28) + (90)^{2}(8)}{120}} - \left(48\frac{1}{3}\right)^{2}$$

$$= \sqrt{2780 - \left(48\frac{1}{3}\right)^{2}}$$

$$= 21.1 \text{ marks } (3 \text{ s.f})$$
A1

Answermarks [2]

(d) The passing mark is 50. Two students are chosen at random. Find the probability that both of them passed.

Within 10% of 50 means from 40 to 60.


Required probability =
$$\left(\frac{40}{120}\right)\left(\frac{39}{119}\right)$$

= $\frac{13}{119}$

B1

Answer [1]

The diagram shows the speed-time graph of a particle. 22



(a) Find the speed of the particle when t = 12s. Let the speed of the particle when t = 12 be v M1: Equating gradients

$$\frac{v-30}{12-6} = \frac{100-30}{13-6}$$

$$\therefore v = 90 \text{ m/s}$$

A1

Answerm/s [2]

(b) Find the distance covered before the particle starts to slow down. M1: Finding area under the graph using

Distance = Area under the graph from 0s to 13s

=
$$(6)(30) + \frac{1}{2}(30+100)(13-6)$$

= 635 m


A1

correct formula


Answer m [2]

(c) Complete the distance-time graph of the particle.

[1]



23



In the diagram, ABC and AED are straight lines and BE is parallel to CD.

(a) Show that triangle ABE is similar to triangle ACD. Give a reason for each statement you make.

 $\angle BAE = \angle CAD$ (Common angle)

 $\angle ABE = \angle ACD$ (Corresponding angles, BE parallel to CD)

 $\angle AEB = \angle ADC$ (Corresponding angles, BE parallel to CD)

 \therefore By AA similarity test, $\triangle ABE$ is similar to $\triangle ACD$

M1: Any two correct statements with complete reasons

A1: Conclusion with test

.....

.....[2

(b) Show that $\frac{BC}{ED} = \frac{AB}{AE}$. [2]

By similar Δs ,

$$\frac{AC}{AB} = \frac{AD}{AE}$$

$$\frac{AB + BC}{AB} = \frac{AE + ED}{AE}$$

$$\frac{AB}{AB} + \frac{BC}{AB} = \frac{AE}{AE} + \frac{ED}{AE}$$

$$1 + \frac{BC}{AB} = 1 + \frac{ED}{AE}$$

$$\therefore \frac{BC}{AB} = \frac{ED}{AE}$$

$$\therefore \frac{BC}{ED} = \frac{AB}{AE}$$

M1: Correct ratio with corresponding sides

A1 (AG)

A circle is drawn with points B, C and D on its circumference. It is further given that angle BCD + angle $BED > 180^{\circ}$. Explain why point E is inside the circle.


If point E is on the circumference of the circle, B1: Accept drawing to aid explanation BCDE will be a cyclic quadrilateral. Due to segments, angles in opposite $\angle BCD + \angle BED = 180^{\circ}$.

Hence, E is not on circumference of the circle.

As E moves further away from the center of circle, ∠BED will decrease in magnitude. Since $\angle BCD + \angle BED > 180^{\circ}$, E will be in the circle.

| |
|-----------|
| |
| |
| |
| ••••• |
| |
| F17 |
| |

24



OBC, ADC, AEB and OED are straight lines.

$$OA = \mathbf{a}$$
 and $OB = \mathbf{b}$.

$$OB = \frac{1}{3}OC$$
, $AE : EB = 3:2$ and $OE : ED = 1:1$.

(a) Find OE in terms of a and/or b as simply as possible.

$$OE = OA + AE$$

$$= OA + \frac{3}{5}AB$$

$$= OA + \frac{3}{5}(OB - OA)$$

$$= \mathbf{a} + \frac{3}{5}(\mathbf{b} - \mathbf{a})$$

$$= \frac{2}{5}\mathbf{a} + \frac{3}{5}\mathbf{b}$$

M1:
$$OA + \frac{3}{5}AB$$

A1

Answer: *OE* =[2]

(b) Find the value of $\frac{\text{area of } \triangle OBE}{\text{area of } BEDC}$

Let area of $\triangle OBE$ be x

∴ area of
$$\triangle OAE = \frac{3}{2}x$$
 (Ratio of bases)

∴ area of $\triangle OAB = \frac{5}{2}x$ and area of $\triangle AED = \frac{3}{2}x$ (Ratio of bases)

∴ area of
$$\triangle ABC = 2\left(\frac{5}{2}x\right) = 5x$$
 (Ratio of bases)

$$\therefore \frac{\text{area of } \triangle OBE}{\text{area of } BEDC} = \frac{x}{5x - \frac{3}{2}x} = \frac{2}{7}$$

M1: Using ratio of bases to find area of $\triangle OAE$


OR

Using ratio of bases to find area of $\triangle AED$ or $\triangle ABC$

A1

Answer[2]

25



AC is a diameter of a circle with centre O. $AC = \sqrt{2}r$ and AB = BC.

COB is a semi-circular arc, centre P.

What percentage of the circle is not shaded?

 $\angle ABC = 90^{\circ}$ (Right angle in semicircle)

B1: Need to show $\angle ABC = 90^{\circ}$

M1: Finding AB or BC

$$AB = BC$$

$$(AB)^2 + (BC)^2 = (\sqrt{2}r)^2$$

$$2(BC)^2 = 2r^2$$

$$(BC)^2 = r^2$$

 $\therefore BC = r \text{ (Reject -ve)}$

$$OC = \frac{1}{2}AC = \frac{\sqrt{2}}{2}r$$

$$CP = PO = \frac{1}{2}BC = \frac{r}{2}$$

$$OC^2 = CP^2 + PO^2$$

.: By converse of Pythagoras' Theorem,

Method 1: $\triangle OPC$ is a right angled \triangle .

Method 2:

B1: Need to show $\triangle OPC$ is a right angled \triangle

Note that P and O are midpoints

of BC and AC respectively.

.. By midpoint theorem,

OP is parallel to AB,

and $\angle OPC = \angle ABC$

 $\angle ABC = 90^{\circ}$ (Right angle in semicircle)

 $\therefore \angle OPC = 90^{\circ}$

 $\therefore \triangle OPC$ is a right angled \triangle .

Percentage of circle shaded

$$= \frac{\text{Sector OPC} - \text{Triangle OPC}}{\text{Area of big circle}} \times 100\%$$

$$= \frac{\frac{1}{2} \left(\frac{r}{2}\right)^{2} \left(\frac{\pi}{2}\right) - \frac{1}{2} \left(\frac{r}{2}\right)^{2} \sin\left(\frac{\pi}{2}\right)}{\pi \left(\frac{\sqrt{2}}{2}r\right)^{2}} \times 100\%$$

$$=\frac{\frac{1}{16}\pi r^2 - \frac{1}{8}r^2}{\frac{1}{2}\pi r^2} \times 100\%$$

$$=\frac{\frac{1}{16}\pi - \frac{1}{8}}{\frac{1}{2}\pi} \times 100\%$$

= 4.542252845%

Percentage of circle shaded = 100% - 4.542252845%

$$=95.5\%$$
 (3 s.f)

M1: Using midpoint theorem

A1: $\angle OPC = \angle ABC$

B1: Need to show $\triangle OPC$ is a right angled \triangle

M1: Correct percentage formula with the relevant shapes found using the correct formula (FT)

A1

Answer% [5]





JURONG SECONDARY SCHOOL 2022 GRADUATION EXAMINATION 2 SECONDARY 4 EXPRESS/ SECONDARY 5 NORMAL ACADEMIC

| NAME NAME | MARKING SCHEME | | |
|-----------|----------------|-----------------|------|
| CLASS | | INDEX NUMBER | |
| MATHEMA | ΓICS | 4048 | 3/02 |
| PAPER 2 | | 24 August | 2022 |

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all the questions.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.


If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For π , use either your calculator value or 3.142, unless the question requires the answer in terms of π .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 100.



2 hour 30 minutes

This document consists of 26 printed pages including this page.

| 1 | (a) | (i) | 3 | B1 |
|---|-----|----------------------------------|-------------------------------------|--------------------------|
| | | | $\frac{3}{2}$ | |
| | | (ii) | $y = \frac{5x - 2}{3x + 1}$ | |
| | | | | |
| | | | 3xy + y = 5x - 2 | |
| | | | 5x - 3xy = y + 2 | M1 – simplify + group |
| | | | x(5-3y) = y+2 | group |
| | | | $x = \frac{y+2}{5-3y}$ | A 1 |
| | | | 5-3y | A1 |
| | (b) | $\frac{x-1}{x}$ | $\frac{5}{2} - \frac{x+3}{2} = -4$ | |
| | | 3 | 2 | |
| | | $\frac{2(x)}{x}$ | $\frac{-5)-3(x+3)}{6} = -4$ | M1 – same denominator |
| | } | 2~- | -10 - 3x - 9 = -24 | denominator |
| | | x = | | A1 |
| - | (c) | | -y=23 | |
| | | i | 23-2x | |
| | | 1 - | -2(23-2x)=10 | M1 - |
| | | | 46+4x=10 | substitution |
| | | 7x = | = 56 | |
| | | x = | | A1 |
| | | y = | | Al |
| - | (d) | 1 - | x^2-9 | |
| | | $\frac{1}{6x^2}$ | $\frac{x}{-x-12}$ | M1 – numerator |
| | | | • | M1 – |
| | | $=\frac{1}{(2)}$ | $\frac{(2x-3)(2x+3)}{(2x-3)(3x+4)}$ | denominator |
| | | | | |
| | | $\int_{0}^{\infty} \overline{3}$ | $\frac{x+3}{x+4}$ | Al |
| | | | | Total: 11 marks |

| 2 | (a) | (28 24) | B1 |
|---|-----|--|------------------------------------|
| | (b) | $\begin{array}{c} (16 12) \\ \hline (70) \end{array}$ | B1 |
| | (c) | (3880) | B1 |
| | | 2080 | |
| | (d) | The elements represent the total amount Mrs Teo collected in a month from the weekday and weekend sessions respectively. | B1 |
| | (e) | Total amount of money per session $=$ $\begin{pmatrix} 10 & 8 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} 70 \\ 80 \end{pmatrix} = \begin{pmatrix} 1340 \\ 370 \end{pmatrix}$ | M1 |
| | | Total after 10% increase = $(1 \ 1.1) \binom{1340}{370} = (1747)$ | M1 |
| | | $1747 \times 4 \times 3 = 20964 | M1 A1 |
| | | | Accept non- matrices working |
| | | | Total: 8 marks |

| 3 | (a) | $\angle XCD = \frac{360^{\circ}}{18} = 20^{\circ}$ | B1 |
|---|--------------------|--|------------------|
| | (b) | $\angle XCD = \angle XDC$ $\angle CXD = 180^{\circ} - 20^{\circ} - 20^{\circ} \text{ (isos. } \Delta)$ | |
| | | =140° | A1 (with reason) |
| | (c) | BC = ED (sides of regular polygon) | M1 – evidence |
| | | XC = XD (given) $\therefore BC + CX = ED + DX$ | A1 |
| | | BX = EX | |
| | (d) | $\angle BXE = \angle CXD \text{ (common)}$ $\angle XBE = \frac{180^{\circ} - 140^{\circ}}{2} = 20^{\circ}$ | M1 – evidence |
| | | $\therefore \angle XCD = \angle XBE$ $\Delta XCD \text{ and } \Delta XBE \text{ are similar (AA)}$ | A1 |
| - | (e) | They are parallel. | B1 |
| | ` ´ - | | Total: 7 marks |

| 4 | (a) | $T_5 = 9^2 - 34 = 47$ | B1 (with |
|---|-----|---|------------------------|
| | ļ | | working) |
| | (b) | $T_n = (n+4)^2 - 18 - 4(n-1)$ | M1 - n+4 and |
| | | $= n^2 + 8n + 16 - 18 - 4n + 4$ | 18+4(n-1) |
| | | $=n^2+4n+2$ | A1 |
| | (c) | $n^2 + 4n + 2 = 962$ | |
| | | $n^2 + 4n - 960 = 0$ | |
| | | $n = \frac{-4 \pm \sqrt{4^2 - 4(1)(-960)}}{2(1)}$ | M1 – find value |
| | | | of n |
| | | = 29.048 or -33.048 | . |
| | | Since n is not an integer, 962 is not a term of the sequence. | A1 |
| | (d) | $T_n - T_{n-1}$ | |
| | | $= (n^2 + 4n + 2) - [(n-1)^2 + 4(n-1) + 2]$ | M1 – correct |
| | | $= n^2 + 4n + 2 - (n^2 - 2n + 1 + 4n - 4 + 2)$ | expression |
| | | $= n^2 + 4n + 2 - n^2 + 2n - 1 - 4n + 4 - 2$ | M1 – correct expansion |
| | | =2n+3 | Aĺ |
| | (e) | The difference between two consecutive terms is $2n+3$. | B1 – always |
| | | Since $2n$ is always even for all values of n , the sum of an even integer and an odd | even + add to |
| | | constant 3 will always give an odd integer. | odd |
| | | | Total: 8 marks |

| 5 | (a) | p = 2.2 | | B1 | |
|---|--------------|----------------------|---|--|--|
| | (b) | Refer to 1 | ast page | G3 | |
| | (c) | The graph | The graph of $y = x + \frac{5}{x^2} - 3$ does not meet the line $y = 0/\text{does not have any } x$ - | | |
| | | | s/ the minimum y-value is, which is more than 0 | D1 | |
| | (d) | Draw tan Gradient | $=-1.96\pm0.7$ | B1 B1 | |
| | (e) | (i) | Draw line with gradient 2 passing through (0.5,2) (y-intercept = 1) | B1 - passes through (0.5,2) B1 - correct gradient (& y- int) | |
| | | (ii) | v = 2x + 1 | Bí | |
| | | (iii) | $y = 2x+1$ $x + \frac{5}{x^2} - 3 = 2x+1$ $\frac{5}{x^2} = x+4$ $5 = x^3 + 4x^2$ | M1 – sim. eqn. | |
| | | | $ \begin{array}{c} x^{2} \\ 5 = x^{3} + 4x^{2} \\ x^{3} + 4x^{2} - 5 = 0 \end{array} $ | A1 — final eqn | |
| | | (iv) | x = 1 | B1 (ecf – accept answer from intersection of curve and line) Total: 13 marks | |
| | 1 | | | | |

| 6 | (a) | OS = | OS = OR (radii of smaller circle) | | | | |
|---|-----|------------|--|--------------------|--|--|--|
| | | ∠PC | $\angle POS = \angle QOR \text{ (vert. opp. } \angle \text{)}$ | | | | |
| | | OP = | OP = OQ (radii of larger circle) | | | | |
| | | ΔI | $^{\circ}SO$ is congruent to ΔQRO (SAS) | A1 | | | |
| | | Alter | Alternatives: RHS, AAS | | | | |
| | (b) | (i) | $\angle ORQ = 90^{\circ} \text{ (tan } \perp \text{ rad)}$ | | | | |
| | | | $QR = \sqrt{8^2 - 4.36^2}$ | M1 – with evidence | | | |
| | | | = 6.7075 | | | | |
| | | | Area of $\triangle QRO = \frac{1}{2}(4.36)(6.7075)$ | M1 | | | |
| | | | =14.62235 | | | | |
| | | | $\approx 14.6 \text{ cm}^2$ | A1 | | | |
| | | (ii) | Area of shaded minor sector | 1.51 | | | |
| | | | $=\frac{1}{2}(4.36)^2(0.995)$ | M1 | | | |
| | | | =9.457276 | | | | |
| | | | Area of ring between the circles | | | | |
| | | | $=\pi(8)^2-\pi(4.36)^2$ | M1 | | | |
| | | | =141.3415 | | | | |
| | | | Area of unshaded in $\triangle QRO$ | | | | |
| | | | =14.62235 - 9.457276 | M1 | | | |
| | | | = 5.165074 | IVII | | | |
| | | | Shaded area | | | | |
| | | | = 141.3415 + 2(9.457276) - 2(5.165074) | | | | |
| | | | =149.925904 | | | | |
| | | | $\approx 150 \text{ cm}^2$ | A1 | | | |
| | 1 | | | Total: 9 marks | | | |

| 7 (a) |) | $\frac{5}{4+4+5} \times 300 \times \$148 = \$18500$ | M1 A1 |
|-------|-------|--|---------------------------------------|
| (b) |) (i) | $\frac{300}{x}$ min | B1 |
| | (ii) | $\frac{300}{x-2} \min$ | B1 |
| | (iii) | $\frac{300}{x-2} - \frac{300}{x} = 3$ $\frac{300x - 300(x-2)}{x(x-2)} = 3$ | M1 – form eqn M1 – combine fractions |
| | (iv) | $x^{2}-2x-200 = 0 \text{ (shown)}$ $x = \frac{-(-2) \pm \sqrt{(-2)^{2}-4(1)(-200)}}{2(1)}$ | A1 M1 |
| | | $2(1)$ =15.17744 or -13.17744 $\approx 15.2 \text{ or } -13.2$ | A1 (3sf, no rej.) |
| | (v) | Reject -13.2 x = 15.2 $15.17744 \times 60 = 910.6464$ | M1 |
| | | ≈910 | A1 (round down) (Reject 3 x 60) |
| | | | Total: 11 marks |

| 8 (a | 1 · · · · · · · · · · · · · · · · · · · | |
|------|--|------------------|
| | $\frac{OA}{\sin 30^{\circ}} = \frac{8}{\sin 120^{\circ}}$ | M1 – sine rule |
| | sin 30° sin 120° | |
| | $OA = \frac{8}{\sqrt{\sin 30^\circ}}$ | |
| | $OA = \frac{8}{\sin 120^{\circ}} \times \sin 30^{\circ}$ | |
| | ≈ 4.6188 cm | |
| | = 4.619 cm (to 3 d.p.) | A1 |
| | - 4.019 cm (to 3 d.p.) | |
| | OR | |
| | | |
| | 2 /2 /2 | |
| | $OA = \frac{2}{3}\sqrt{8^2 - 4^2}$ | M1 |
| | = 4.61880 | |
| | ≈ 4.619 | A1 |
| (b | | |
| ` | Base area $\triangle ABC = \frac{1}{2}(8)(8)\sin 60^{\circ}$ | M1 – base |
| | $\approx 27.7128 \text{ cm}^2$ | |
| | Height of $\triangle AXB$, $l = \sqrt{18^2 - 4^2}$ | M1 – lateral |
| | reight of $\triangle AAB$, $t = \sqrt{16} - 4$ $\approx 17.5499 \text{ cm}$ | height |
| | ≈ 17.3499 cm Total surface area | 8 |
| | | |
| | $=3\times\frac{1}{2}(8)(17.5499)+27.7128$ | |
| | $\approx 238.31 \text{ cm}^2$ | |
| | $= 238 \text{ cm}^2 \text{ (to 3 s.f.)}$ | A1 |
| (c | $5^2 + 4^2 - 3.5^2$ | |
| | $\cos \angle PQR = \frac{5^2 + 4^2 - 3.5^2}{2(5)(4)}$ $\angle PQR = \cos^{-1}\left(\frac{23}{32}\right)$ | M1 – cosine rule |
| | (23) | |
| | $\angle PQR = \cos^{-1}\left(\frac{2S}{22}\right)$ | |
| | | A1 |
| (4 | $= 44.0^{\circ} \text{ (to 1 d.p.)}$ $AQ = 6 \text{ cm}$ | |
| | $OQ = \sqrt{4.619^2 + 6^2 - 2(4.619)(6)\cos 30^\circ}$ | |
| | | M1 – find OQ |
| | ≈ 3.055 cm | |
| | $XO = \sqrt{18^2 - 4.619^2}$ | |
| | ≈ 17.397 cm | M1 – |
| | $\tan \theta = \frac{17.397}{3.055}$ | Pythagoras' thm |
| | | |
| | Angle of elevation of X from Q | |
| | $=\tan^{-1}\left(\frac{17.397}{3.055}\right)$ | Mi |
| | | |
| | $= 80.0^{\circ} \text{ (to 1 d.p.)}$ | A1 |
| | | Total: 11 marks |

| | (a) | (i) | $100 < x \le 120$ | B1 |
|----------|--------------|----------|--|------------------------|
| | | (ii) | 22% | B1 |
| | | (iii) | 809000 ,6280,2 | M1 – working |
| | | | S.D. = $\sqrt{\frac{809000}{50} - (\frac{6280}{50})^2}$ | (show substitution) |
| | | | = 20.11566 | substitution) |
| | | | | A1 |
| | | | ≈ 20.1 | |
| | | (iv) | The blood pressures of the second group of patients is less consistent than that of the first group of patients, since it has a SD of 22.1 which is higher than the SD | B1 – compare SD and |
| | | | of the first group (20.1) | comment on |
| |] | | of the first group (20.1) | consistency |
| | (bi) | | | B2 |
| | | | blue | -1 for each |
| | | | | wrong pair of |
| | | | blue | branches |
| | | |) blue | |
| | | | $\frac{2}{8}$ white | |
| | | | Winte | |
| | | | white | |
| | | | | |
| ļ | | < | | |
| | | | | |
| | | | blue | |
| | | | blue < | |
| | | | | |
| | | | white white | |
| | | | | |
| | | | | |
| | | | white | |
| <u> </u> | | (ii) | (a) 6 5 2 1 4 | M1 – either both |
| | | (11) | $\binom{(a)}{8} \left(\frac{6}{8} \times \frac{5}{7} \right) + \left(\frac{2}{8} \times \frac{1}{7} \right) = \frac{4}{7}$ | W or both B |
| | | | | A1 |
| | | | , | 777.6 |
| | | | | FT from tree |
| | | <u> </u> | (1) (2 (1) (6 2 5) | diagram M1 – either |
| | | | (b) $\left(\frac{2}{8} \times \frac{6}{7} \times \frac{1}{6}\right) + \left(\frac{6}{8} \times \frac{2}{7} \times \frac{5}{6}\right)$ | WBW or all |
| | | | (8 7 6) (8 7 6) | BWB |
| | | | 3 | A1 |
| | | | $=\frac{1}{14}$ | |
| | | 1 | | Total: 11 marks |

| 10 | (a) | $3.42 \times 2.7 = 9.234 \text{ m}^2$ | B1 Reject 3s.f. |
|----|-----|--|---------------------------------------|
| | (b) | 10 cm represents 168 cm | M1 – measure |
| | (0) | 1 cm represent 16.8 cm | distance from |
| | | Elbow to ground = 6.4 cm (accept 6.3cm and 6.5cm) | elbow to ground |
| | | Elbow to ground – 0.4 cm (accept 0.5cm and 0.5cm) | eloow to glound |
| | | $6.4 \times 16.8 = 107.52$ cm | |
| | | Range of optimal kitchen cabinets height | |
| | | 107.52 - 10 = 97.52 cm to 107.52 cm | A1 Measured 6.3 cm, 95.84 cm to |
| | | | 105.84 cm |
| 1 | | | Measured 6.5 |
| | | | cm, 99.2 cm to |
| | | | 109.2 cm |
| | (c) | Kitchen Cabinet | |
| | | Compulsory: | |
| | | Sink + corner cabinet + wire basket shelves | · |
| | | Cabinet 1 + Cabinet 2 (must go at the junction of the wall) + Cabinet 5 | M1 (Cabinet |
| | | Remaining length = $342-50-60-50=172$ cm | 1+2+5) |
| | | Remaining width = $135-60=75$ cm | M1 (remaining |
| | | Current total cost = $$156 + $166 + $166 = 488 | space) |
| | | Along the length – 1 Cabinet 3 + 1 Cabinet 4 (90 cm + 80 cm) | M1 |
| | | Along the width – 1 Cabinet 6 (70 cm) | (combination of |
| | | in order to optimise storage | cabinets) |
| | | | M1 – total cost |
| | | Total cost of cabinets = $$488 + $193 + $260 + $290 = 1231 | of cabinets |
| | | Height customisation = $$25 \times 6 = 150 | M1 – height |
| | | Remaining budget = $$2200 - $1231 - $150 = 819 | M1 – budget + |
| | | Ceramic or acrylic for materials | conclusion for |
| | | | type of material |
| | | Mr Graham should choose the Ceramic counter top and his kitchen cabinet | |
| | | combination is one of cabinets 1, 2, 3, 4, 5, 6 with \$19 remaining in his budget. | A1 (relate to |
| | ļ | | budget) |
| | | | Total: 10 marks |