



--

2	0	J		
---	---	---	--	--

S				
---	--	--	--	--

--	--	--	--

9729/03

21 September 2021

2 hours

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions in the spaces provided on the Question Paper. If additional space is required, you should use the pages at the end of this booklet. The question number must be clearly shown.

Section A

Answer **all** the questions.

Section B

Answer **one** question.

A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use								
Question Number	1	2	3	4	5	s.f.	units	Total
Marks	20	22	18	20	20			80

This document consists of **32** printed pages and **0** blank page.

Section A

Answer **all** the questions in this section.

- 1 In a 2014 paper published in the Journal of Agricultural and Food Chemistry, Hendon and Colonna-Dashwood discovered the effect of water hardness on coffee flavour. Compounds in hard water tend to attach to the flavourful elements in roasted coffee beans during brewing. Water with higher levels of magnesium will likely extract more flavour from a coffee bean.

Water described as "hard" is high in concentration of Total Dissolved Solids (TDS), specifically calcium and magnesium. The hardness of water may be reported in parts per million (ppm). The solute concentration of a dilute aqueous solution in units of mg dm^{-3} is called parts per million, or ppm.

Classification	ppm
Soft	0 – 17.1
Slightly hard	17.1 – 60.0
Moderately hard	60.0 – 120
Hard	120 – 180
Very hard	> 180


- (a) In a sample of Singapore's tap water, the concentration of magnesium and calcium ions present are found to be $5.97 \times 10^{-5} \text{ mol dm}^{-3}$ and $5.49 \times 10^{-4} \text{ mol dm}^{-3}$ respectively. These two ions can be separated by selective precipitation with potassium hydroxide. The numerical values of solubility product of magnesium hydroxide and calcium hydroxide at 25°C are 1.50×10^{-11} and 5.50×10^{-6} respectively.

- (i) Calculate the total concentration of magnesium and calcium ions in ppm, and hence classify the hardness of water in this sample of tap water. [2]
- (ii) Calculate the minimum pH of the solution at which the magnesium ion precipitates as magnesium hydroxide. [2]
- (iii) The magnesium hydroxide continues to precipitate out of the solution as potassium hydroxide is being added continuously. Eventually, the concentration of the hydroxide becomes high enough to precipitate the calcium ions as well.

What is the concentration of magnesium ions when calcium ions begin to precipitate?

[2]

- (b) A balanced extraction is a well-brewed cup of coffee that is aromatic and rich in flavours. Eugenol is a flavour note with a “woody” taste found in coffee, wine and whisky.



Eugenol

Like other alkenes, it undergoes hydrohalogenation when treated with hydrogen halides.

- (i) Draw a labelled diagram showing the orbital overlap between the carbon atoms C1 and C2 and state the hybridisation involved.

Do **not** include other atoms.

[2]

- (ii) Hydrohalogenation of unsymmetrical alkenes results in a mixture of products. In such cases, the major product can be predicted using Markovnikov's rule. Describe the mechanism of the reaction between eugenol and hydrogen chloride.

You may represent eugenol using R_1 .

[2]

- (iii) With reference to your mechanism in (b)(ii), explain why the major product is formed.

[2]

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

- $$\begin{array}{c} \text{R} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array} \xrightarrow[\text{Step 1}]{\text{BH}_3} \begin{array}{c} \text{H} \quad \text{BH}_2 \\ | \quad | \\ \text{R}-\text{C}-\text{C}-\text{H} \\ | \quad | \\ \text{H} \quad \text{H} \end{array} \xrightarrow[\text{Step 2}]{\text{H}_2\text{O}_2} \begin{array}{c} \text{H} \quad \text{OH} \\ | \quad | \\ \text{R}-\text{C}-\text{C}-\text{H} \\ | \quad | \\ \text{H} \quad \text{H} \end{array}$$

It is suggested that the mechanism goes through the formation of the intermediate below.



- Suggest the mechanism showing the formation of the intermediate given above when eugenol reacts with borane. Indicate clearly the polarity of the B–H bond in borane by drawing $\delta+$ and $\delta-$ on the appropriate atoms.

[1]

- [illegible]

- The table shows the results of experiments in which the halogens \mathbf{X}_2 , \mathbf{Y}_2 and \mathbf{Z}_2 were added to separate solutions containing \mathbf{X}^- , \mathbf{Y}^- and \mathbf{Z}^- ions.

	X⁻(aq)	Y⁻(aq)	Z⁻(aq)
X₂	no reaction	no reaction	no reaction
Y₂	X₂ formed	no reaction	Z₂ formed
Z₂	X₂ formed	no reaction	no reaction

With reference to the table above, identify the halogens **X**, **Y** and **Z**. Explain your reasoning. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (e) A glass rod was heated in a Bunsen burner flame and placed into a jar of hydrogen chloride gas. The experiment was repeated using a jar of hydrogen iodide gas. A colour change was observed in one of the samples.

Using relevant data from *Data Booklet*, explain these observations.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

[Total: 20]

- 2 Nitrous acid, HNO_2 , is a weak monoprotic acid. HNO_2 is unstable and decomposes readily.

It can be prepared by acidification of aqueous solutions of potassium nitrite with a mineral acid. The acidification is usually conducted at low temperatures, and the HNO_2 is consumed *in situ*.

An equilibrium exists as follows.



$$K_c = 1.66 \times 10^3 \text{ mol}^{-1} \text{ dm}^3$$

- (a) Calculate the value of the acid dissociation constant, K_a , of HNO_2 . [1]

.....

.....

.....

.....

.....

.....

.....

.....

- (b) A solution of HNO_2 was prepared by mixing equal volumes of 0.40 mol dm^{-3} $\text{HCl}(\text{aq})$ with 0.40 mol dm^{-3} $\text{KNO}_2(\text{aq})$.

Calculate the pH of this solution.


[2]

.....

.....

-
-
-
-
-
-
-
-
- (c) Calculate the pH of the aqueous mixture when 10.0 cm³ of 0.400 mol dm⁻³ HCl(aq) is added to 30.0 cm³ of 0.400 mol dm⁻³ KNO₂(aq). [2]

-
-
-
-
-
-
-
-
-
- (d) Gaseous nitrous acid decomposes into nitrogen dioxide, nitric oxide, and water:



- (i) Draw the dot-and-cross diagram of NO₂.
Explain the difference in bond angles in H₂O and NO₂. [2]
- (ii) Calculate the increase in pressure when 2.00 g of HNO₂ decomposes under 1 atm and 150 °C in a 1 dm³ container. [2]
-
-


- (e) Similar to HNO_2 , 3-chloropropanoic acid is a weak monobasic organic acid. Compounds **E** and **F** are constitutional isomers of 3-chloropropanoic acid. When the same amount of 3-chloropropanoic, **E** and **F**, is each added to separate portions of water of equal volume, the three solutions obtained have pH values as shown in the table below.

compound	pH of aqueous solution
3-chloropropanoic acid	2.3
E	1.9
F	1.0

Suggest structural formulae for compounds **E** and **F** and hence, account for the difference in pH values of the three solutions obtained. One of the compounds has the structure $\text{R}-\text{O}-\text{R}$.

[4]

- (f) Hexanedioic acid, a diprotic acid, can be synthesised from succindialdehyde by the following synthesis pathway.



- (i) Suggest reagents and conditions for each of the Steps 1, 2 and 3. [3]
- (ii) Suggest the structures for **K** and **L**. [2]

- (g) Carbonyl compounds can undergo the Aldol reaction under basic conditions. The mechanism is shown below.

Step 1	
	One of the hydrogen atoms, bonded to the carbon next to the carbonyl carbon, forms a bond with a hydroxide ion via an acid–base reaction.
Step 2	
	Nucleophilic attack by enolate anion.
Step 3	
	Acid–base reaction.
$R_1, R_2, R_3 = \text{H, alkyl or aryl}$	

- (i) Suggest the structure of the compound formed from the Aldol reaction between 1 molecule of succindialdehyde and 1 molecule of methanal. [1]

(ii) Compound **M** is a compound with 8 carbon atoms and undergoes the following reactions.

- **M** gives an orange precipitate with 2,4-DNPH.
- **M** does not react with Tollens' reagent.
- **M** reacts with alkaline aqueous iodine to give a yellow precipitate and product **N**, $C_7H_5O_2^-$.
- **M** undergoes Aldol reaction to form **O**, $C_{16}H_{16}O_2$, under basic conditions.

Suggest the structures for **M**, **N** and **O**.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

[Total: 22]

- 3 (a) The following scheme illustrates a series of reaction involving $\text{Fe}^{3+}(\text{aq})$.



- (i) Define transition element. [1]
- (ii) State the type of reaction which occurred in reaction I and write an equation for the reaction. [2]
- (iii) Write an equation for reaction II. [1]
- (iv) Given that $\text{C}_2\text{O}_4^{2-}$ is a bidentate ligand, draw the structural formula of the complex formed. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Ferromanganese is an alloy added to steels to improve their mechanical properties. A 15.0 g sample of ferromanganese was dissolved in 250 cm³ of dilute sulfuric acid to give an solution containing iron(II) sulfate and manganese(II) sulfate. 25.0 cm³ of the resulting solution required 20.0 cm³ of 0.0360 mol dm⁻³ of potassium manganate(VII) solution for complete reaction.

(i) By using the *Data Booklet*, construct a balanced equation for the reaction between resulting solution and potassium manganate(VII) solution. [1]

(ii) Calculate the percentage by mass of iron in ferromanganese. [3]

-
-
-
-
-
-
-
- (c) Neutral FeCl_3 remains yellow when reacted with compound **S**, $\text{C}_{11}\text{H}_{10}\text{O}_3$. **S** gives a silver mirror with silver diammine complex. However, **S** does not give a precipitate with hot alkaline Cu^{2+} solution.

S is heated with acidified KMnO_4 for several hours to give benzene-1,2,3-tricarboxylic acid as one of the organic products. When heated with $\text{NaOH}(\text{aq})$ followed by acidification, **S** forms **T**, $\text{C}_{11}\text{H}_{12}\text{O}_4$. **T** reacts with alkaline aqueous iodine to give a yellow precipitate. **S** reacts with NaBH_4 to give **U**, $\text{C}_{11}\text{H}_{12}\text{O}_3$.

Suggest the structures for **S**, **T** and **U**. Explain the reactions described.

[9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[Total : 18]

Section B

Answer **one** question from this section.

- 4 (a) Alkaline earth metals, also known as Group 2 elements, are highly metallic and are good conductors of electricity. They have a grey-white lustre when freshly cut but tarnish readily in air.

(i) Describe and explain the trend in thermal stability of the Group 2 carbonates.

[2]

(ii) When ozone (O_3) is passed over dry powdered BaO at $-10.0^\circ C$, barium ozonide, $Ba(O_3)_2$, is formed as a red-brown solid.

Adding water to the solid and warming to room temperature causes a reaction to occur. Oxygen gas is produced and an alkaline solution is left.

Write a balanced equation for the reaction between barium ozonide and water.

[1]

(iii) Suggest a suitable temperature to produce calcium ozonide by passing ozone through powdered CaO.

[1]

.....

.....

.....

.....

.....

.....


.....

.....

.....


.....

- (b) Ozone is usually produced by passing oxygen gas through two highly-charged electrical plates.




The reaction does not go to completion and a mixture of gases will be produced.

The concentration of O_3 in the mixture can be determined by its reaction with aqueous KI.




The iodine formed can be estimated by its reaction with sodium thiosulfate.




When 300 cm^3 of an oxygen/ozone gaseous mixture at s.t.p. was passed into an excess of aqueous KI, and the iodine formed was titrated against $\text{Na}_2\text{S}_2\text{O}_3$. 24.0 cm^3 of $0.100 \text{ mol dm}^{-3}$ $\text{Na}_2\text{S}_2\text{O}_3$ was required to discharge the iodine colour.

- (i) Suggest a suitable indicator can be used in the titration and state the colour change at end point. [1]
- (ii) Calculate the percentage of O_3 in the gaseous mixture. [2]


- (c) Ozone adds rapidly to alkenes at low temperature to give cyclic intermediates, called molozonides. Once formed, molozonides then rapidly rearranges to form ozonides. The reaction scheme is shown below.



The reaction can also be expressed as



- (i) State the type of reaction occurring in Step I of the reaction scheme. [1]
- (ii) Suggest the role of zinc in the reaction scheme. [1]
- (iii) β -Damascenone is a chemical compound found in whisky, which is an alcoholic liquor. Also known as rose ketones, β -damascenone is a key compound that contributes a floral note to whisky.



[3]

β -damascenone

Predict the organic products formed when β -damascenone reacts with ozone as shown by the reaction scheme above.

This image shows a full page of a handwriting practice worksheet. It consists of multiple rows of horizontal dotted lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no margins or additional markings.

(d) Apart from β -damascenone, whisky contains an array of compounds that affect its taste and flavour, which include phenolic compounds, aldehydes and esters. The use of different grains, distillation process and wood used in the ageing process can also change the flavour profile.

- (i) Phenolic compounds in general contributes smoky flavours and bitterness in whisky. In Scotch whisky, the use of peat fires to dry the barley grains creates a class of medicinal-smelling compounds known as cresols.


Cresols exists as constitutional isomers with the formula C_7H_8O . When added to aqueous bromine, rapid decolourisation followed by the formation of a white precipitate is observed for all isomers of cresol.

Draw all the possible isomers of cresol.

[3]

- (ii) Whisky lactones, such as trans-3-methyl-4-octanolide, are responsible for the woody, spicy and coconut flavour notes.

A student wanted to synthesise an amide from this lactone.



trans-3-methyl-4-octanolide


The following steps were proposed.

1. heat with dilute H_2SO_4
2. addition of ethylamine


Comment on the feasibility of this reaction scheme.

[2]

- (iii) Esters in whisky can also contribute fruity flavours such as ethyl hexanoate, which imparts a sweet apple flavor, and isoamyl acetate, which gives a banana aroma.



ethyl hexanoate



isoamyl acetate

[2]

Describe a simple chemical test that can be used to distinguish ethyl hexanoate from isoamyl acetate.

State any observations you would make with each compound.

- (iv) Whisky is a mixture of water, ethanol and organic compounds like whisky lactones. When the alcoholic whisky is stored for a long period of time, as the ethanol content decreases, a cloudiness can be observed in the whisky.

Suggest a reason why the cloudiness occurs.

[1]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

.....

.....

.....

.....

.....

.....

.....

.....

[Total: 20]

- 5 Cyclopropane is an explosive, colourless gas that was discovered by August Freund in 1881, and was once used as a general anesthetic in clinical practice.

(a) Cyclopropane can undergo isomerisation to propene at 298 K.




Table 5.1

	$\Delta H / \text{kJ mol}^{-1}$
standard enthalpy change of formation of $\text{CO}_2(g)$	-394
standard enthalpy change of formation of $\text{H}_2\text{O}(l)$	-286
standard enthalpy change of combustion of cyclopropane	-2091

Using the data given above and in Table 5.1, construct a suitable energy cycle and calculate the enthalpy change of formation of propene at 298 K.

[3]

.....

.....

.....

.....

.....

.....

(b) 100 cm³ of a mixture of methane and cyclopropane was completely burnt in 300 cm³ of oxygen and then cooled to room temperature.

The total volume of the gaseous reaction mixture decreases by 215 cm^3 .

On passing the resultant gaseous mixture through potassium hydroxide, the final volume was found to be 25.0 cm^3 .

What is the ratio of methane to cyclopropane in the original mixture?

[2]

.....

.....

.....


.....

.....

.....

.....

Cyclopropane can undergo free radical substitution with chlorine, similar to an aliphatic alkane. Chlorocyclopropane is the major product formed under a certain set of conditions.




- (c) (i) State the conditions needed to produce chlorocyclopropane as the major product. [1]
- (ii) Describe the mechanism for this reaction. [3]

Trichlorocyclopropane, $\text{C}_3\text{H}_3\text{Cl}_3$, is one of the possible polysubstituted by-products of the reaction under a different set of conditions.

This tri-substituted cycloalkane can exist as 3 constitutional isomers, all of which exhibit stereoisomerism.

- (iii) Define the term *stereoisomerism*. [1]
- (iv) Draw the structures of the 3 isomers of trichlorocyclopropane and suggest the type of stereoisomerism associated with each of them.


Note: You may refer to the diagram of cyclopropane shown below to draw the isomers.



[4]


- (d) However, in the absence of ultraviolet light, cyclopropane can undergo addition reactions similarly as alkenes. This leads to the opening of the ring structure.

For example,




- (i) Suggest a possible reason why cyclopropane can undergo addition reactions, unlike aliphatic alkanes. [1]
- (ii) Propane can be produced from cyclopropane in a three-step synthesis as shown below.

Suggest the reagents and conditions you would use in steps 2 and 3, and identify the intermediates **P** and **Q**.



[4]

- (e) A possible amino acid derivative of cyclopropane is shown below.



Draw the structure of the zwitterion formed by this amino acid derivative.

[1]

[Total: 20]

Additional answer space

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

.....

.....

.....

.....

.....

.....

.....

.....

.....