ANGLO-CHINESE JUNIOR COLLEGE DEPARTMENT OF CHEMISTRY **Preliminary Examination**

CHEMISTRY **Higher 2**

9729/03

Paper 3

18 August 2017 2 hours

Additional Materials: Writing Paper Data Booklet **Cover Page**

READ THESE INSTRUCTIONS FIRST

Write your index number and name, form class and tutorial class on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, glue or correction fluid.

Answer all questions in Section A. Answer either question 4 or 5 in Section B. Start each question on a new sheet of writing paper.

A Data Booklet is provided.

The use of an approved calculator is expected, where appropriate. You are reminded of the need for good English and clear presentation in your answers.

At the end of the examination, fasten all your work securely behind a cover sheet. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 17 printed pages, including this cover page.

9729/03/Prelim/17

© ACJC 2017



ANGLO-CHINESE JUNIOR COLLEGE



Section A – Answer **ALL** questions. Begin each question on a fresh piece of writing paper.

- 1 This question is about Period 3 and Group 2 elements.
 - Phosphorus sulfide, P_4S_3 , is used in small amounts in the tip of a matchstick. (a) On striking a matchstick, this compound burns to form sulfur dioxide and phosphorus pentoxide.
 - (i) Write the equation for this reaction.
 - (ii) The melting points of the two oxides formed in (a)(i) differ significantly from that of silicon(IV) oxide (SiO₂). Account for this difference in terms of structure and bonding of each oxide.
 - (iii) The solid oxide formed in (a)(i) dissolves in water to give an acidic solution.

Write the equation for this reaction and give an approximate pH of the solution formed.

- (b) A Group 2 nitrate exists as a hydrate $M(NO_3)_2 \times H_2O$. On heating, 1.80 g of this hydrate lost 0.55 g in the form of steam, as it was converted into the anhydrous nitrate, $M(NO_3)_2$. Subsequent heating to constant mass produced a white residue and mixture of two gases. The gases produced were passed through aqueous sodium hydroxide. The remaining gas occupied 95 cm³ at 101 kPa and 30 °C.
 - Write an equation, with state symbols, representing the thermal (i) decomposition of Group 2 nitrates, $M(NO_3)_2$. [1]
 - Assuming that the gas behaves ideally, calculate the amount of remaining (ii) gas formed. [1]
 - (iii) Hence, calculate the value of x and deduce the identity of metal **M**. [3]
- (c) Calcium oxide is the key ingredient for the process of making cement.

When 1.50 g calcium is burned in air, calcium oxide is formed together with a red brown solid.

The red brown solid has the following composition by mass: Ca, 81.1%; N, 18.9%. Adding water to the red brown solid produces calcium hydroxide and 19.2 cm³ of ammonia gas at room temperature and pressure.

- (i) Deduce the formula of the red brown solid. [2]
- Write the chemical equation for the reaction between the red brown solid (ii) with water. [1]
- (iii) Write the chemical equation for the reaction of calcium with O_2 and hence calculate the mass of CaO formed when 1.5 g of calcium is burnt in O_2 . [2]

[1]

[2]

[2]

1 (d) A diagonal relationship is said to exist between certain pairs of diagonally adjacent elements in the second and third Periods of the Periodic Table.

For instance, lithium shows similar chemical properties to magnesium.

- (i) Write the equation for the thermal decomposition of lithium carbonate. [1]
- (ii) Explain why this is unlike that of the other Group 1 carbonates. [1]
- (e) Lithium nitride, Li₃N, is a red solid. It is the only stable Group 1 nitride. The nitrides of all the Group 2 elements are known.

Li₃N is currently investigated as a storage medium for hydrogen gas.

 $Li_3N + 2H_2$ $2LiH + LiNH_2$

State the role of hydrogen.

(f) A diagonal relationship also exists between boron and silicon.

 B_2O_3 is acidic, like SiO₂ but unlike the oxides of the other Group 13 elements.

- Write a balanced chemical equation, with state symbols, that illustrates the acidic nature of SiO₂.
- (ii) Write a balanced chemical equation, with state symbols, that illustrates the acidic nature of B_2O_3 , given that it forms a similar anion as in (i). [1]

[Total: 20]

2 (a) Divalent metal cations like tin and cadmium form insoluble precipitates with NaOH (aq).

The K_{sp} of Sn(OH)₂ is 5.45 x 10⁻²⁷ mol³ dm⁻⁹.

- (i) If the concentration of $Cd(OH)_2$ in a saturated solution is 1.217×10^{-5} mol dm⁻³, determine the solubility product of $Cd(OH)_2$. [1]
- (ii) Calculate the molar solubility of $Sn(OH)_2$ in a solution of pH 8. [2]
- (iii) A certain solution has 0.002 mol dm⁻³ each of Cd²⁺ and Sn²⁺.

Calculate the pH range over which the two cations can be effectively separated. [2]

(b) Propanone, also known as "acetone", is one of the most important solvents in organic chemistry – it can be used to dissolve many things from fats and waxes to airplane glue and nail polish.

It decomposes to **H** and ketene (H₂C=C=O). At 600 °C, the decomposition rate constant is 8.7 x 10^{-3} s⁻¹.

- (i) Suggest the identity of **H**. [1]
- (ii) Determine the half-life of the reaction at 600 °C. [1]
- (iii) How much time is required for 75% of a sample of propanone to decompose at 600 °C? [1]
- (iv) The Arrhenius equation links different parameters (its activation energy, the absolute temperature, and its rate constant at that temperature) of a reaction together.

$$k = Ae^{-\frac{E_a}{RT}}$$

Given that the half-life of the reaction at 500 °C is 8700 s and using your answer to **(ii)**, calculate the activation energy (including its units) of this reaction.

4

[2]

2 Under appropriate conditions, butane can be made to isomerise reversibly to (C) 2-methylpropane at 298K.



2-methylpropane

(g)

	butane	2-methylpropane
S° / J K ⁻¹ mol ⁻¹	310	295
ΔH _f ° / kJ mol⁻¹	-127.2	-135.6

Calculate the ΔG° for the forward isomerisation reaction.

- (i) [2] (ii) Given that $\Delta G^{\circ} = -RT \ln K_{p}$, calculate the equilibrium constant, K_{p} , for the isomerisation reaction. [1] (iii) Write the K_p expression for the isomerisation equilibrium. [1]
- Determine the mole fractions of the two gases at equilibrium. (iv) [3]
- (v) In the industry, 2-methylpropane is used to make compound Y, C₈H₁₈.

Y has the whole range of carbon atoms; it contains primary, secondary, tertiary and quaternary carbon atoms.

Draw the structural formula of Y, assuming that the original carbon skeleton did not rearrange. [1] 2 (d) Life in the universe is widely thought to have originated from methanal, HCHO, through the Formose reaction.

In this reaction, methanal is converted into a vast range of sugars like ribose and from there to RNA, an important hereditary material.



(i) X and Y are functional group isomers.

Y does not rotate the plane of plane-polarised light. Fehling's and Tollens' tests are the only tests which can differentiate between **X** and **Y**.

Draw the structural formula of Y.

- Suggest a reason why the interconversion of C to C' <u>cannot</u> be done in a single step.
- (iii) Draw the structural formula of **D**.

[Total: 21]

[1]

3 This question is on Group 17, the halogens.

Chlorate(V), ClO_3^{-} , reacts with chloride according to the equation:

 $2ClO_3^{-}(aq) + 2Cl^{-}(aq) + 4H^{+}(aq) \rightarrow Cl_2(g) + 2ClO_2(aq) + 2H_2O(l)$

(a) To study the kinetics of chlorate(V)-chloride reaction, an experiment was conducted using a mixture in which the concentrations of the reactants are as follows: 0.000480 mol dm⁻³ of ClO₃⁻, 0.1 mol dm⁻³ of Cl⁻ and 0.4 mol dm⁻³ of H⁺.

At five-minute intervals, small samples of the reaction mixture were withdrawn, quenched and placed into the UV-vis spectrometer to record its absorbance value. The absorbance value corresponds to the concentration of the product ClO_2 .

The results of the above experiment are shown below.

Time/min	0	5	10	15	20	25
Absorbance/A	0.000	0.211	0.348	0.436	0.494	0.531



The graph of absorbance/A against time/min is plotted below.

3 (a) (i) Beer-Lambert's Law states that the absorbance values, A, is directly proportional to the concentration of absorbing species, c, as shown below.

$$A = \epsilon c l$$

where ϵ is the molar extinction coefficient and *l* is the path length, which is usually 1.0 cm.

This equation can be used to calculate the absorbance value when maximum amount of C_1O_2 was formed.

Show that the maximum absorbance value in the above experiment is 0.600, given that ε of ClO₂ is 1250 mol⁻¹ dm³ cm⁻¹.

(ii) From the graph provided on the previous page and the information given in (a)(i), determine the half-life with respect to ClO_3^- and hence the order of reaction with respect to ClO_3^- .

To obtain the full rate law, a further experiment was conducted.

The following graph was obtained.



- (iii) Using the above data, determine the order of the reaction with respect to H⁺.
- (iv) Given that the units of k is mol⁻³ dm⁹ s⁻¹, write the rate equation of the chlorate(V)-chloride reaction.
 [1]
- (v) State a physical property that can be monitored as the reaction progresses.
 [1]

[2]

[2]

3	(b)	Heating solid halides with concentrated H ₂ SO ₄ is one of the ways to obtain
		hydrogen halides.

However, the halides have different reactivities with concentrated H₂SO₄.

- (i) Write an equation to show the reaction of KCl with concentrated H_2SO_4 . [1]
- (ii) However, when KI is treated with concentrated sulfuric acid, HI will be contaminated with other gaseous products. Identify these gaseous products.
- (iii) Suggest another reagent that can be reacted with potassium iodide to obtain pure hydrogen iodide. [1]

The interhalogens are compounds that are made up of two or more different halogens.

ICl and IBr are two such examples.

- (c) Even though ICl is more polar than IBr, IBr has a higher boiling point than ICl.
 Explain this dilemma as clearly as you can.
 [2]
- (d) IC*l* reacts with water in which water is acting as the nucleophile.

The equation for the reaction is as follows.

$$ICl + H_2O \rightarrow HX + HOY$$

	(i)	By comparing relative electronegativities, identify X and Y .	[1]
	(ii)	Hence propose a two-step mechanism for this reaction. Include curly arrows and partial charges.	[2]
(e) IC l_3 exists as a planar dimer I_2Cl_6 in solid state. The two iodine atoms are central atoms.			
	(i)	Draw the structure of the dimer.	[1]
	(ii)	The molten form of ICl_3 is able to conduct electricity because it undergoes auto-ionisation.	
		One species is square planar and the other is bent with respect to the central iodine atom.	
		Deduce the formulae of the cation and the anion.	[2]
		[Total	: 191

Section B - Choose <u>either</u> question 4 or 5. Begin it on a fresh piece of writing paper.

- 4 α-carotene is a red-orange pigment found in plants and fruits. It is the yellow/orange pigment that gives vegetables and fruits their rich colors. The name "carotene" came about when it was first discovered in carrot roots in 1831.
 - (a) α -carotene is metabolised to form α -ionone and β -ionone, which share the same molecular formula, C₁₃H₂₀O.



- (i) Explain why hot acidified KMnO₄ cannot be used in the conversion of α -carotene to β -ionone.
- (ii) Based on the structure of α -carotene, draw the structural formula of α -ionone.
- (iii) Specify what type of constitutional isomerism is shown between α-ionone and β-ionone.
- (iv) Give the structural formula of a constitutional isomer of β -ionone, where neutral iron(III) chloride solution is the <u>only</u> chemical test that can distinguish between itself and β -ionone.
- (v) Give the structural formula of a constitutional isomer of β-ionone, which can be distinguished from β-ionone by warm alkaline iodine.
 [1]

[1]

[1]

[1]

4 (b) β -ionone forms β -damascone via an electrochemical route.



It involved the formation of ketoxime in step I, followed by an anodic oxidation in step II. In the last step, the oxygen-nitrogen bond was cleaved to give β -damascone.

(i)	Name the type of reaction in step I.	[1]
-----	--------------------------------------	-----

(ii) Construct a balanced half-equation for the oxidation of ketoxime in step II under acidic conditions.

You are to use "ketoxime" and "**K**" in your half-equation. [1]

(iii) β -damascone exists as a pair of cis-trans isomers. Draw the cis isomer. [1]

4 (c) Ethylenediaminetetraacetic acid (EDTA) is a colourless, water-soluble solid. It is on the World Health Organization's List of Essential Medicines.

EDTA can be synthesised via the following reaction route.



- (i) X gave a white precipitate when aqueous silver nitrate was added to it. Give the structural formula of X.
- (ii) Give the structural formula of **Y**. Hence, state a potential problem that might happen when it reacts with chloroethanoic acid.
- (d) Chloroethanoic acid can be synthesised from methanal via a three-step route. State the reagents and conditions used in each step and the intermediates involved.
- (e) The Kolbe electrolysis involves conversion of carboxylate salts into alkanes. It can be represented by the following half-equation.

 $3 R_1 COO^- + 3 R_2 COO^- \rightarrow R_1 - R_1 + R_1 - R_2 + R_2 - R_2 + 6CO_2 + 6e^-$

where R_1 and R_2 may not represent the same alkyl group.

In a certain electrolysis, two carboxylate salts were electrolysed, forming three alkanes.

The relative molecular masses are 58.0, 86.0 and 114.0. The alkane with M_r 58.0 is non-chiral. The alkane with M_r 114.0 exists as three stereoisomers - two of which are chiral and the third is non-chiral.

- (i) Identify the two carboxylate salts used. [2]
- (ii) Draw the stereochemical formula of any of the two chiral stereoisomers. [1]

[1]

[2]

[4]

4 (e) In a special application of the Kolbe electrolysis, fumarate was electrolysed into ethyne.



(iii) Kolbe electrolysis proceeds via radical intermediates, which are formed and reacted via homolytic fission and fusion.

Explain the term homolytic fission.

[1]

(iv) The decomposition of fumarate into ethyne and carbon dioxide happens through two steps – the first of which forms the neutral diradical species below as the intermediate.



Use curly arrows to show how the intermediate decomposes to ethyne and carbon dioxide. [1]

[Total: 20]

5 Melanin is a naturally-occurring organic compound that is responsible for the colour of the skin. The higher the melanin content, the darker the skin is. In the Asian market, skin-lightening lotions have grown to be the most on-demand skin care cosmetic products.

One common ingredient of such lotions is kojic acid, which works by slowing down the rate of formation of melanin. It is obtained from the fermentation of rice malt.

The structure of kojic acid is shown below with the two hydroxyl groups labelled **a** and **b**.



kojic acid

(a) For each of the hydroxyl groups **a** and **b**, explain if it will be substituted with a chlorine atom when one mole of kojic acid is reacted with two moles of PCl_5 at room temperature.

[2]

(b) Kojic acid also functions as an antioxidant, scavenging hydroxyl radicals ('OH) that are responsible for ageing. One mole of kojic acid can react with a total of four moles of the radicals to form comenic acid via an intermediate Z as shown in the scheme below.



(i) Given the following information, use curly arrows to draw the mechanism for the formation of comenic acid from compound Z in two separate steps.

Step 1:

An •OH radical reacts with compound Z to form an organic radical intermediate with H₂O being formed as a by-product.

<u>Step 2</u>:

Another •OH radical react with the intermediate formed from the first step. [2]

(ii) Draw another possible organic by-product in the formation of comenic acid from compound **Z**.

5 (c) Besides kojic acid, vitamin E is also an essential ingredient that acts as an antioxidant in skin-lightening lotions.

One class of vitamin E is tocopherol, the general structure of which is shown below.



tocopherol

The table below shows the substituent R of two different types of tocopherol, namely α - and β -tocopherol.

	–R
α-tocopherol	–CH ₃
β-tocopherol	_H

Arrange phenol, α -tocopherol and β -tocopherol in an increasing order of pK_a values. Hence, explain the difference in acidity among the three compounds. [3]

- ility of skin-lightening lations
- (d) To increase the effectiveness and marketability of skin-lightening lotions, titanium dioxide, TiO₂, is often added as a sunscreen agent due to its reflective property.

Solid titanium dioxide reacts with hydrofluoric acid, HF, to form a coordination compound, $H_2[TiF_6]$, and water.

- (i) Write a balanced chemical equation, with state symbols, for the above reaction. [1]
- (ii) Explain whether the above reaction is a redox reaction. [1]
- (iii) State the electronic configuration of titanium in H₂[TiF₆]. Hence, explain whether the complex ion is coloured. [3]

5 (e) Hydroquinone and catechol are isomers.

Hydroquinone has been used for more than half a century in the formulation of skin lotions to help fade uneven skin tone and dark spots. Catechol, on the other hand, is an important synthetic precursor to pesticides, flavours, and fragrances.

The structures of both compounds, together with their boiling points, are shown below.



hydroquinone (boiling point: 287 °C)



catechol (boiling point: 246 °C)

- (i) Explain why catechol has the lower boiling point. [1]
- (ii) Catechol is able to function as a bidentate ligand, but not hydroquinone.

Explain why hydroquinone cannot function as a bidentate ligand. [1]

5 (f) Catechol is used industrially to make ethylvanillin, a flavourant reputedly three times more potent than vanillin itself.



Assume that the ether functional group (R–O–R') is inert.

(i)	Explain the purpose of adding NaOH in step I.	[1]
(ii)	Name reagent S in step I .	[1]
(iii)	Outline the synthetic pathway to convert ethylvanillin back to compound T . Draw the intermediate formed.	[3]
	[Total:	201

END OF PAPER