Name: ...... ( ) Class: Sec 4 SG 1 / 2

# **Queenstown Secondary School**



### Preliminary Examination 2024 Secondary Four Express Chemistry 6092/01

28 August 2024 Time: 1145 – 1245h Wednesday Duration: 1 hour

Setter:

Additional Materials: Multiple Choice Answer Sheet

#### READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write your name, class and index number on the Answer Sheet in the spaces provided.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A**, **B**, **C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

#### Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer. Any rough working should be done in this booklet.

A copy of the Periodic Table is printed on page 21.

The use of an approved scientific calculator is expected, where appropriate.


1 The reaction of manganate(VII) ions with ethanedioate ions in acid solution may be represented by the following equation.

$$2MnO_4^{-1}$$
 (aq) +  $16H^+$  (aq) +  $5C_2O_4^{2-1}$  (aq)  $\rightarrow 2Mn^{2+1}$  (aq) +  $8H_2O_1(l)$  +  $10CO_2_1(g)$ 

A student is exploring various ways to measure the rate of the reaction by measuring changes in different variables of the reaction.

Which of the following methods of monitoring the rate of reaction are suitable?


- 1 volume of gas produced
- 2 pH of the reaction mixture
- 3 mass of the reaction mixture
- 4 amount of precipitate obtained
- 5 intensity of the purple colour of the reaction mixture
- **A** 1, 2 and 3
- **B** 1, 3 and 4
- **C** 1, 2, 3 and 5
- **D** 2, 3, 4 and 5
- **2** The heating curve of an impure sample of substance X is shown below.



What are the melting point and boiling point of pure substance X?

	melting point / °C	boiling point / °C	
Α	45	129	
В	47	128	
С	48	127	
D	49	130	

**3** A student carried out an experiment using the set-up shown below to separate a mixture containing four alcohols.



The table gives the boiling points of the four alcohols used.

alcohol	butanol	ethanol	pentanol	propanol
boiling point / °C	117	79	138	97

Despite repeated attempts, the student only managed to obtain two alcohols from the mixture.

Which alcohols did he fail to obtain?

**A** butanol and ethanol


**C** ethanol and propanol

**B** butanol and pentanol

**D** pentanol and propanol

4 Two experimental set-ups used to demonstrate the diffusion of gases are shown in the diagrams below. In each porous pot is carbon monoxide.

In the first experiment, the gas introduced into the beaker is fluorine gas, while in the second experiment, hydrogen gas was introduced.




What changes, if any, to the water levels X and Y would you expect to see in both experiments?


	experiment 1	experiment 2	
Α	Y is higher than X	X is higher than Y	
В	X is higher than Y	Y is higher than X	
С	X and Y remain the same	Y is higher than X	
D	X and Y remain the same	X and Y remain the same	

**5** An ion, X<sup>2-</sup>, has a mass number of m and it contains n electrons. What does the nucleus of an atom of X contain?

	number of protons	number of neutrons	
Α	n – 2	m – n	
В	n – 2	m – n + 2	
С	n + 2	m – n + 2	
D	n + 2	m – n – 2	

6 Part of the giant lattice structure of magnesium oxide and sodium chloride is shown.





The structure repeats to make a giant lattice.

In the giant lattice, how many negative ions directly surround each positive ion?

- **A** 3
- **B** 4


- **C** 5
- **D** 6

7 The diagram shows the structural formula of the covalent molecule hydrazine, N<sub>2</sub>H<sub>4</sub>.

Which row is correct?

	total number of electrons	total number of electrons
	involved in bonding	<b>not</b> involved in bonding
Α	5	4
В	5	8
С	10	4
D	10	8

8 The structures of three substances P, Q and R are shown below.



Which statements are correct?

- 1 P and R can conduct electricity in the molten state.
- 2 Q is an element while P and R are compounds.
- 3 P has a giant structure while Q has a simple structure.
- **A** 1 and 2
- **B** 1 and 3
- **C** 2 and 3
- **D** 1, 2 and 3
- 9 The information below shows the percentage of carbon in two samples of steel, Q and R. Q consists of 60% carbon while R consists of 20% carbon.

Which statement is correct about the two samples of steel?

- **A** Q has higher strength and less brittle than R.
- $\boldsymbol{\mathsf{B}}\ \ \mathsf{R}$  has higher strength and more brittle than  $\mathsf{Q}.$
- **C** Q has lower strength and more brittle than R.
- **D** R has lower strength and less brittle than Q.
- **10** Upon strong heating, a metal nitrate compound undergoes decomposition according to the following equation:

$$2XNO_{3}\left( s\right) \rightarrow2X\left( s\right) +2NO_{2}\left( g\right) +O_{2}\left( g\right)$$

Complete decomposition of 3.40 g of the nitrate gives 240 cm<sup>3</sup> of oxygen, measured at room temperature and pressure.

What is the relative atomic mass of X?

- **A** 85
- **B** 108
- **C** 133
- **D** 170

11 The structure of oxalic acid is shown.

A 25.0 cm<sup>3</sup> solution of oxalic acid reacts completely with 15.0 cm<sup>3</sup> of 2.50 mol/dm<sup>3</sup> aqueous sodium hydroxide.

What is the concentration of the oxalic acid?

- **A** 0.750 mol/dm<sup>3</sup> **B** 2.08 mol/dm<sup>3</sup> **C** 1.50 mol/dm<sup>3</sup> **D** 4.17 mol/dm<sup>3</sup>

Aspirin,  $C_9H_8O_4$ , is made from salicylic acid,  $C_7H_6O_3$  according to the equation: 12

$$C_7H_6O_3 + C_4H_6O_3 \rightarrow C_9H_8O_4 + CH_3COOH$$

Assuming a 70% yield, what is the mass of salicylic acid required to make an aspirin tablet of 325 mg?

 $[M_r: C_7H_6O_3, 138; C_9H_8O_4, 180]$ 

- **A** 174 mg
- **B** 249 mg **C** 356 mg **D** 424 mg

13 50.0 cm<sup>3</sup> of hydrochloric acid has a pH of 1.0.

This acid requires 25.0 cm<sup>3</sup> of aqueous sodium hydroxide to be neutralised.

A second 50.0 cm<sup>3</sup> solution contains the weak acid, ethanoic acid.

The hydrochloric acid and ethanoic acid have the same concentration.

How will the pH of ethanoic acid and the volume of NaOH needed for neutralisation differ, if at all, from the hydrochloric acid?

	рН	volume of NaOH needed
		for neutralisation
Α	higher than HCl	lower than for HCl
В	higher than HCl	equal to HC <i>l</i>
С	lower than HCl	lower than for HCl
D	lower than HCl	equal to HC <i>l</i>


- **14** What is a characteristic property of all bases?
  - **A** Alkalis are bases which are insoluble in water.
  - **B** They dissolve in water to produce hydroxide ions.
  - **C** They form salts with acids.
  - **D** They react with ammonia to form ammonium salts.
- 15 Which statements about oxides are correct?
  - 1 An aqueous solution of sulfur dioxide has a pH less than 7.
  - 2 An aqueous solution of potassium oxide turns blue litmus paper red.
  - 3 Carbon dioxide reacts with ammonia to make a salt.
  - 4 Carbon monoxide reacts with hydrochloric acid to make a salt.
  - **A** 1 and 2 **B** 1 and 3 **C** 2 and 3 **D** 3 and 4
- 16 Which of the following method could **not** be used to prepare a dry sample of lead salt?

	name of salt	method
Α	lead(II) carbonate	add aqueous sodium carbonate to aqueous lead(II)
^	lead(II) carbonate	nitrate
В	lead(II) chloride	add hydrochloric acid to aqueous lead(II) nitrate
	lood/II) indide	add nitric acid to lead(II) carbonate, then add aqueous
С	lead(II) iodide	potassium iodide
D	lead(II) sulfate	add sulfuric acid to lead(II) carbonate

17 Ammonia is produced from Haber Process using a suitable catalyst.

$$N_2(g) + 3H_2(g) \Longrightarrow 2NH_3(g)$$

The following graph shows the different yields of ammonia at different temperature and pressure.



Which of the following is **not** true?

- **A** A higher percentage yield of ammonia can be obtained at higher pressure.
- **B** A higher percentage yield of ammonia can be obtained at lower temperature.
- **C** Some of the ammonia formed will decompose to form hydrogen and nitrogen.
- **D** At the right conditions of temperature and pressure, all of the hydrogen and nitrogen can be converted into ammonia.

**18** Three tests are carried out on salt X.

S/N	test	result
1	To 2 cm <sup>3</sup> of salt X, add aqueous sodium	White precipitate, soluble in
	hydroxide until no further change. Warm	excess giving a colourless
	the mixture. Keep the mixture for Test 4.	solution.
2	To 2 cm <sup>3</sup> of salt X, add aqueous	White precipitate, soluble in
	ammonia until no further change.	excess giving a colourless
		solution.
3	To 2 cm <sup>3</sup> of salt X, add dilute nitric acid	No visible reaction.
	followed by aqueous silver nitrate.	
4	To 2 cm <sup>3</sup> of the mixture in test 1, add	Gas evolved turns damp litmus
	aluminium foil.	paper blue.

What is the identity of salt X?

A aluminium chloride C zinc chloride

B ammonium nitrate D zinc nitrate

19 A disproportionation reaction occurs when the same element undergoes both oxidation and reduction simultaneously in a chemical reaction.

Which of the following is **not** a disproportionation reaction?


**A** Mg + FeC
$$l_2 \rightarrow$$
 MgC $l_2$  + Fe

$$\label{eq:Bound} \textbf{B} \quad H_2SO_4 + Cu_2O \rightarrow Cu + CuSO_4 + H_2O$$

$$\label{eq:continuous} \textbf{C} \quad 2H_2O_2 \rightarrow O_2 + 2H_2O$$

$$\textbf{D} \quad 2\text{NO}_2 + \text{H}_2\text{O} \rightarrow \text{HNO}_2 + \text{HNO}_3$$


20 The diagrams show an electrolysis set-up using inert electrodes.



Which could be liquid Y?

- 1 aqueous magnesium nitrate
- 2 aqueous copper(II) sulfate
- 3 concentrated hydrochloric acid
- 4 dilute sulfuric acid
- **A** 1 and 4 only **B** 2 and 4 only **C** 1, 2 and 4 only **D** 1, 3 and 4 only

21 A student set up two simple cells as shown below.



He recorded four statements in his notebook.

statement 1: In cell 1, sodium ions gain electrons to form sodium.

statement 2: In cell 2, copper(II) ions gain electrons to form copper.

statement 3: In both cells, magnesium loses electrons to form magnesium ions.

statement 4: The voltage of cell 1 is greater than cell 2.

#### Which statements are incorrect?

- **A** 1 and 2
- **B** 1 and 4
- **C** 1, 2 and 4
- **D** 2, 3 and 4
- 22 Which statement regarding the Periodic Table is correct?
  - **A** The elements are arranged by increasing relative atomic mass.
  - **B** Across a period from left to right, elements have weaker reducing power.
  - **C** Down Group 1, the elements become stronger oxidising agents.
  - **D** Down Group 17, the elements become weaker reducing agents.
- 23 Excess bromine is shaken with a mixture of sodium chloride and sodium iodide solutions. Which substances will the final mixture contain?
  - A bromine, iodine, sodium bromide
  - **B** bromine, iodine, sodium bromide, sodium chloride
  - **C** bromine, iodine, sodium bromide, sodium iodide
  - **D** iodine, sodium bromide, sodium chloride

The properties of the element vanadium, V, can be predicted from its position in the Periodic Table.

Which row identifies the properties of vanadium?

	can be used	conducts	forms coloured	has a law density
	as a catalyst	electricity at r.t.p.	compounds	has a low density
Α	✓	✓	✓	×
В	✓	✓	×	✓
С	✓	×	✓	✓
D	*	✓	✓	✓

[key:  $\checkmark$  = yes;  $\times$  = no]


25 The table below shows three unknown metals and their method of extraction.

metal	method of extraction
Р	reduction of ore by carbon
Q	electrolysis
R	mining from the ground

Which statement regarding the three metals is true?

- **A** Metal Q should be found high up in the reactivity series.
- **B** Metal R is most likely to be found above hydrogen in the reactivity series.
- **C** Metal R is the most reactive metal among all three.
- **D** The oxide of P must be the most stable compared to the oxides of the other two metals.

**26** The diagram shows the energy profile for a chemical reaction.



What is the correct description of the reaction?

	sign of $E_A$	sign of $\Delta H$	overall energy change
Α	-	-	exothermic
В	+	+	endothermic
С	-	+	endothermic
D	+	+	exothermic

27 Methane burns in excess oxygen to produce carbon dioxide and water.

$$CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(g)$$

Given the following information of the bond energies, calculate the enthalpy change of the reaction.

bond	C — H	O = O	C = 0	O – H
bond energy / kJmol <sup>-1</sup>	410	496	805	460

**A** −359 kJ/mol

**B** –818 kJ/mol

**C** +102 kJ/mol

**D** +818 kJ/mol

28 In which reaction is pressure least likely to affect the rate of reaction?


**A**  $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$ 

**B**  $HCl(g) + NH_3(g) \rightarrow NH_4Cl(s)$ 

**C**  $CO_2(g) + Ca(OH)_2(aq) \rightarrow CaCO_3(s) + H_2O(l)$ 

**D**  $K_2CO_3$  (s) +  $H_2SO_4$  (aq)  $\rightarrow K_2SO_4$  (aq) +  $H_2O$  (l) +  $CO_2$  (g)

29 In the first experiment, excess magnesium reacts with 25 cm<sup>3</sup> of 1.0 mol/dm<sup>3</sup> hydrochloric acid to produce hydrogen gas. The volume of hydrogen produced is measured and shown in the graph.



Graphs P and Q show the volume of hydrogen produced under different conditions. What changes in conditions produce graphs P and Q?

- **A** P uses a higher temperature and Q uses a lower temperature.
- **B** P uses 25 cm³ of 1.5 mol/dm³ hydrochloric acid and Q uses 25 cm³ of 0.5 mol/dm³ hydrochloric acid.
- **C** P uses a catalyst and Q uses 25 cm³ of 0.5 mol/dm³ hydrochloric acid.
- **D** P uses smaller strips of magnesium of the same mass and Q uses a lower temperature.
- **30** The table shows the boiling points of four fractions P, Q, R and S, obtained when crude oil is distilled.

fraction	boiling point range / °C
Р	35 – 75
Q	80 – 145
R	150 – 250
S	> 250


How is fraction P different from fraction S?

- A Fraction P is in less demand than S.
- **B** Fraction P is more viscous than fraction S.
- **C** Fraction P is more flammable than fraction S.
- **D** Fraction P contains molecules of larger relative molecular masses than fraction S.

31 Biodiesel, an alternative fuel made from vegetable oil, can be used as a fuel for vehicles. Although carbon dioxide is released during the combustion of biodiesel, scientists still claim that it is a carbon neutral fuel.

Which is the basis for this claim?

- A Biodiesel is not a carbon compound.
- **B** Biodiesel produces less carbon dioxide when it burns.
- **C** Plants release carbon dioxide during respiration.
- **D** Plants absorb carbon dioxide during photosynthesis.
- **32** Pentane, C<sub>5</sub>H<sub>12</sub>, and octane, C<sub>8</sub>H<sub>18</sub>, are alkanes present in the petrol fraction. Which statements about alkanes are correct?
  - 1 They are unsaturated hydrocarbons.
  - 2 Their general formula is  $C_nH_{2n+2}$ .
  - 3 Pentane has a higher boiling point than octane.
  - 4 Both pentane and octane undergo substitution reaction with chlorine in the presence of light.
  - **A** 1 and 2
- **B** 1 and 4
- **C** 2 and 3
- **D** 2 and 4
- 33 Squalene, a naturally occurring polyunsaturated oil present in sharks can be reduced to form a saturated hydrocarbon using titanium as a catalyst.




A 0.100 mol sample of squalene reacted with 14.4 dm<sup>3</sup> of hydrogen at room temperature and pressure to form a saturated hydrocarbon, C<sub>30</sub>H<sub>62</sub>.

What is the molecular formula of squalene?


- **A**  $C_{30}H_{50}$
- **B**  $C_{30}H_{52}$
- **C** C<sub>30</sub>H<sub>54</sub>
- $D C_{30}H_{56}$

**34** The displayed formulae of two compounds are shown.



Which statement(s) about the compounds is/are correct?

- 1 Both compounds are from the same homologous series.
- 2 Both compounds have the same molecular formula.
- 3 Both compounds have the same percentage mass of carbon.
- 4 Both compounds undergo the same type of reactions.
- A 1 and 2 only B 1 and 4 only C 2 and 3 only D 3 and 4 only
- **35** The structures of compounds W, X, Y and Z are shown below.



What reactions do compounds W, X, Y and Z undergo?

	decolourises	decolourises acidified	effervescence with				
		aqueous potassium	aqueous sodium				
	aqueous bromine	manganate(VII)	carbonate				
Α	X and Y	X and Z	W and Y				
В	X and Y	W, X and Y	W and Y				
С	W and Z	X and Z	W, X and Y				
D	W and Z	W, X and Y	W, X and Y				

**36** Two esters have the same molecular formula,  $C_3H_6O_2$ .

	Wha	at are the names	of these two e	esters?				
	1	methyl ethanoa	ite					
	2	ethyl methanoa	ite					
	3	ethyl propanoat	te					
	4	propyl methano	ate					
	Α	1 and 2	<b>B</b> 1 and 3	С	2 and 4	D	3 and 4	
37	·	ine oil is used to viscosity of engir –CH <sub>2</sub> –CH(CH		n of the cha	in of one s	such polym	er is shown	below
		olecule of this ponomer are require	ed to form one	molecule o	f this polyr	•	nolecules of	f
	Α	4	<b>B</b> 5	С	8	D	10	

**38** The structure below shows part of a polymer.

Which one of the following options show the correct monomers?

A	HO NH <sub>2</sub>	OH O=O NH <sup>2</sup>
В	H I N N C OH	HO NH2
С	HO C N	H − N OH C OH
D	H N OH CE O	$H_2N$ $N$ $N$ $NH_2$

- **39** A sample of air is slowly passed through aqueous sodium hydroxide and then over heated copper. Which gases are removed by this process?
  - A carbon dioxide and water vapour
- **C** nitrogen and oxygen
- **B** carbon dioxide and oxygen
- **D** nitrogen and water vapour

**40** Nitrogen monoxide, NO, is formed in the engines of petrol-powered cars. One constituent of petrol is pentane, C<sub>5</sub>H<sub>12</sub>. Nitrogen monoxide is removed from exhaust fumes by catalytic converters.

Which row identifies the reactants that produce nitrogen monoxide and a reaction that removes it in the catalytic converter?

	reactants that produce NO	reaction that removes NO
Α	pentane + one gas found in air	$NO + CO \rightarrow \frac{1}{2}N_2 + CO_2$
В	pentane + one gas found in air	$NO + CO_2 \to NO_2 + CO$
С	two gases found in air	$NO + CO \rightarrow \frac{1}{2}N_2 + CO_2$
D	two gases found in air	$NO + CO_2 \rightarrow NO_2 + CO$

The Periodic Table of Elements

	18	He Helium	4 5	10	Š	neon 20	2 6	Ā	argon 40	36	궇	krypton 84	24	Xe	xenon	131	98	쮼	radon	1	118	ő	oganesson	E)	
	17			თ	L.	fluorine 10	17	C C	chlorine 35.5	35	ā	bromine 80	53	Н	iodine	127	85	¥	astatine	1	117	<u>s</u>	tennessine	Ĺ	
	16			œ	0	oxygen	2 4	်	sulfur 32	34	Se	selenium 79	52	Te	tellurium	128	84	Ъ	polonium	ı	116	_	livermorium	t	
	15		,	/	z	nitrogen 1.4	15	<u>.</u>	phosphorus 31	33	As	arsenic 75	51	Sp	antimony	122	83	ā	bismuth	209	115	Mc	moscovium	t	
	14			9	ပ	carbon 12	1 4	Si	silicon 28	32	Ge	germanium 73	20	S	£	119	82	Ъ	lead	207	114	Fl	flerovium	Ü	
	13			2	മ	boron 11	- (2)	Al	aluminium 27	31	Ga	galfium 70	49	디	indium	115	81	11	thallium	204	113	R	nihonium	Ę	
									12	30	Zu	zinc 65	48	ပ္ပ	cadmium	112	80	Ε̈́	mercury	201	112	ວົ	copernicium	ı	
									F	29	C	copper 64	47	Ag	silver	108	79	Au	plog	197	111	Rg	roentgenium	Ē.	
dno									10	28	Z	nickel 59	46	Pd	pelladium	106	78	₹	patinum	195	110	Ds	darmstadtium	t)	
Group									6	27	ပိ	cobalt 59	45	몺	modium	103	77	ī	iridium	192	109	¥	meitnerium	Ē	
		Hydrogen							œ	56	Fe	iron 56	44	R	ruthenium	101	92	Os	osmium	190	108	£	hassium	ŀ	
									7	25	M	manganese 55	43	ည	technetium	1	75	Re	rhenium	186	107	В	pohrium	I	
				mber	mper	lod Do	2000	2		9	24	ပ်	chromium 52	42	Mo	molybdenum	96	74	≥	tungsten	184	106	Sg	seaborgium	Ē
			į	ney	proton (atomic) number	atomic symbol	name relative atomic mass			2	23	>	vanadium 51												E
				proton	ator	relativ			4	22	ı	titanium 48	40	Zr	zirconium	91	72	Ξ	hafnium	178	104	抷	rutherfordium	Ü	
									က	21	Sc	scandium 45	39	>	yttrinm	89	57-71	lanthanoids			89-103	actinoids			
	2			4	Be	beryllium	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium	88	56	Ba	barium	137	88	Ra	radium	t)	
	-			က	:=	lithium 7	- 11	Na	sodium 23	19	¥	potassium 39	37	R <sub>b</sub>	rubidium	82	55	S	caesium	133	87	ŭ	francium	ſ	

71 Lu	lutetium 175	103	۲	lawrencium	1		
6 <del>Q</del>	ytterbium 173	102	8	nobelium	1		
69 Tm	thulium 169	101	Md	mendelevium	1		
8 <u>P</u>	167	100	F	fermium	1		
67 Ho	holmium 165	66	Es	einsteinium	1		
96 Dv	dysprosium 163	98	ರ	californium	1		
65 Tb	terbium 159	26	ă	berkelium	ı		
64 Gd	gadolinium 157	96	S	curium	1		
63 Eu	europium 152	92	Am	americium	ı		
62 Sm	samarium 150	94	Pu	plutonium	1		
61 Pm	promethium	93	ď	neptunium	ī		
9 N	neodymium 144	92	⊃	uranium	238		
59 Pr	praseodymium 141	91	Ъа	protactinium	231		
Çe 28	140	06	드	thorium	232		
57 La	lanthanum 139	88	Ac	actinium	1		
lanthanoide	200	actinoids					

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.). The Avogadro constant,  $L = 6.02 \times 10^{23} \, \text{mol}^{-1}$ .

## MARK SCHEME

1	2	3	4	5	6	7	8	9	10
С	С	В	В	В	D	D	В	D	В
11	12	13	14	15	16	17	18	19	20
Α	С	В	С	В	D	D	D	Α	Α
21	22	23	24	25	26	27	28	29	30
С	В	В	Α	Α	В	В	D	С	С
31	32	33	34	35	36	37	38	39	40
D	D	Α	С	С	Α	С	Α	В	С

<u>Tally</u>

A - 8

B – 12

C – 11

D – 9