

1 How many hydrogen atoms are present in 4.0 g of methane? (L = Avogadro constant)

- A $\frac{L}{16}$ B $\frac{L}{4}$ C L D $4L$

2 The table shows the relative abundance of a sample of naturally occurring isotopes of zinc.

| isotope | relative abundance |
|-------------------------|--------------------|
| ${}^{64}_{30}\text{Zn}$ | 10 |
| ${}^{65}_{30}\text{Zn}$ | 8 |
| ${}^{67}_{30}\text{Zn}$ | 2 |
| ${}^{68}_{30}\text{Zn}$ | 1 |

What is the relative atomic mass of this sample of zinc?

- A 64.0 B 64.9 C 68.1 D 72.2

3 Analysis of a mixture of two sulfur-containing gases show that H_2S and CS_2 are present in a 3 : 1 mole ratio.

This mixture is burned in excess oxygen to give CO_2 and SO_2 gas.

What is the mole ratio of the gases CO_2 : SO_2 obtained after complete combustion?

- A 1 : 2 B 1 : 3 C 1 : 4 D 1 : 5

4 2 moles of an oxidising agent, XO_4^- , in the presence of excess acid, oxidised 96.0 dm³ of nitrogen dioxide gas to NO_3^- at room temperature and pressure.

What is the number of moles of electrons accepted by one mole of XO_4^- ?

- A 1 B 2 C 3 D 4


- 5 The first six ionisation energies of an element, **Y**, in kJ mol^{-1} are shown.

| | 1 st | 2 nd | 3 rd | 4 th | 5 th | 6 th |
|--|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Ionisation Energy / kJ mol^{-1} | 738 | 1451 | 7733 | 10543 | 13630 | 18020 |

Y forms an oxide by heating **Y** with oxygen gas.

What is the molecular formula of the oxide of **Y** formed?

- A** YO **B** YO₂ **C** Y₂O **D** Y₂O₃
- 6 The radioactive isotope ${}^{223}_{88}\text{Ra}$ decays to give **Q** and emits a high energy α -particle, ${}^4_2\text{He}$. No other particle is produced.



How many neutrons are present in **Q**?

- A** 86 **B** 133 **C** 135 **D** 219
- 7 Why is the second ionisation energy of fluorine lower than that of oxygen?
- A** There are more paired electrons in the 2p orbitals of fluorine than in oxygen.
- B** The ionic radius of O⁺ is greater than F⁺.
- C** Fluorine has a lower nuclear charge compared to oxygen.
- D** All 2p orbitals of O⁺ are singly filled but one of the 2p orbitals of F⁺ is doubly filled.
- 8 Chlorine atoms in the PCl₅ molecule can be successively replaced by fluorine atoms, with the axial chlorine atoms replaced before the equatorial ones.


Which of the possible molecules formed in the above reaction does **not** have a net dipole moment?

- A** PClF₄ **B** PCl₂F₃ **C** PCl₃F₂ **D** PCl₄F

- 9 **Q** has the following physical properties.
- It is non-volatile.
 - It does not conduct electricity in its standard state.
 - It dissolves in water.

What is the identity of **Q**?

- A** Magnesium
- B** Carbon dioxide
- C** Silicon dioxide
- D** Sodium chloride
- 10 Which statement best explains why the boiling point of butanone (80 °C) is higher than that of pentane (36 °C)?
- A** The covalent bonds in the butanone molecule are stronger than those in the pentane molecule.
- B** The relative molecular mass of butanone is higher than that of pentane.
- C** There are permanent dipole-permanent dipole forces between butanone molecules, but not between pentane molecules.
- D** There are hydrogen bonds between butanone molecules, but not between pentane molecules.
- 11 Which option is correct for the following organic molecule?



| | No. of σ bond | No. of π bond |
|----------|----------------------|-------------------|
| A | 9 | 2 |
| B | 18 | 2 |
| C | 19 | 4 |
| D | 21 | 2 |

- 12 *Use of the Data Booklet is relevant to this question.*

Hexamine has an enthalpy change of combustion of $-4288 \text{ kJ mol}^{-1}$.

12.4 g of hexamine tablets were burnt to heat up 850 cm^3 of water. Given that the process was 75 % efficient and temperature of the water increased from $10 \text{ }^\circ\text{C}$ to $90 \text{ }^\circ\text{C}$, what is the molar mass of hexamine?


- A 105.2 g mol^{-1} B 140.3 g mol^{-1} C 187.1 g mol^{-1} D 249.4 g mol^{-1}
- 13 *Use of the Data Booklet is relevant to this question.*

A reaction which causes the presence of oxides of nitrogen in car exhausts is the formation of NO.



What is the bond energy in NO, in kJ mol^{-1} ?


- A 630 B 810 C 1260 D 1440
- 14 The reaction of a compound **XY** is shown below.



The rate equation for the reaction is $\text{rate} = k[\text{XY}]$ and the half-life is 193s. If the initial concentration of **XY** is $2.0 \times 10^{-2} \text{ mol dm}^{-3}$, what will be the concentration of **XY** after 770 seconds?

- A $1.0 \times 10^{-2} \text{ mol dm}^{-3}$
B $5.0 \times 10^{-3} \text{ mol dm}^{-3}$
C $2.5 \times 10^{-3} \text{ mol dm}^{-3}$
D $1.25 \times 10^{-3} \text{ mol dm}^{-3}$

- 15 The graph represents the decomposition of a sample of hydrogen peroxide in the presence of manganese(IV) oxide.




Which conclusions can be drawn from the graph?

- 1 The rate of decomposition of hydrogen peroxide depends on its concentration in the sample.
- 2 The half-life of the hydrogen peroxide in the sample is 200s.
- 3 The reaction is first order with respect to hydrogen peroxide.

A 1, 2 and 3 **B** 2 and 3 only **C** 1 and 2 only **D** 1 and 3 only

- 16 The diagram shows the structure of a catalytic converter fitted in the exhaust system of a car where harmful gases are converted into carbon dioxide, nitrogen and water vapour.



Which reactions would occur on the surface of the catalyst in the catalytic converter?


- 1 hydrocarbons + oxides of nitrogen \longrightarrow carbon dioxide + water + nitrogen
- 2 carbon monoxide + oxides of nitrogen \longrightarrow carbon dioxide + nitrogen
- 3 carbon monoxide + hydrocarbon \longrightarrow carbon dioxide + water

A 1, 2 and 3 **B** 2 and 3 only **C** 1 and 2 only **D** 1 and 3 only

17 Which pairs of molecules are constitutional isomers?

1 $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ and $(\text{CH}_3)_4\text{C}$

2 butanone and 2-methylpropanal



A 1, 2 and 3 **B** 2 and 3 only **C** 1 and 2 only **D** 1 and 3 only


18 How many isomeric alkenes with formula C_5H_8 are present in the mixture produced when 1,4-dibromopentane is heated with NaOH in ethanol?

A 1 **B** 2 **C** 3 **D** 4

19 Which compound is a product of the hydrolysis of $\text{CH}_3\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$ by boiling aqueous sodium hydroxide?

- A** CH_3OH
B CH_3COOH
C $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
D $\text{CH}_3\text{CH}_2\text{CH}_2\text{COO}^-\text{Na}^+$

- 20 'Slug-bait' is used for killing slugs and contains the compound shown.



This compound is prepared by a sequence of addition reactions of a simple carbonyl compound, **X**, using concentrated sulfuric acid as a catalyst at 0 °C.

What is **X**?

- A ethanal
- B methanal
- C propanal
- D propanone