YISHUN INNOVA JUNIOR COLLEGE JC 2 PRELIMINARY EXAMINATION

CANDIDATE NAME

CG

INDEX NO

CHEMISTRY

Higher 2

Paper 2 Structured Questions

Candidates answer on the Question Paper Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand
in.PaWrite in dark blue or black pen on both sides of the paper.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.12Answer all questions in the spaces provided on the Question3

Answer **all** questions in the spaces provided on the Question Paper.

The use of an approved scientific calculator is expected, where appropriate.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
Pa	aper 1
	/30
Pa	aper 2
1	/7
2	/5
3	/7
4	/17
5	/11
6	/22
7	/6
Penalty	
	/75
Pa	aper 3
	/80
Pa	aper 4
	/55
Overall Percentage (%)	

30 August 2021

2 hours

9729/02

Answer **all** the questions in this section in the spaces provided.

1 Use of the Data Booklet is relevant to this question.

This question is about period 3 elements.

(a) (i) State the electronic configuration of argon.

......[1]

(ii) Using the Cartesian axes shown in Fig. 1.1, draw a fully labelled diagram of the valence orbitals of argon.





[1]

(iii) Two glass vessels **B** and **C** are connected by a closed valve as shown in Fig 1.2.



B contains argon at 20 °C at a pressure of 1×10^5 Pa. **C** has been evacuated and has three times the volume of **B**. In an experiment, the valve was opened and the temperature of the whole apparatus was raised to 100 °C. Calculate the final pressure in the system.

(b) Fig 1.3 shows the fifth, sixth, seventh, eighth and ninth ionisation energies of another element **T** in Period 3.



(i) State and explain which period 3 element has these ionisation energy values.



2 Diamine can ionise in stages.

$$H_2N(CH_2)_nNH_2 \xrightarrow{+H^+} H_2N(CH_2)_nNH_3^+ \xrightarrow{+H^+} H_3N^+(CH_2)_nNH_3^+$$

_ . .

(a) Table 2.1 compares the p K_b values of ethylamine and 1,2-ethanediamine at 25 °C.

Table 2.1						
Base	Formula	р <i>К</i> 1	p <i>K</i> ₂			
ethylamine	CH ₃ CH ₂ NH ₂	3.19	-			
1,2-ethanediamine	$H_2NCH_2CH_2NH_2$	4.11	7.39			

(i) Suggest a reason why the pK_1 value of ethylamine is less than the pK_1 value of 1,2-ethanediamine.

......[1]

(ii) Suggest why the pK_2 value of 1,2–ethanediamine is higher than its pK_1 value.

.....

-[1]
- (iii) 0.10 mol dm⁻³ of HCl (aq) was added to ethylamine solution at 25 °C.

 $CH_3CH_2NH_2(aq) + H_2O(l) \rightleftharpoons CH_3CH_2NH_3^+(aq) + OH^-(aq)$

Deduce without calculation, what happens to the position of equilibrium and the value of pK_1 .

 (b) Adding equimolar of 1,2–ethanediamine and phosgene, $COCl_2$ produced a cyclic compound V with molecular formula $C_3H_6ON_2$. Suggest the structure of compound V.

[1]

[Total: 5]

3 Lysine is an essential amino acid that cannot be synthesised by the human body and must be obtained from the diet. It is found in legumes such as peas, and animal products such as beef and fish.

The structure of lysine is given below.



lysine

(a) A solution was prepared by reacting 1 mole of lysine with 2 moles of hydrochloric acid. This solution was titrated with aqueous sodium hydroxide to obtain the titration curve below.



(i) Draw the structure of the species present at points W, X, Y and Z on the titration curve.



(ii) Explain why the melting point of lysine is high, in terms of structure and bonding.

(b) Leucine is another essential amino acid that is mainly found in legumes.



leucine Write an equation to explain how the zwitterionic form of leucine behaves as a buffer when a small amount of base is added.

.....[1]

[Total: 7]

- 4 Benzene is a natural constituent of crude oil and is mainly used as an intermediate to produce other chemicals.
 - (a) In the presence of anhydrous AlCl₃, benzene can undergo Friedel-Crafts alkylation with CH₃Cl to form methylbenzene. The alkyl side chain can further react to form compound **A**, C₁₄H₁₄, along with side products like hydrogen chloride gas.
 - (i) Suggest the structure of compound A.

(ii) Unlike benzene, phenylamine does **not** undergo Friedel-Crafts alkylation with halogenoalkanes in the presence of $AlCl_3$. This is because the amine reacts with $AlCl_3$ to form a neutral compound **B**. Draw the **displayed** formula of **B**.

[1]

(b) Friedel-Crafts alkylation can also be achieved using alkenes, such as the example below.



(i) In the first step, $(CH_3)_3C^+$ is produced in the presence of $AlCl_3$ and HCl.

Write an equation to show the formation of $(CH_3)_3C^+$.

(ii) Name and outline the mechanism of the reaction to form compound C.

Show all charges and relevant lone pairs and show the movement of electron pairs by using curly arrows.

(iii) Compound **D** is a constitutional isomer of compound **C** and it can exhibit stereoisomerism. Draw diagrams to illustrate the type of stereoisomerism present in **D**.

- [2]
- (c) Chloromethane, CH₃Cl, can be hydrolysed by aqueous sodium hydroxide. The rate equation was experimentally determined to be rate = k [CH₃Cl] [OH⁻]
 - (i) Outline the mechanism of this hydrolysis reaction that is consistent with the rate equation.

(ii)	Explain why the mechanism in (c)(i) is consistent with the rate equation.	
		[1]

[3]

(d) When aqueous sodium hydroxide is added dropwise to aqueous copper(II) nitrate, a pale blue precipitate, Cu(OH)₂, is formed.

The numerical value of the solubility product, K_{sp} , of Cu(OH)₂ is 2.2 × 10⁻²⁰ at 25 °C.

(i) Calculate the solubility of $Cu(OH)_2$ in g dm⁻³.

(ii) If equal volumes of a sample of copper(II) nitrate solution and 0.250 mol dm⁻³ aqueous sodium hydroxide are mixed, calculate the minimum concentration, in mol dm⁻³, of copper(II) ions that must be present in the sample to cause precipitation of Cu(OH)₂.

[1]

(e) A student suggested two methods to prepare 4-methoxymethylbenzene.



BLANK PAGE

5 (a) Table 5.1 below gives the pK_b values of some weak bases.



Table 5.1

(i) Arrange the bases in ascending order of base strength, explaining your reasoning.

(ii) Calculate the pH of 0.200 mol dm^{-3} phenylamine.

[2]

(b) Phenylamine can be synthesised from benzene in a three-step process as shown below.



(i) Draw the structure of intermediate E in the box above.

[1]

- (ii) Identify the type of reaction in step II, and name the reagents and conditions used.
 type of reaction:
 reagents and conditions:
- (iii) Explain why ammonia can be used in place of NaOH in step III.
-[1]
- (c) Phenylamine can undergo the following reactions to form compound F.



(i) State the reagent used in step 1.

.....[1]

(ii) Besides F, another possible product G is also formed. Suggest the structural formula of G.

[1]

[2]

[Total: 11]

6 Wearing of masks can be used for either protection of healthy persons or to prevent onward transmission of Coronavirus disease, COVID-19.

Tea polyphenols extracted from green tea possess antiviral properties. A dip coating of nonwoven fabric mask into tea polyphenol extract can inactivate >99% of tested viruses.

The chemical synthesis of a new type of polyphenol was achieved by reaction of 4–((2–phenylhydrazono)methyl)phenol (4–PHMP) to form Poly(4–PHMP) using sodium hypochlorite, NaOC*l* as oxidising agent in an aqueous alkaline medium.



Fig. 6.1

Data about these 4 compounds are given in Table 6.1.

Table 6.1					
	Molecular Mass /g mol ⁻¹	Solubility in hexane / g cm ⁻³			
4-hydroxybenzaldehyde	122.0				
phenylhydrazine	108.0				
4-PHMP	212.0	0.99			
Poly(4–PHMP)	22065.0	0.00			

(a) The monomer 4–PHMP was prepared by the reaction between 4–hydroxybenzaldehyde and phenylhydrazine in ethanol as shown in Fig.6.1.

15

1.221 g of 4–hydroxybenzaldehyde in 3 cm³ of ethanol and 1.282 g of phenylhydrazine in 2 cm³ of ethanol and were mixed and stirred for 30 mins at room temperature. The precipitated product was purified by recrystallisation to provide 2.06 g of product 4–PHMP.

(i) State the type of reaction occurring.
 (ii) Use the data above to determine, by calculation, whether 4–hydroxybenzaldehyde or phenylhydrazine is the limiting reagent.

(iii) Using answer in (a)(ii), calculate the percentage yield of 4-PHMP.

[2]

(iii) Identify one possible by-product of this reaction in aqueous medium.

- (b) The reaction of 4-PHMP to form Poly(4-PHMP) using NaOC*l* was carried out as follows
 - 1. 0.53 g of 4–PHMP was dissolved in 1 cm³ of aqueous KOH and placed into a 25 cm³ three-necked round-bottomed flask that was fitted with a condenser, a thermometer and a stirrer, in addition to a funnel containing NaOC*I*.
 - 2. After heating to 50°C, NaOC/ was added dropwise over 30 min with stirring and the mixture was heated under reflux for 5 hours.
 - 3. The mixture was allowed to cool to room temperature and 0.5 cm³ of concentrated HC*l* was added.
 - 4. The mixture was filtered and washed with 25 cm³ hot water three times, and the removal of mineral salts was confirmed using an AgNO₃ solution.
 - 5. The unreacted 4-PHMP was separated from the reaction products by washing with hexane and dried in an oven at 105 °C.
 - (i) Suggest why 1 cm³ of aqueous KOH is added to dissolve 4–PHMP in step 1.

(ii) State the purpose of adding concentrated HC*l* in step 3 and suggest why the mixture was cooled to room temperature before addition.

.....

-[2]
- (iii) Suggest the identity of the mineral salt which is removed when the mixture was washed with 25 cm³ hot water three times after the reaction is completed in step 4.

......[1]

(iv) In step 5, the unreacted 4–PHMP was separated from the reaction products by washing with hexane. Using data from Table 6.1 and knowledge of structure and bonding, suggest how the product, Poly(4–PHMP) is separated from unreacted 4–PHMP.

[3]

(c) A researcher carried out the reaction of 4-PHMP described in (b) by repeating the reaction with different parameters such as reaction temperature and reaction time, on the yield of poly 4-PHMP. He also decided to use O₂ instead of NaOC*l* as the oxidising agent. The results are summarised in Fig. 6.2a (reaction temperature) and Fig. 6.2b (reaction time).



- Source: Demir, H. A novel polyphenol: synthesis, characterization and investigation of its thermal and electrochemical properties. Polym J44, 699–705 (2012). https://doi.org/10.1038/pj.2012.17
- (i) From Fig. 6.2a and 6.2b, the optimal reaction condition for producing Poly(4-PHMP) using O₂ is 60 °C for 40 h.

State the optimal reaction condition in terms of temperature and time, for producing Poly(4-PHMP) using NaOC*l*.

(ii) Hence suggest with reasoning which oxidising agent should be more suitable for use in commercial large scale production of Poly(4-PHMP).





Describe the reagents and conditions needed for steps 1 and 2 and draw the structure of the intermediate formed in the box above.



(e) Patients with COVID-19 may use a pulse oximeter to measure the oxygen level in their body. However, the oxygen level measured by a pulse oximeter is not the only way to know how sick someone is.

Pulse oximeters measure how much of the haemoglobin in blood is carrying oxygen (oxygen saturation). Oxygen saturation refers to the percentage of the available haemoglobin that carries oxygen.

(i) One molecule of haemoglobin molecule can bind up to four molecules of oxygen according to the following equation.

 $Hb(aq) + 4O_2(aq) \rightleftharpoons Hb(O_2)_4(aq) \qquad \qquad \mathcal{K}_c = 3.00 \times 10^{20}$

Write an expression for K_c for the reaction, and calculate the ratio of $[Hb(O_2)_4]$ to [Hb], given that the $[O_2]$ is 9.6 x 10⁻⁶ mol dm⁻³.

(ii) Calculation of oxygen saturation (SaO₂) is given by the following formula:

$$SaO_2(\%) = \frac{[HbO_2]}{[Hb]+[HbO_2]} \times 100\%$$

where

[HbO₂] is the concentration of haemoglobin [Hb] is the concentration of deoxyhaemoglobin

[Hb] + [HbO₂] is the total concentration of haemoglobin capable of binding oxygen

Given that the $[Hb(O_2)_4] = [HbO_2]$, using answers in **(d)(i)**, calculate the oxygen saturation in blood for a man who has the same $[O_2]$ as in **(d)(i)**.

- [1]
- (iii) A normal oxygen level measured by a pulse oximeter is around 97%. Doctors start to worry when this level drops under 90% because this can affect the amount of oxygen going to the brain and other vital organs. People can experience confusion and lethargy at low levels. Levels below 80% are considered dangerous and increase the risk of organ damage.

Based on you answer in (e)(ii), suggest whether the man needs to seek doctor's advice based on his oxygen saturation in blood.

.....[1]

[Total: 22]

- 7 (a) Phosphorus triiodide, PI_3 , is a red solid that is widely used in organic synthesis to convert alcohols into iodoalkanes
 - (i) Draw a 'dot-and-cross' diagram of PI_3 and state its shape.

		Shape:	[2]
	(ii)	Unlike phosphorus pentachloride, PCl_5 , phosphorus pentaiodide, PI_5 , is highly unsta and cannot be prepared at room temperature. Suggest a reason for this.	ble
			. [1]
(b)	The	e boiling points of PC l_5 and PI ₃ are 167 °C and 200 °C respectively.	
	Exp	plain, in terms of structure and bonding, the difference in boiling point between PC l_5 and F	ין₃.
			. [2]
(c)	Wri	te an equation to describe the reaction of P_4O_{10} with water.	
			. [1]
		[Total	: 6]