

DARRELL ER (COPYRIGHTED) ©

TOPIC 7.3: AMMONIA

THE ABOUT

CHAPTER ANALYSIS



TIME

- Relatively straight forward chapter
- 1 **key** concept



EXAM

- Usually tested in MCQs
- Tested together with chapters like Rate of Reaction & Energy Changes



WEIGHTAGE

- Light overall weightage
- Constitute to **1.5%** of marks for past 5 year papers

AMMONIA

RAW MATERIALS (H_2 & N_2)

HABER PROCESS



AMMONIA

Raw materials

Nitrogen and hydrogen are the raw materials that are used in the manufacturing of ammonia, via the Haber process.


Nitrogen is obtained through the process of **fractional distillation of liquid air**.

Hydrogen is obtained through the **cracking of crude oil**.

Iron would act as a **catalyst** to increase the rate of reaction.

AMMONIA

Ammonia (NH₃) is a **weak alkali** when it is in its aqueous state, as it partially dissociates in water to produce low concentration of OH⁻ ions.




\rightleftharpoons **reversible reactions** will never be fully completed.

Displacement of ammonia from its salts

An alkali has the ability to displace the ammonia from an ammonium salt.


For example, potassium hydroxide **displaces ammonia** from ammonium carbonate when the solution is gently heated:



**Chemical reaction for alkali (acid & bases)!*

HABER PROCESS

Haber process



In the process, nitrogen and hydrogen gases are mixed together in the ratio of 1:3.

Conditions

The Haber process is usually carried out at a **temperature of 450°C**, at a **pressure of 200 atm** and with finely divided iron catalyst.

HABER PROCESS

Analysis:


As seen from the graph, the **yield of ammonia increases when pressures are higher and temperatures are lower**.

Hence, to maximise the yield of ammonia, *theoretically*, the pressure levels should be increased and the temperature should be decreased.


However in reality, optimal conditions are kept at 450°C and pressure of 200 atm.

This is because:


- At pressures higher than 200 atm, the machines would be **more costly** and outweigh the benefits of that incremental yield. Also, there will be greater **safety risks at higher pressures**.
- **At temperatures lower than 450°C**, the **rates of reaction** would be **slowed down too much**. It would be more **cost efficient** to use a higher temperature to **increase the rate** despite lowering the percentage yield.
- Due to the **recycling of reactants**, **98% of the reactants** are eventually **converted into ammonia**.



HABER PROCESS



Haber process



In the process, nitrogen and hydrogen gases are mixed together in the ratio of 1:3.

Conditions

The Haber process is usually carried out at a **temperature of 450°C**, at a **pressure of 200 atm** and with **finely divided iron catalyst**.

HABER PROCESS

- **Nitrogen and hydrogen gases** are mixed in a ratio of **1:3**.
- The mixture would be passed through a compressor, where a **pressure of 200 atm** is applied to the gas mixture and then passed through the converter **containing iron catalyst at 450°C to increase the rate of reaction**.
- The ammonia gas formed would be directed into a cooler, **condensing it into a liquid**, while **unreacted nitrogen and hydrogen gases are recycled**.
- The Haber process is **efficient and relatively cheap**, as the starting materials required (nitrogen, hydrogen and iron) are readily available at a low cost.
- **Heat is produced during the reaction (exothermic)**. It **maintains the temperature of the catalyst chamber**.

ADVANCED

things to note

Understanding Haber Process

Rate of reaction is more important than yield

Temperatures lower than 450°C will result in very **slow rates of reaction**. It is more **cost efficient** to use a higher temperature that **increases the rate of reaction**.

Only 15% of the reactants are converted into ammonia. But that is okay because **98% of the reactants** are eventually **reacted to form ammonia**.

Recall how 'pressure' increases rate of reaction

At a higher pressure, the reactants are brought closer together. There are **more reactants per unit volume**.

As a result, there are more collisions between reactants and thus a higher frequency of effective collisions. This causes the rate of reaction to increase.

Recall how 'temperature' increases rate of reaction

A higher temperature of a system means that:

- 1) Reactants have **higher kinetic energy and move faster**
- 2) The **fraction of reactant particles** in the system that have energy **more than or equal to the activation energy** is higher

These two factors increase the **frequency of effective collisions** and essentially result in an increase in the rate of reaction.

Try it yourself! (TYS Question)

4. Which statement about ammonia is correct?

(N2017/P1/Q20)

- A It decomposes on heating to a high temperature to form nitrogen and hydrogen.
- B It dissolves in water to form an acidic solution.
- C It is formed when ammonium salts are heated with sulfuric acid.
- D It reacts with alkalis to form salts.

()

Answer:

4. A

The reaction for the manufacture of ammonia $[\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})]$ is a reversible reaction. It means the backward reaction will take place under certain conditions. Ammonia is an alkali as it dissolves in water to form an alkaline solution. It is formed by heating ammonium salts with an alkali. It reacts with acids to form salts.

Try it yourself! (TYS Question)

5. Ammonia can be displaced from its salts.
Which equation is correct?

(N2020/P1/Q22)

- A $\text{Ca(OH)}_2(\text{s}) + \text{NH}_4\text{Cl}(\text{s}) \xrightarrow{\text{heat}} \text{CaCl}(\text{s}) + \text{NH}_3(\text{g}) + \text{H}_2\text{O}(\text{g})$
- B $\text{Ca(OH)}_2(\text{s}) + (\text{NH}_4)_2\text{SO}_4(\text{s}) \xrightarrow{\text{heat}} \text{CaSO}_4(\text{s}) + 2\text{NH}_3(\text{g}) + 2\text{H}_2\text{O}(\text{g})$
- C $\text{NaOH}(\text{s}) + 2\text{NH}_4\text{NO}_3(\text{s}) \xrightarrow{\text{heat}} \text{NaNO}_3(\text{s}) + 2\text{NH}_3(\text{g}) + \text{H}_2\text{O}(\text{g})$
- D $\text{NaOH}(\text{s}) + (\text{NH}_4)_2\text{SO}_4(\text{s}) \xrightarrow{\text{heat}} \text{NaSO}_4(\text{s}) + 2\text{NH}_3(\text{g}) + 2\text{H}_2\text{O}(\text{g})$ ()

Answer:

5. **B**

Ammonia is formed when an ammonium salt reacts with a base. The chemical formula of the salt is not correct in options A and D while the equation in option C is not balanced.

About Us

OVERMUGGED is a learning platform created by tutors, for students.

Our team of specialist tutors offer 1-to-1 private tuition, group tuitions and crash courses.

Follow us on [IG](#) and join our [Telegram channel](#) to get the latest updates on our free online revision sessions, webinars and giveaways!

If you would want to join Darrell's group tuition, contact him at:

Whatsapp: [8777 0921](#)

Telegram: [@DarrellEr](#)

Website: <https://www.overmugged.com/darrell>

For more free notes & learning materials, visit: www.overmugged.com

Notes prepared by:



Darrell Er
'O' Levels Chemistry & Physics



OVERMUGGED's Curated Notes

Found the free notes useful? We got something better!

OVERMUGGED's curated notes is a **highly condensed booklet** that **covers all content within the MOE syllabus**.

This booklet consist of **key concept breakdowns**, **worked examples** and **exam tips/ techniques** to required to ace your exams.

Get an **upgraded version** of the free notes and supercharge your revision!

Purchase [here](https://www.overmugged.com).



Crash courses

Check out our upcoming crash courses at:
<https://www.overmugged.com/crashcourses>

'O' levels subject available:

- Pure Chemistry
- Pure Physics
- Pure Biology
- Combined Science
- E-Math
- A-Math
- English
- History
- Geography
- Combined Humanities
- Principles of Accounts (POA)