

NANYANG JUNIOR COLLEGE
JC2 PRACTICAL PRELIMINARY EXAMINATION
Higher 2

CANDIDATE
NAME

--

CLASS

--	--	--	--

CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

CHEMISTRY

Paper 4 Practical

9729/04

14 August 2024

2 hour 30 minutes

Candidates answer on the Question Paper

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number, name and class on all the work you hand in.
Give details of the practical shift and laboratory, where appropriate, in the boxes provided.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs
Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions in the spaces provided on the Question Paper.
The use of an approved scientific calculator is expected, where appropriate.
You may lose marks if you do not show your working or if you do not use appropriate units.
Qualitative Analysis Notes are printed on pages 23 and 24.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [] at the end of each question or part question.

Shift
Laboratory

For Examiner's Use	
1	/ 16
2	/ 16
3	/12
4	/ 11
Total	/55

Answer **all** questions in the space provided.

1 An investigation of a gemstone

Spinel minerals form crystals that are often used as gemstones. The general formula of a spinel mineral is AB_2O_4 , where **A** and **B** are different metal ions. The presence of different combinations of metal ions accounts for the range of colours found in these gemstones. One example of a spinel gemstone is **hercynite**, which is blue-green in colour.

As they are essentially composed of metal oxides, spinel crystals may be reacted with strong acids to form solutions containing salts of the metal ions present. The rarity and high cost of spinel gemstones, however, prohibit the preparation of such solutions for this task.

The **FA 1** solution you are supplied with is a mixture of two salts. It is intended to duplicate a solution that could have been produced by reacting hercynite crystals with a strong mineral acid. Clearly, non-metal ions are also present in **FA 1**.

FA 1 is a solution containing the metal ions, **A** and **B**, together with non-metal ions.

FA 2 is sodium carbonate, Na_2CO_3 .

Solid FA 3 is granulated zinc.

You will perform tests to identify the metal ions, **A** and **B**, present in **FA 1**.

You are **not** expected to identify the anions.

You are advised to consider the general formula of the spinel minerals, AB_2O_4 , and the likely oxidation states of the metal ions, **A** and **B**, before you start this experiment.

Unless otherwise stated, the volumes given below are approximate and should be estimated rather than measured.

Test and identify any gases evolved.

(a) Carry out the following tests. Carefully record your observations in Table 1.1

Table 1.1

test		observations
1	Test the FA 1 solution using Universal Indicator paper.	Obs 1. pH 2–4 Or Yellow universal indicator paper changes to orange/red (1)
2	<p>Using a measuring cylinder, transfer 10 cm³ of FA 1 to a clean boiling tube.</p> <p>Use a dropping pipette to transfer 1 cm³ of FA 2 to the boiling tube.</p> <p>Mix the contents of the boiling tube thoroughly.</p> <p>Repeat the transfer of 1 cm³ of FA 2 with thorough mixing of the contents in</p>	<p>Obs 2. (slight) white ppt formed (1)</p> <p>Obs 3. effervescence/bubbles but not 'gas evolved' (1)</p> <p>Obs 4. gas evolved gives a white ppt with limewater (1)</p> <p>gas evolved is CO₂</p> <p>Obs 5. green/grey-green/blue-green/grey-green precipitate (1) (indicates initial precipitate becomes darker)</p> <p>Obs 6. then turns brown (this is observed at the inner side of the boiling tube) (1)</p>

	the boiling tube until all of the FA 2 is used up.	
3	<p>Place about 2 cm³ of FA 1 into a test-tube. Carefully add sodium hydroxide, dropwise with shaking, until no further changes are seen.</p> <p>Swirl and filter the mixture, collecting the filtrate in a clean boiling tube. Keep the filtrate for use in Test 6.</p> <p>Wash the residue thoroughly with deionised water. Discard the washings. Keep the residue for use in Test 4.</p>	<p>Obs 7. white ppt formed (soluble in excess NaOH) (1) Obs 8. then green/grey-green/blue-green/dirty green/light blue ppt formed (1) Obs 9. (green) ppt insoluble in excess NaOH (1)</p> <p>Obs 10. green residue/ppt (1) Obs 11. residue then then turns brown (1) Obs 12. colourless/pale yellow filtrate (1)</p>
Note: When aqueous potassium manganate(VII) is added in Test 4, the endpoint is a permanent pale pink colour.		
4	<p>Transfer a spatula load of the residue collected in Test 3 into a clean test-tube.</p> <p>Carefully add hydrochloric acid, dropwise with shaking, until no further changes are seen.</p> <p>Add aqueous potassium manganate(VII), dropwise with shaking, until the endpoint is reached.</p> <p>Record the number of drops of aqueous potassium manganate(VII) you have added to reach the endpoint.</p> <p>Keep this mixture for use in Test 5.</p>	<p>Obs 13. solid dissolves in acid to form an orange/orange-brown/brown/yellow solution (1)</p> <p>Obs 14. after <u>≥ 3 drops</u>, it turns pink/endpoint is reached (1)</p>
5	<p>To the final mixture in Test 4, add one piece of FA 3. Gently warm the mixture and set aside for a few minutes. From time to time, observe the mixture. While you are waiting, continue with Test 6.</p> <p>When no further changes are observed, decant the solution into a second test-tube.</p>	<p>Obs 15. the solution gradually lightens in colour/becomes colourless/orange solution turns yellow (1) Obs 16. many bubbles/effervescence but not 'gas evolved' (1) Obs 17. 'pops' with a lighted splint (1) gas evolved is H₂</p> <p>Obs 18. decolourisation of the purple KMnO₄(aq) (1) Obs 19. after a few drops, the solution darkens or turns orange/orange-brown/brown (1)</p>

	Add a few drops of aqueous potassium manganate(VII) to the second test-tube, with shaking.	
6	Place about 1 cm ³ of the filtrate collected in Test 3 in a clean test-tube. Carefully add hydrochloric acid, dropwise, until no further change is seen before you shake the mixture. The mixture is then use for Test 7.	Obs 20. white ppt forms (1) Obs 21. that dissolves in excess HCl (1) to form colourless solution
Note: You should NOT do the following test. Observations have been recorded for you.		
7	Place about 1 cm ³ of the final mixture in Test 6 in a clean test-tube. Add aqueous ammonia, dropwise with shaking, until no further change is seen.	white ppt. insoluble in excess

[8]

Obs points		marks	
------------	--	-------	--

Observation points

See the responses listed below. There are 21 observational points identified in Tests 1 to 6, each indicated by (1).

Record the total number of observational points awarded in the "Obs points" box at the end of question 1. Use the scaling grid below to determine the number of marks awarded. Write this number in the "marks" box at the end of question 1.

>= 18 = 8
 16 – 17 = 7
 13 – 15 = 6
 10 – 12 = 5
 7 – 9 = 4
 5 – 6 = 3
 3 – 4 = 2
 1 – 2 = 1

- (b) By consider your observations in Tests 1 and 2, suggest an explanation for the observations in Test 2 in terms of the chemistry involved.

.....
 [2]

Test 1 indicates the presence of acidic cation in **FA 1**.

The acidic cation (which is Al^{3+}) has high charge density, hence hydrolyses in water to produce acidic solution OR acidic cation has high charge density and polarises electron cloud of H_2O which weakens the O–H bond which is broken more easily to donate H^+ .

Na_2CO_3 reacts with the acid to produce CO_2 . [1]

White ppt is $\text{Al}(\text{OH})_3$, green ppt is FeCO_3 OR hydroxide and carbonate ppt is formed when carbonate reacts with metal ions in **FA 1**. [1] (FYI: The white ppt appeared first as it has a lower K_{sp} .)

- (c) (i) By considering your observations in Tests 3, 6 and 7, identify the metal-containing complex ion present in the filtrate collected in Test 3.

..... [1]

$[\text{Al}(\text{OH})_4]^-$ (aq) [1]

Explanation for understanding:

Al^{3+} ion present gives a white precipitate, $\text{Al}(\text{OH})_3$ with $\text{NaOH}(\text{aq})$, which is soluble in excess in Test 3 (complex ion formed is $[\text{Al}(\text{OH})_4]^-$ and a white precipitate, $\text{Al}(\text{OH})_3$ with $\text{NH}_3(\text{aq})$, which is insoluble in excess in Test 7.

- (ii) Identify the residue collected in Test 3.

..... [1]

$\text{Fe}(\text{OH})_2$

Accept $\text{Fe}(\text{OH})_3$ as oxidation by O_2 occurred/was observed.

- (iii) **FA 1** is intended to duplicate a solution that could have been produced by dissolving hercynite in a strong acid.

Suggest a formula for hercynite.

..... [1]

hercynite formula = FeAl_2O_4 [1] ecf so long charges balanced $(+2) + (+3)(2) \leftrightarrow (-2)(4)$.

- (d) Explain the following reactions involving the metal ions in the reaction mixture used in Test 5.
- reaction between **FA 3** and resultant mixture from Test 4
 - reaction between solution in the second test-tube and aqueous potassium manganate(VII)

.....
 [1]

Zn reduces Fe^{3+} to Fe^{2+}

MnO_4^- oxidises Fe^{2+} back to Fe^{3+} [1]

(hydrogen formed by reaction between Zn and the excess HCl acid (not acidic Fe^{3+}) added in Test 4)

- (e) Explain in terms of the chemistry involved and with the aid of relevant equations, your observations in Test 6.


.....
 [2]

- precipitate forms due to removal of OH^- by HCl, causing the POE of reaction (1) to shift to the left.
 $\text{Al}(\text{OH})_3 + \text{OH}^- \rightleftharpoons [\text{Al}(\text{OH})_4]^-$ ———(1) [1]
- precipitate dissolves due to further removal of OH^- by the excess HCl, causing the POE of reaction (2) to shift to the left
 $\text{Al}^{3+} + 3\text{OH}^- \rightleftharpoons \text{Al}(\text{OH})_3$ ———(2) [1]


[Total: 16]

2 Determination of the molar enthalpy change of a reaction by an indirect method

It is not possible to measure experimentally the enthalpy change, ΔH_1 , for the following reaction as it does not take place in the laboratory.



Sodium carbonate and sodium hydrogencarbonate each react with hydrochloric acid.



In this question, you will perform an experiment to determine a value for ΔH_3 . You will use the results of the experiments you have carried out, data provided for ΔH_2 and an energy cycle to calculate a 'theoretical' value for ΔH_1 .

You are provided with:

- **FA 4**, 2.00 mol dm^{-3} hydrochloric acid, HCl
- solid **FA 5**, NaHCO_3 .

(a) (i) Determination of the molar enthalpy change of reaction, ΔH_3

In this experiment, you will measure the temperature of the contents of a polystyrene cup at timed intervals, both before and after solid NaHCO_3 is added. You will analyse your results graphically in order to determine an accurate value for the temperature change of the mixture, caused by the reaction between solid NaHCO_3 and HCl.

You will use this value to calculate the heat change, q , for the experiment and hence determine a value for the molar enthalpy change of reaction, ΔH_3 .

In an appropriate format in the space provided on page 8, prepare tables in which to record results for your experiment:

- all weighings to an appropriate level of precision,
- all values of temperature, T , to an appropriate level of precision,
- all values of time, t , recorded to the nearest 0.5 min.

It is important that you measure each temperature at the specified time.

Procedure

1. Weigh the capped bottle containing solid **FA 5**. Record the mass in your table on page 8.
2. Place one polystyrene cup inside a second polystyrene cup. Place these in a glass beaker to prevent them from tipping over.
3. Use a measuring cylinder to transfer 50 cm^3 of **FA 4** into the first polystyrene cup. Cover the cup with a lid.
4. Carefully stir the solution of **FA 4** in the cup with the thermometer. Read and record the temperature, T . Start the stopwatch ($t = 0.0 \text{ min}$). The stopwatch must be left to run for the rest of the experiment.

5. Continue to stir the solution. Read and record T every minute.
6. At **exactly** three minutes, remove the lid and transfer the solid **FA 5** cautiously to the polystyrene cup in three separate lots, taking care that the mixture does not overflow. Replace the lid. Stir the mixture but do not read T .
7. Continue to stir the mixture. Read and record T at $t = 3.5$ min.
8. Continue to stir the mixture. Read and record T at $t = 4.0$ min and every minute until $t = 9.0$ min.
9. Reweigh the emptied bottle with its cap. Record this mass in your table below.

Results

[4]

Mass of bottle and solid FA 5 / g	a	10.009
Mass of bottle and residual FA 5 / g	b	5.512
Mass of FA 5 used / g	c	4.497

t / min	0.0	1.0	2.0	3.5	4.0	5.0	6.0	7.0	8.0	9.0
T / °C	29.2	29.2	29.2	23.4	23.2	23.4	23.6	23.6	24.0	24.2

1 mark

Tabulates mass **and** temperature data in **(a)**. Students should record all data in a table or tables. Given the large amount of data to be recorded, students should be allowed to record this data in multiple tables. All table(s) should contain correct headers and units. Data headers to be included in the table(s) are *as shown in the suggested answers above*.

'Table(s)' need not have lines but there should be no absences of headers or units.

Tabulation may be vertical or horizontal.

The mark is for drawing the table(s); the table(s) need not be populated.

1 mark

All temperature readings are recorded to 1 decimal place.

And

All mass readings and mass used values are recorded to 3 decimal places.

1 mark

Check the followings:

- 1) $(a) > (b)$
- 2) Correctly calculates $(a) - (b)$

1 mark


No temperature recording at $t = 3.0$ min + complete set of data

- (ii) Plot a graph of temperature, T , on the y -axis, against time, t , on the x -axis on the grid in Fig. 2.1.


Draw a best-fit straight line taking into account all of the points before $t = 3.0$ min.

Draw another best-fit straight line taking into account all of the points after the temperature of the mixture has started to rise steadily.

Extrapolate (extend) both lines to $t = 3.0$ min.

**Fig. 2.1**

[3]



1 mark

Axes correct way round + correct labels + units + scale.

*Note: Sensible scales must be used **and** must allow for the lines to be extrapolated to cross each other. Awkward scales (e.g. 3:10) are not allowed. The plotted points must occupy at least half of the graph grid in both x and y directions.*

1 mark

Plotting – all points correctly plotted within $\pm\frac{1}{2}$ small square.

Check all of the points and put ticks if correct.

1 mark

Graph lines must be best-fit lines, drawn so as to best reflect the distribution of points **before** and **after** the time = 3 minute. The line before and after time = 3 minute should be extrapolated until it **crosses** the vertical line drawn at time = 3 minute. The second line after time = 3 minute should be an increasing trend, not decreasing trend.

*Markers **must not** alter or adjust a poorly drawn student's line.*

- (iii) From your graph, read the minimum temperature, T_{\min} , and the maximum temperature, T_{\max} , at $t = 3.0$ min. Record these values in the spaces provided.

Calculate the temperature change, ΔT , at $t = 3.0$ min.

T_{\min} =
 T_{\max} =
 ΔT = [2]

$$T_{\min} = 23.05\text{ }^{\circ}\text{C}$$

$$T_{\max} = 29.20\text{ }^{\circ}\text{C}$$

$$\Delta T = 6.15\text{ }^{\circ}\text{C}$$

1 mark

Correct reading, to $\pm\frac{1}{2}$ small square, of T_{\min} , T_{\max} **and** correct calculation of ΔT

Could be 1 d.p or 2 d.p; be consistent

1 mark

Compared $\Delta T/m$ of student with $\Delta T/m$ of Teacher.

Ensure $\Delta T/m$ of student is rounded to 1 d.p before checking accuracy

Award 1 mark for accuracy if student's $\Delta T/m$ is ± 0.2 of $\Delta T/m$ of Teacher

- (iv) Calculate the heat change, q , for your experiment using the value you deduced in (a)(iii).

Assume that the specific heat capacity of the reaction mixture is $4.18\text{ J g}^{-1}\text{ K}^{-1}$, and that the density of the reaction mixture is 1.00 g cm^{-3} .

$q = \dots\dots\dots$ [1]

$$q = 50 \times 1.00 \times 4.18 \times 6.15 \times 10^{-3} = 1.285 \approx 1.29\text{ kJ}$$

units can be in J or kJ

(v) Determine the molar enthalpy change, ΔH_3 , for the reaction in equation 3.

Include the sign of ΔH_3 in your answer.


[Ar: Na, 23.0, H, 1.0, C, 12.0, O, 16.0]

$\Delta H_3 = \dots\dots\dots$ [2]

$$\Delta H_3 = + \frac{1.285}{\frac{4.497}{84.0}} = +24.00 \approx +24.0 \text{ kJ mol}^{-1}$$

- Knowing that the limiting reagent is NaHCO_3 (accept no calculation to prove this) and hence calculating $n(\text{NaHCO}_3)$
- Calculate the numerical value of ΔH_2 correctly by taking the ans in (iv) divide by $n(\text{NaHCO}_3)$
- Include the correct sign and correct units
3 pts – 2 marks, 2 pts – 1 mark

- (b) Use the value of ΔH_2 given and your value of ΔH_3 calculated in (a)(v), calculate a value for ΔH_1 for the reaction in equation 1.



$$\Delta H_1 = -36.0 - 2(+24.00) = -84.00 \approx -84.0 \text{ kJ mol}^{-1}$$

$$\Delta H_1 = -84.0 \text{ kJ mol}^{-1} \dots\dots\dots [2]$$

- energy cycle
 - equations correctly balanced and with state symbols
 - arrows correctly labelled
- applying Hess' Law correctly
- correct sign + units + correct numerical answer presented to 3 s.f. for (a)(iv), (v) and (b)

Or

- applying $\Delta H_1 = \Delta H_2 - 2\Delta H_3$
- correct numerical answer + units
- 3 s.f. for (a)(iv), (v) and (b)

3 marking points – 2 marks, 2 marking points – 1 mark

- (c) A student performed the same experiment using 40 cm³ of **FA 4**.

Explain how this will affect the temperature change per gram of **FA 5** and hence the accuracy of the experiment.

.....
 [2]

$$\Delta H = \frac{V \times \rho \times c \times \Delta T}{\frac{m}{M_r}} \text{ where } V \text{ is the volume of FA 4 used and } m \text{ is the mass of FA 5 used}$$




From the above relationship, $V \propto \frac{1}{\Delta T/m}$ (or $\frac{\Delta T}{m} \propto \frac{1}{V}$), hence, when volume of FA 4 decreased, temperature change per gram of FA 5 used will increase. [1]

Since percentage error $\propto \frac{1}{\Delta T}$, the percentage error will decrease and increase the accuracy of the experiment. [1]


[Total: 16]

3 Analysis of a redox reaction between hydroxylamine and iron(III) ion

A redox reaction takes place between hydroxylamine, NH_2OH , and iron(III) ion, Fe^{3+} , in acidic conditions. Iron(III) ion is reduced to an iron(II) ion, Fe^{2+} . The reaction is slow at room temperature but is completed in a few minutes at $100\text{ }^\circ\text{C}$. The reaction is shown by one of the following equations.



You will carry out a titration to determine which of the three suggested equations best represents the reaction. Iron(II) ions formed in the reaction with hydroxylamine are oxidised by manganate(VII) ions.



FA 6 is $0.0200\text{ mol dm}^{-3}$ potassium manganate(VII), KMnO_4 .

FA 7 is a solution prepared by boiling a 1.00 dm^3 aqueous mixture containing 3.30 g of hydroxylamine hydrochloride, $\text{NH}_2\text{OH}\cdot\text{HCl}$, excess iron(III) chloride, FeCl_3 , and excess sulfuric acid. Any water lost by evaporation was replaced after cooling.

FA 8 is dilute sulfuric acid.

Assume that one mole of hydroxylamine hydrochloride gives one mole of hydroxylamine in solution.

(a) (i) Procedure

1. Fill the burette with **FA 6**.
2. Pipette 25.0 cm^3 of **FA 7** into a conical flask.
3. Use a 10 cm^3 measuring cylinder, add 10 cm^3 of **FA 8** into the same conical flask.
4. Titrate the solution in the conical flask with **FA 6**. The end-point is reached when a permanent pale pink colour is obtained.
5. Record your titration results to an appropriate level of precision in the space on page 13.
6. Repeat steps 2 to 5 until consistent results are obtained.

Titration results

initial burette reading / cm ³	0.00	0.00
final burette reading / cm ³	24.55	24.50
volume of FA 6 added / cm ³	24.55	24.50
consistent results	√	√

[3]

1 mark

Tabulates initial and final burette readings and volume added in the titration table.

Table has correct headers and units.

Tabulation may be vertical or horizontal; lines are not essential but there should be no absences of headers.

Where units have not been included in the header, there should be appropriate unit for each entry in the table.

Do NOT award this mark if any final and initial burette readings are inverted or 50 is used as the initial burette reading.

1 mark

All the final/initial burette readings, for all accurate titres in the titration table, are recorded to the nearest 0.05 cm³.

Treat all titres as “accurate” unless labelled ‘rough’ or first titre is recorded to a lower precision than subsequent titres.

1 mark

Has at least two uncorrected titres for end-point within 0.10 cm³

Note:

- Do not award this mark if having performed two titres within 0.1 cm³ a further titration is performed which is more than 0.10 cm³ from the closer of the initial two titres, unless a fourth titration, within 0.1 cm³ of the third titration or of the first two titres has also been carried out. [9701_w10_ms31]*
- A student's ‘rough’/‘trial’ titre value can be considered by the examiner when selecting titre values for the mean titre calculation **if** the student has ‘validated’ this value either by ticking it or by using it in an expression in **(a)(ii)**. (By doing either of these, the student has declared it to be no longer a ‘rough’/‘trial’ value).*

- (ii) From your titrations, obtain a suitable volume of **FA 6** to be used in your calculations. Show clearly how you obtained this volume.

$$\text{volume of FA 6} = \frac{24.55 + 24.50}{2} = 24.525 \approx 24.53 \text{ cm}^3$$

volume of **FA 6** = [3]

1 mark

Student obtains appropriate “average”, to 2 d.p., from any experiments with uncorrected end-point titre values within 0.20 cm³ with units “cm³” included.

Do not award this mark if the titres used are not identified either in the table (by, for example, a tick) or in a calculation.

*Do **not** award this mark if there are arithmetic errors in the table.*

2 marks

Round burette readings to the nearest 0.05 cm³. Check and correct titre subtractions where necessary. Marker selects the best mean titre.

Apply hierarchy: 2 identical, titres within 0.05cm³, titres within 0.10cm³ etc. Examiner subtracts (corrected) candidate's titre from Supervisor's titre, δ .

Award accuracy marks as follows:

Award if $\delta \leq 0.30 \text{ cm}^3 \rightarrow 1 \text{ mark}$

Award if $\delta \leq 0.20 \text{ cm}^3 \rightarrow 2 \text{ marks}$

- (b) (i) Calculate the amount in moles of potassium manganate(VII) present in the volume of **FA 6** in (a)(ii).

$$n(\text{KMnO}_4) = \frac{24.53}{1000} \times 0.0200 = 4.906 \times 10^{-4} \approx 4.91 \times 10^{-4} \text{ mol}$$

amount of potassium manganate(VII) = 4.91×10^{-4} mol [1]

Correctly calculates amount of $\text{KMnO}_4 = 0.0200 \times \text{(a)(ii)} / 1000 \text{ mol}$

- (ii) Use your answer to (b)(i) to calculate the amount in moles of iron(II) ions in 25.0 cm³ of solution **FA 7**.

$$n(\text{Fe}^{2+}) = 4.906 \times 10^{-4} \times 5 = 2.453 \times 10^{-3} \approx 2.45 \times 10^{-3} \text{ mol}$$

amount of iron(II) ions = 2.45×10^{-3} mol [1]

Correctly calculates amount of $\text{Fe}^{2+} = 5 \times \text{(b)(i)} \text{ mol}$

- (iii) Calculate the amount in moles of hydroxylamine hydrochloride that has reacted with the **FA 7** pipetted into the conical flask. Show your working. [A: N, 14.0; H, 1.0; O, 16.0; Cl, 35.5]

$$n(\text{NH}_2\text{OH}\cdot\text{HCl}) = \frac{25.0}{1000} \times \frac{3.30}{69.5} = 1.187 \times 10^{-3} \approx 1.19 \times 10^{-3} \text{ mol}$$

amount of hydroxylamine hydrochloride = 1.19×10^{-3} mol [2]

Correctly calculates

M1 $M_r(\text{NH}_2\text{OH}\cdot\text{HCl}) = 69.5$; award credit if calculate correctly and 1 d.p

M2 amount of $\text{NH}_2\text{OH}\cdot\text{HCl} = 3.3/69.5 \times 25.0/1000 = 1.187 \times 10^{-3} / 1.19 \times 10^{-3} \text{ mol}$

- (iv) Use your answer to **(b)(iii)** to deduce which of the three suggested equations corresponds to your results. Show your working.

$$\frac{n(\text{Fe}^{2+})}{n(\text{NH}_2\text{OH})} = \frac{2.453 \times 10^{-3}}{1.187 \times 10^{-3}} = 2.066 \approx 2$$

The correct equation number is 2..... [2]


M1 Correctly uses mole ratio $n(\text{Fe}^{2+}) : n(\text{NH}_2\text{OH}\cdot\text{HCl}) = (\text{b})(\text{ii}) / (\text{b})(\text{iii})$ AND states equation number that fits their data

M2 final ans for part (b)(i) to (iii) given to 3 significant figures

[Total: 13]

4 Planning

Baker's yeast is a useful enzyme which can be used to catalyse the decomposition of hydrogen peroxide.



You are provided with

apparatus:

- 100 cm³ conical flask, fitted with rubber bung and rubber tubing
- gas syringe (possible capacities of 10, 50 or 100 cm³)
- stopwatch
- all other common laboratory equipment

reagents:

- 50 cm³ of yeast suspension
- 30 cm³ of 3% (by weight) hydrogen peroxide
- Distilled water

When 8.0 cm³ of the yeast suspension, 4.0 cm³ of H₂O₂ and 18.0 cm³ of distilled water were mixed, 10 cm³ of oxygen gas was produced in 90 s.

Use the above information and the reagents provided to design an experiment to determine the order of reaction with respect to yeast and hydrogen peroxide respectively.

In your experiment, you should perform a total of five experimental runs (including the above run) to measure the volume of oxygen gas produced at regular intervals in each run. The volume of yeast suspension used should be varied in all the runs while keeping volume of H₂O₂ constant at 4.0 cm³. One of the experimental runs should be conducted to allow the reaction to go to completion.

- (a) Calculate the maximum volume of oxygen gas produced for an experimental run if the reaction was allowed to go to completion. In your calculations, assume that the density of H₂O₂ is 1.00 g cm⁻³ and the experiment is conducted under r.t.p. condition.

[A_r: H, 1.0; O, 16.0 and V_m = 24 dm³ mol⁻¹ at r.t.p.]

maximum volume of oxygen gas produced =

Hence, state the appropriate capacity of the gas syringe to be used.

capacity of gas syringe =

[2]

- Mass of 4.0 cm³ of H₂O₂ = 1.00 x 4.0 = 4.000 g
- Amount of H₂O₂ = (4.000 x 0.03) / 34.0 = 0.00353 mol
- Amount of O₂ gas = ½ (0.00353) = 0.00176 mol
- Volume of O₂ gas = 0.00176 x 24000 = 42.4 cm³
- Hence appropriate capacity of gas syringe to be used is either 50 cm³ or 100 cm³ size if the reaction is allowed to go to completion

2 pts – 1 mark, 5 pts – 2 marks

- (b) Plan an experiment to collect sufficient data to plot suitable graphs to prove that the reaction is first order with respect to hydrogen peroxide and first order with respect to yeast.

Your plan should include details of:

- a table to tabulate the volumes of each of the reagents used that you will be using for each run of the experiment
- the apparatus you would use
- the procedure you would follow
- the measurement you would take

You may find it useful to indicate in your procedure the experimental run that would be conducted to allow the reaction to go to completion and the measurement to be taken when the reaction has reached completion.

[5]

Table of volumes

Run	V(yeast) /cm ³	V(H ₂ O ₂) /cm ³	V(distilled water) /cm ³	Total volume /cm ³ (optional to show this column)
1	4.0	4.0	22.0	30.0
2	6.0	4.0	20.0	30.0
3	8.0	4.0	18.0	30.0
4	10.0	4.0	16.0	30.0
5	12.0	4.0	14.0	30.0

- volume of yeast is between 1 – 20 cm³; total volume of yeast provided for all expts is ≤ 50cm³
 - volume of H₂O₂ is constant at 4.0 cm³
 - volume of distilled water added accordingly to keep total volume constant
- 2 pts – 1 mark, 3 pts – 2 marks

Procedure for measuring the volume of oxygen gas produced at fixed intervals

1. Connect the rubber tubing to the gas syringe.
2. Record the initial reading on the syringe.
3. Using a 25 cm³ measuring cylinder, measure 4.0 cm³ of the yeast suspension and add into the conical flask
4. Using a 25 or 50 cm³ measuring cylinder, measure 22.0 cm³ of the distilled water and add into the conical flask.
5. Using a 10 cm³ measuring cylinder, measure 4.0 cm³ of the H₂O₂.
6. Pour the H₂O₂ into the conical flask, stopper the conical flask with the rubber bung and start the stop watch immediately. Swirl the conical flask a few times to ensure complete mixing.
7. Record the reading on the syringe every 10 seconds (or other regular time intervals, until 5 readings are collected)
8. Repeat steps 2 – 7 for the next 4 experimental runs, changing the volumes according to the table given.
9. For run 5 (or any other run), allow the reaction to go to completion i.e. no more gas produced. Note the final reading on the syringe.

3 steps – 1 mark, 4-6 steps – 2 marks, 7-9 steps – 3 marks

- (c) (i) Sketch a graph you would expect to show how the volume of O_2 will vary with time on Fig. 4.1. Using the maximum volume of oxygen gas calculated in (a), describe how you would use your graph to prove that the reaction is first order with respect to hydrogen peroxide.





Fig. 4.1

[2]



- Correct shape of graph
 - Since it is first order reaction with respect to hydrogen peroxide, find 2 consecutive half-lives and the half-lives should be constant i.e. $t_1 = t_2$.
 - Correct construction lines to show half-lives and the labelling.
- 3 pts – 2 marks, 2 pts – 1 mark

- (ii) In Fig. 4.1 above, show how the initial rate can be determined. Hence, sketch the graph you would expect on Fig. 4.2 to prove that the reaction is first order with respect to yeast.





Fig. 4.2

[2]

[1] for drawing the tangent on Fig. 4.1



[1]

[Total: 11]

BLANK PAGE

BLANK PAGE

Qualitative Analysis Notes

[ppt. = precipitate]

(a) Reactions of aqueous cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	ammonia produced on heating	–
barium, Ba ²⁺ (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt., turning brown on contact with air insoluble in excess	green ppt., turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt., rapidly turning brown on contact with air insoluble in excess	off-white ppt., rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

(b) Reactions of anions

<i>anion</i>	<i>reaction</i>
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$)
bromide, $\text{Br}^-(\text{aq})$	gives pale cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$)
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$)
nitrate, $\text{NO}_3^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil
nitrite, $\text{NO}_2^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil; NO liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown NO_2 in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	SO_2 liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in dilute strong acids)

(c) Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	"pops" with a lighted splint
oxygen, O_2	relights a glowing splint
sulfur dioxide, SO_2	turns aqueous acidified potassium manganate(VII) from purple to colourless

(d) Colour of halogens

<i>halogen</i>	<i>colour of element</i>	<i>colour in aqueous solution</i>	<i>colour in hexane</i>
chlorine, Cl_2	greenish yellow gas	pale yellow	pale yellow
bromine, Br_2	reddish brown gas / liquid	orange	orange-red
iodine, I_2	black solid / purple gas	brown	purple