

Anglican High School Secondary Four Preliminary Examination 2020



CANDIDATE NAME		
CLASS 4 CLASS	S INDEX	
PHYSICS		6091/03
Paper 3 Practical Test		24 July 2020
	1 ho	ur 50 minutes
Candidates answer on the Question Paper.		
Additional Materials: As listed in the Confidential Instructions.		
READ THESE INSTRUCTIONS FIRST		
Write your Centre number, index number and name on all the work you hand in. Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid.		
Answer all questions. All of your answers should be written in this Question Paper: scrap paper must not Graph paper is provided in this Question Paper. Additional sheets of graph paper enecessary to do so.		ised only if it is
You will be allowed to work with the apparatus for a maximum of 55 minutes for each You are expected to record all your observations as soon as they are made. An account of the method of carrying out the experiments is not required.	ch section.	
The use of an approved scientific calculator is expected, where appropriate.		
At the end of the examination, fasten all your work securely together.		
The number of marks is given in brackets [] at the end of each question or part question.	For Exa	miner's Use
	1	
	2	
	3	
	Total	

This document consists of 13 printed pages.

Section A

1 In this experiment, you will investigate the amplitude of a simple pendulum.

You are provided with

- · a pendulum bob attached to a length of string,
- a split cork,
- · a stand, boss and clamp,
- a stopwatch,
- · a metre rule,
- · a counterweight to prevent the stand from toppling,
- a piece of card marked with a series of parallel lines, drawn at 2cm intervals.

The apparatus has been set up for you, as shown in Fig. 1.1(a) and Fig. 1.1(b).




Fig. 1.1(a) (front view)

Fig. 1.1(b) (side view)

- (a) (i) The distance from the bottom of the split cork to the centre of the pendulum bob is *H*. Loosen the clamp and adjust the string until *H* is equal to 1.25 m. Tighten the clamp.
 - (ii) Adjust the boss so that the distance h from the bottom of the pendulum bob to the floor is 2.0 cm.

(b) View the pendulum from the front, as shown in Fig. 1.1(a). Move the pendulum bob a few centimetres to the side. Release the pendulum bob so that it swings freely.

Determine an accurate value for the period *T* of the pendulum.

		٠
Г_	כיו	ŀ
_	 -	ı

(c) (i) Stop the pendulum.

Place the piece of card on the floor beneath the pendulum bob. The lines on the card are drawn at 2cm intervals.

Adjust the position of the card until the right hand edge of the pendulum bob is aligned with the 0cm mark on the card, when viewed from above, as shown in Fig. 1.2.




Fig. 1.2 (view from above)

(ii) Move the pendulum bob to the side until the right hand edge of the bob is aligned with the 12 cm mark on the card, as shown by the dotted line in Fig. 1.2.

Release the pendulum bob. Measure and record the time t for the amplitude of the pendulum to decrease from 12 cm to 10 cm.

	[=[1]
(iii)	Suggest why the amplitude of the pendulum decreases.

.....[1]

(d)	Plan

A student suggests that the time taken for the amplitude to decrease, as in (c)(ii), is directly proportional to the mass of the pendulum bob.

Plan an experiment to investigate this relationship.

Your plan should include

- the quantities that you will keep constant,
- a detailed description of how you will perform the investigation,
- an indication of how you will achieve accurate results,
- a statement of the graph you would plot to test the relationship,

a sketch of the graph you would obtain if the suggested relationship is correct.

	[C]

2 In this experiment, you will investigate the extension of a spring.

You have been provided with

- · a length of string tied to a spring,
- an S-hook,
- a stand, boss and clamp,
- · a piece of plastic pipe,
- a mass of weight 1.5 N,
- · a set square,
- a 30 cm ruler,
- a 50 cm rule,
- a metre rule fixed to the bench with Blu-tac,
- a loop of string on the metre rule.
- (a) (i) Measure and record the unstretched length l_0 of the spring, as shown in Fig. 2.1.




Fig. 2.1

$$l_0 = \dots [1]$$

(ii) Place one of the loops of the spring onto the rod of the clamp, as shown in Fig. 2.2. Suspend the mass on the other loop.




Fig. 2.2

Measure and record the stretched length $l_{\rm s}$ of the spring. Determine the extension $e_{\rm o}$ of the spring using the equation $e_{\rm o} = l_{\rm s} - l_{\rm o}$.

$l_s =$	
e _o =	 [2]

(b)	Use your value of e_0 from (a)(ii) and the equation $F = ke_0$, where $F = 1.5$ N, to determine the
	spring constant <i>k</i> for the spring.

-		14	1
- 1	***************************************	11	1

(c) Remove the spring from the rod of the clamp. Place the plastic pipe in the clamp. Adjust the height of the boss so that the centre of the plastic pipe is 50 cm above the bench. Adjust the position of the clamp stand so that the centre of the plastic pipe is aligned with the 50 cm mark on the metre rule.

Attach the S-hook to the loop of string on the metre rule. Attach the spring to the S-hook. Place the string over the plastic pipe and suspend the mass from the loop at the end of the string, as shown in Fig. 2.3.




Fig. 2.3

	(i)	Adjust the position of the loop of string on the metre rule until the spring is vertical.
		Measure and record the stretched length l_1 of the spring. Determine the extension e_1 of the spring using the equation $e_1 = l_1 - l_0$.
		,
		<i>l</i> ₁ =
		e ₁ =[1]
	(ii)	With the mass still suspended from the string, carefully move the loop of string on the metre rule so that it is at the 15 cm mark.
		Measure and record the stretched length l_2 of the spring. Determine the extension e_2 of the spring using the equation $e_2 = l_2 - l_0$.
		<i>l</i> ₂ =
		e ₂ =
		[1]
(d)	Use	your value of k , values of e_1 and e_2 and the equation from (b) to calculate the forces F_1 F_2 in the spring with the loop in the two different positions on the metre rule.
		<i>F</i> ₁ =
		$F_2 = \dots $ [2]
(e)	Exn	lain why neither F_1 nor F_2 are equal to the weight of the mass on the spring.
(0)	ΕλΡ	itali why hold of the regard to the weight of the mass on the spining.
		[2]

Section B

3 In this experiment, you will investigate an electrical circuit containing a light-dependent resistor (LDR).

You are provided with

- a lamp,
- a light-dependent resistor (LDR),
- a power supply,
- a fixed resistor of resistance 470 Ω,
- a ruler,
- a wooden rod,
- a stand and boss,
- Blu-Tack,
- a switch,
- a voltmeter and two connecting leads,
- eight connecting leads.

Set up the circuit as shown in Fig. 3.1.




Fig. 3.1

Set up the light-dependent resistor (LDR) and lamp as shown in Fig. 3.2. Secure the LDR to the wooden rod using Blu-Tack. The rod can be moved up and down to change the distance between the LDR and the lamp.





Fig. 3.2

Adjust the position of the wooden rod so that the LDR is directly above the lamp and the distance d is zero.

(a)	(i)	Connect the voltmeter across the terminals of the power supply. Close the switch.
		Record the voltmeter reading $V_{\rm s}$.
		V _s =[1]
	(ii)	Open the switch. Disconnect the voltmeter from the power supply. Connect the voltmeter across points A and B. Close the switch.
		Record the voltmeter reading $V_{\rm AB}$.
		V _{AB} =[1]
(b)	Op	en the switch.
	The	e current I in the fixed resistor is given by the equation
		$I = \frac{V_{AB}}{R}$
	wh	ere $R = 470 \Omega$.
	Ca	culate I.
		<i>I</i> =[1]
(c)	Ca	Iculate the resistance $R_{\rm L}$ of the LDR using the equation
		$R_{\rm L} = \frac{V_{\rm L}}{I}$
	wh	ere the potential difference (p.d.) V_L across the LDR is given by $V_L = (V_s - V_{AB})$.
		R _L =[1]

(d)	(i)	Adjust the position of the wooden rod so that the distance d between the LDR and the top of the lamp is 1.0 cm.
		Close the switch. Record the new voltmeter reading $V_{\rm AB}$.
		V _{AB} =
	(ii)	Open the switch.
		Repeat (b) and (c) to calculate the new current I in the resistor and the new resistance $R_{\rm L}$ of the LDR. Use your value of $V_{\rm s}$ from (a)(i).
		<i>I</i> =
		R _L =[2]
	_	
(e)		peat (d) for further values of d.
	Re	cord your results for d , V_{AB} , V_{L} , I and R_{L} . Include all of your results.
		[6]
,	_	•
(f)	Oı	the grid plot a graph of $R_{\rm L}$ against d .



(g)	(i)	Determine the change in resistance $R_{\rm L}$ between $d=1.0{\rm cm}$ and $d=4.0{\rm cm}$.
		change in resistance =[1]
	(ii)	Calculate the average change in resistance per cm of the LDR between $d = 1.0 \mathrm{cm}$ and $d = 4.0 \mathrm{cm}$.
		average change in resistance per cm =[1]
	(iii)	Explain how the graph shows that the average change in resistance per cm decreases as d increases.
		[1]
(h)	(h) A student has a voltmeter that gives readings to 0.1 V.	
	Explain why the values for the change in resistance $R_{\rm L}$ obtained when d is large would not be accurate using the voltmeter.	
		,
		[1]
		[Total: 20]

End of paper -

This paper is adapted from the 2018 & 2019 GCE O Level Physics Examinations.