

NANYANG JUNIOR COLLEGE
JC 2 PRELIMINARY EXAMINATION
Higher 2

CANDIDATE
NAME

--

CLASS

--

TUTOR'S
NAME

--

CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

PHYSICS

9749/01

Paper 1 Multiple Choice

20 September 2022

1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write your name, class, Centre number and index number in the spaces at the top of this page.

There are **thirty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in **soft pencil** on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

The use of an approved scientific calculator is expected, where appropriate.

This document consists of **16** printed pages.

Data

speed of light in free space

permeability of free space

permittivity of free space

elementary charge

the Planck constant

unified atomic mass constant

rest mass of electron

rest mass of proton

molar gas constant

the Avogadro constant

the Boltzmann constant

gravitational constant

acceleration of free fall

$$c = 3.00 \times 10^8 \text{ m s}^{-1}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$$

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$$

$$(1 / (36\pi)) \times 10^{-9} \text{ F m}^{-1}$$

$$e = 1.60 \times 10^{-19} \text{ C}$$

$$h = 6.63 \times 10^{-34} \text{ J s}$$

$$u = 1.66 \times 10^{-27} \text{ kg}$$

$$m_e = 9.11 \times 10^{-31} \text{ kg}$$

$$m_p = 1.67 \times 10^{-27} \text{ kg}$$

$$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$$

$$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$$

$$g = 9.81 \text{ m s}^{-2}$$

Formulae

uniformly accelerated motion

$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

$$W = p\Delta V$$

$$p = \rho gh$$

$$\phi = -Gm/r$$

$$T / \text{K} = T / ^\circ\text{C} + 273.15$$

$$p = \frac{1}{3} \frac{Nm}{V} < c^2 >$$

$$E = \frac{3}{2} kT$$

$$x = x_0 \sin \omega t$$

$$v = v_0 \cos \omega t$$

$$= \pm \omega \sqrt{x_0^2 - x^2}$$

$$I = Anvq$$

$$R = R_1 + R_2 + \dots$$

$$1/R = 1/R_1 + 1/R_2 + \dots$$

$$V = \frac{Q}{4\pi\epsilon_0 r}$$

$$x = x_0 \sin \omega t$$

$$B = \frac{\mu_0 I}{2\pi d}$$

$$B = \frac{\mu_0 NI}{2r}$$

$$B = \mu_0 nI$$

$$x = x_0 \exp(-\lambda t)$$

$$\lambda = \frac{\ln 2}{t_{\frac{1}{2}}}$$

electric current

resistors in series

resistors in parallel

electric potential

alternating current/voltage

magnetic flux density due to a long straight wire

magnetic flux density due to a flat circular coil

magnetic flux density due to a long solenoid

radioactive decay

decay constant

- 1 Which of the following is not a reasonable estimate?
- A Sound of frequency in the order of 10^9 Hz is audible to human beings.
 - B Light of wavelength in the order of 10^{-7} m is visible to human beings.
 - C The volume of an average-sized orange is in the order of 10^{-4} m³.
 - D The speed of beta particles emitted from a nucleus is in the order of 10^7 m s⁻¹.
- 2 The kinetic energy E_k of a rotating solid sphere is given by the following expression


$$E_k = \frac{2\pi^2 I}{T^2}$$

where I is the moment of inertia and T is the period of rotation of the solid sphere.

What is the unit for the moment of inertia I , expressed in SI base units?

- A kg m² s⁻¹
- B kg m²
- C J s²
- D N m s²

- 3 A ruler is supported horizontally by two pivots as shown.



The vertical displacement y at the centre of the ruler can be used to measure the mass loaded on it and is given by the equation

$$y = \frac{kML^3}{wt^3}$$

where

k is a constant,

L is the distance between the pivots,

M is the mass loaded onto the ruler,

t is the thickness of the ruler and

w is the width of the ruler.

When a particular M is loaded onto the ruler, the following results are obtained:

$$y = (0.25 \pm 0.01) \text{ mm}$$

$$L = (80.0 \pm 0.2) \text{ cm}$$

$$t = (6.0 \pm 0.1) \text{ mm}$$

$$w = (23.0 \pm 0.5) \text{ mm}$$

Which measurement contributes the most to the uncertainty of M ?

- A** y **B** L **C** t **D** w


- 4 The speed of an aeroplane in still air is 200 km h^{-1} . The wind pushes it from the west at a speed of 85.0 km h^{-1} .

In which direction must the pilot steer in order to fly due north?

- A** 23.0° east of north
B 23.0° west of north
C 25.2° east of north
D 25.2° west of north

- 5 A radio-controlled toy car travels along a straight line for a time of 15 s.

The variation with time t of the velocity v of the car is shown below.



What is the average velocity of the toy car for the 15 s journey?

- A** -1.5 m s^{-1} **B** 0.0 m s^{-1} **C** 4.0 m s^{-1} **D** 4.5 m s^{-1}

- 6 A tennis ball is thrown horizontally in air from the top of a tall building. The effect of air resistance is not negligible.

Which of the following correctly describes the subsequent change in the horizontal and vertical components of the ball's velocity?


	horizontal component of velocity	vertical component of velocity
A	constant	constant
B	constant	increases at a constant rate
C	decreases to zero	increases at a constant rate
D	decreases to zero	increases to a maximum value

- 7 A rock of mass $2m$ in deep space, initially travelling at velocity v , explodes into two parts of equal mass, one of which becomes stationary immediately after the explosion.

What is the kinetic energy of the moving part after the explosion?

- A** $\frac{1}{2}mv^2$ **B** mv^2 **C** $\frac{3}{2}mv^2$ **D** $2mv^2$

- 8 A single horizontal force F is applied to a block X which is in contact with a separate block Y, as shown.




The blocks remain in contact as they accelerate along a horizontal frictionless surface. X has a greater mass than Y.

Which statement is correct?


- A The acceleration of X is equal to force F divided by the mass of X.
 - B The force that X exerts on Y is equal to F .
 - C The force that X exerts on Y is less than F .
 - D The force that X exerts on Y is less than the force that Y exerts on X.
- 9 A full barrel of weight W is being rolled up a ramp. The force P is required to hold the barrel at rest on the ramp. Friction between the barrel and the ramp stops the barrel from slipping.

Which arrow shows the resultant force that the ramp exerts on the barrel?




- 10** An isolated disc is subjected to three forces, each given in terms of units of magnitude F .

In which situation will the disc experience both a resultant force and a resultant torque?




- 11** A block is being pulled up a rough inclined plane using a rope at a constant speed.



Which of the following statements is correct?


- A** The weight of the block does no work because the block is not moving in the direction of its weight.
- B** The work done by the force of the rope is equal to the work done against the frictional force from the plane because there is no gain in kinetic energy.
- C** The normal reaction force from the plane does positive work because it has an upward component and the block is moving upwards.
- D** The negative work done by the weight of the block is equal to the gain in gravitational potential energy of the block.

- 12 The figure below shows a light inextensible rope that passes over a light smooth pulley with two blocks of masses 4.0 kg and 2.0 kg attached to its two ends. The two blocks are initially at rest with the 4.0 kg block 10.0 cm above the 2.0 kg block.



What is the speed of the blocks when they are at the same height above the ground?

- A** 0.57 m s^{-1} **B** 0.81 m s^{-1} **C** 0.99 m s^{-1} **D** 1.4 m s^{-1}
- 13 A rod PR of length L is turned about the point P through an angle θ .




The end R of the rod moves through a distance D . Both D and L are measured in metres.

What is the angle θ , expressed in radians?

- A** L/D **B** D/L **C** DL **D** $1/(DL)$

- 14 The gravitational potential ϕ along the line joining the centres of a planet and its moon varies with the distance r from the centre of the moon as shown.



Which of the following expressions gives a value that is closest to that of the gravitational force acting on a 1 kg mass at a distance of r_2 ?

- A $\frac{\phi_1 - \phi_2}{r_2 - r_1}$
- B $-\frac{\phi_2}{r_2}$
- C $\frac{\phi_1 - \phi_3}{r_3 - r_1}$
- D $\frac{\phi_1 - \phi_4}{r_4 - r_1}$


- 15 Earth has a mass M and radius R . X is a point $5R$ from the center of the Earth.

An object of mass m falls freely from rest at X and hits the surface of the Earth.

Which of the following statements is false?

- A The change in gravitational potential is $\frac{4}{5} \frac{GM}{R}$.
- B The work done by the gravitational field is $\frac{4}{5} \frac{GMm}{R}$.
- C The speed of impact is $\sqrt{\frac{8}{5} \frac{GM}{R}}$.
- D The change in the magnitude of gravitational field strength is $\frac{24}{25} \frac{GM}{R^2}$.

- 16 The diagram shows the displacement-time graphs of two sound waves P and Q at a point in space. The graphs have the same scales for the time axes.



The frequency of Q is 125 Hz. The waves are in phase at time = 0.

At what time are the waves next in phase?

- A 32 ms B 36 ms C 64 ms D 72 ms
- 17 Which of the following correctly states the changes, if any, in the potential energy and kinetic energy of the molecules of a solid as it melts?

	potential energy	kinetic energy
A	decreases	increases
B	increases	remains the same
C	remains the same	decreases
D	remains the same	remains the same

- 18 The frequency of a wave is 600 Hz and its speed is 330 m s⁻¹.

What is the phase difference between the oscillation of two points on the wave that are 0.275 m apart?


- A 0 B $\frac{\pi}{4}$ rad C $\frac{\pi}{2}$ rad D π rad

- 19 A point source emits 50.0 W of sound energy in all directions. A small microphone of area 0.85 cm^2 detects the sound at 4.0 m from the source.

What is the power received by the microphone?


- A $1.6 \times 10^{-5} \text{ W}$
- B $2.1 \times 10^{-5} \text{ W}$
- C $2.1 \times 10^{-1} \text{ W}$
- D $2.5 \times 10^{-1} \text{ W}$

- 20 The diagram shows a standing wave on a string. The standing wave has three nodes N_1 , N_2 and N_3 .



Which statement is correct?


- A All points on the string vibrate in phase.
 - B All points on the string vibrate with the same amplitude.
 - C Points equidistant from N_2 vibrate with the same frequency and in phase.
 - D Points equidistant from N_2 vibrate with the same frequency and the same amplitude.
- 21 A particle of charge $-5.0 \mu\text{C}$ is projected from X towards Y with kinetic energy $250 \mu\text{J}$. When the particle is at Y which is 0.25 m away from X, its kinetic energy decreased to $150 \mu\text{J}$.



Which of the following statements may not be correct?


- A X is at a higher electric potential than Y.
- B The potential difference between X and Y is 20 V.
- C The electric field between X and Y is of uniform strength 80 N m^{-1} .
- D The electric field between X and Y is directed towards Y.

- 22** The diagram shows an electrical circuit consisting of a 1.5 V cell of negligible internal resistance and two resistors P and Q.



Which statement about this circuit is correct?

- A** The cell converts 1.5 J of electrical energy to chemical energy for each coulomb of charge passing through it.
 - B** The energy dissipated per unit charge passing through P and Q is the same.
 - C** The potential difference across P and the potential difference across Q add up to 1.5 V.
 - D** The rate of flow of charge at point X is greater than the rate of flow of charge at Y.
- 23** In the circuit shown, the cells have negligible internal resistance and the reading on the ammeter is zero.




What is the resistance of R ?

- A** $2.0\ \Omega$
- B** $6.0\ \Omega$
- C** $12\ \Omega$
- D** $18\ \Omega$

- 24** Electrons, each of mass m and charge q , are accelerated from rest in a vacuum through a potential difference V .


The accelerated electrons are then projected at point S into a region of uniform magnetic field of flux density B , as shown. The electrons move in a circular path of radius r .



Which of the following expressions represents the specific charge $\frac{q}{m}$ of the electrons?

- A** $\frac{V}{2B^2r}$ **B** $\frac{2V}{B^2r}$ **C** $\frac{V}{2B^2r^2}$ **D** $\frac{2V}{B^2r^2}$


- 25** A rectangular coil made of 100 turns of wire with cross sectional area 30 cm^2 is placed within a uniform magnetic field of 0.80 T . The coil is rotated with an angular velocity of 100 rad s^{-1} . At different stages of its rotation, the cross sectional area of the coil can be parallel or perpendicular to the magnetic field, as shown.



What is the maximum e.m.f. induced and the corresponding orientation of the coil?

	maximum e.m.f. / V	orientation of coil
A	0.24	parallel to magnetic field
B	0.24	perpendicular to magnetic field
C	24	parallel to magnetic field
D	24	perpendicular to magnetic field


26 The graph below shows how the current in a coil varies with time.



What is the value of a steady current that will dissipate heat in the coil at the same average rate as the current above?

- A** 1.5 A **B** 2.3 A **C** 2.5 A **D** 2.6 A


27 An ideal transformer steps down the 240 V sinusoidal voltage from the mains to 12 V, which is then applied to a variable load.



What is the change in the current supplied by the mains when the resistance of the load is increased from $20\ \Omega$ to $50\ \Omega$?

- A** Decrease from 30 mA to 12 mA.
B Increase from 12 mA to 30 mA.
C Decrease from 12 mA to 4.8 mA.
D Increase from 0.24 A to 0.60 A.

28 The diagram shows two spectra of X-rays from an X-ray tube.



Which of the following statements is true?

- A The accelerating potential to produce spectrum B is lower than that to produce spectrum A.
- B A different target metal is used to produce spectra A and B as shown from the existence of additional peaks in spectrum A.
- C With the same accelerating potential, the temperature of cathode used to produce spectrum A is higher than that to produce spectrum B as shown from the higher intensity of X-ray photons in spectrum A.
- D The temperature of cathode used to produce spectrum A is higher than that to produce spectrum B as shown from the existence of additional peaks in spectrum A.

29 Which of the following radioactive samples has the greatest activity?

	nuclide	amount / mole	half-life / day
A	$^{225}_{89}\text{Ac}$	0.003	10
B	$^{228}_{90}\text{Th}$	0.1	400
C	$^{228}_{88}\text{Ra}$	0.6	2100
D	$^{241}_{94}\text{Pu}$	1.0	4800

- 30** A radioactive source is placed 1 cm from a Geiger-Muller tube, and various absorbers are placed between them, one at a time. For each absorber, a one-minute count is taken of the total number of decays, and this is repeated several times. The table shows the results of the experiment.

absorber	average number of decays detected in one minute
none	1043
0.1 mm paper	1040
1 mm aluminium	497
1 cm lead	6

The average background count per minute is 5.

What nuclear radiation do the results suggest the source was emitting?

- A** alpha and beta only
- B** beta only
- C** beta and gamma only
- D** alpha, beta and gamma

End of Paper