

XINMIN SECONDARY SCHOOL

新民中学 SEKOLAH MENENGAH XINMIN

Preliminary Examinations 2022

CANDIDATE NAME

CLASS

INDEX NUMBER

4049/01

ADDITIONAL MATHEMATICS

Paper 1

29 August 2022 2 hour 15 minutes

Secondary 4 Express/ 5 Normal Academic Setter : Mr Johnson Chua Vetter : Mrs Loh Si Lan Moderator: Ms Pang Hui Chin

Candidates answer on the Question Paper. No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, register number and class in the spaces at the top of this page. Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question. The use of an approved scientific calculator is expected, where appropriate. You are reminded of the need for clear presentation in your answers.

At the end of the paper, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is **90.**

Errors	Qn No.	Errors	Qn No.
Accuracy		Simplification	
Brackets		Units	
Geometry		Marks Awarded	
Presentation		Marks Penalised	

For Examiner's Use		
90		

Parent's/Guardian's Signature:

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the quadratic equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Expansion

$$(a+b)^n = a^n + {n \choose 1} a^{n-1}b + {n \choose 2} a^{n-2}b^2 + \dots + {n \choose r} a^{n-r}b^r + \dots + b^n$$

where *n* is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)...(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$
$$\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$
$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^{2} = b^{2} + c^{2} - 2bc \cos A$$
$$\Delta = \frac{1}{2}bc \sin A$$

1 Solve the equation $\sqrt{2x^2 - 5x + 3} - 2x = 0$.

2 (a) Express $3x^2 - 24x + 53$ in the form $a(x+b)^2 + c$ where a, b and c are constants. [2]

(b) Hence explain why there is no intersection between the graph $y = 3x^2 - 24x + 53$ [1] and the line y = 4.

[3]

3 Given that $f'(x) = \frac{6}{3x+5}$ and that f(x) passes through the origin, find the equation of f(x).

[3]

Turn over for Question 4



The diagram shows part of the graph $y = (2x-3)^2 \sqrt{x-1}$. Find the range of values of x for which y is increasing. [5]

5 (a) Solve
$$3^{2x-1} = 4^{2-x}$$
 and show that $x = \frac{\lg 48}{2\lg 6}$. [3]

(b) In order to obtain a graphical solution of the equation $x = \ln\left(\frac{7-x}{5}\right)^2$, a suitable straight line can be drawn on the same axes as the graph $y = 5e^{\frac{x}{2}} - 4$. Find the equation of this line. [2]

6 Solve the equation $3\sin A = \csc A - 5\cot A$ for $-180^\circ < A < 180^\circ$. [5]

7 Express $\frac{3x^3 + x^2 + 13x + 8}{x(x^2 + 4)}$ in partial fractions.

[6]

8 The diagram below shows the base of a hexagonal pyramid. The base is a regular hexagon, made up of six equilateral triangles of sides $(1+\sqrt{3})$ cm.



(a) Show that the area of the hexagon is $(6\sqrt{3}+9)$ cm². [3]

(b) Given that the volume of the hexagonal pyramid is $(9+4\sqrt{3})$ cm³, find the exact height of the hexagonal pyramid. (Volume of pyramid = $\frac{1}{3}$ × base area × height) [3]

- The function $f(x) = 2x^3 + ax^2 + bx + 2$, where *a* and *b* are constants, is exactly 9 divisible by (2x-1). Given that f(x) leaves a remainder of 9 when divided by (x+1), [4]
 - find the value of *a* and *b*. **(a)**

Using the value of a and b found in (**a**), solve the equation **(b)** $2x^3 + ax^2 + bx + 2 = 0.$

[4]

10 A gardener wants to use 40 m of fence to form a flower bed, which is made up of a rectangle of length 2x m and two identical semicircles of radius r m.



(a) Express r in terms of x.

[2]

[2]

(b) Show that the total area, $A m^2$, of the flowerbed is given by $A=\frac{400-4x^2}{\pi}\,.$

(c) Given that x can vary, find the value of x which gives a stationary value of A. [2]

(d) The gardener's wife claimed that he can optimise the length of the fencing to obtain a maximum area by forming only a circular flowerbed of radius $\frac{20}{\pi}$ m. Do you agree with her? Explain your answer with relevant workings. [2]

- 11 The negative x –axis and positive y –axis are tangents to circle C. Given that the radius of the circle is a units and that centre of circle is (-a, a),
 - (a) write down the equation(s) of the tangent(s) to the circle. [2]

- P(-1,8) and Q(-8,25) lie on the circumference of circle C.
- (b) Find the equation of the perpendicular bisector of *PQ*.

[3]

(c) Show that the equation of the circle is $x^2 + 26x + y^2 - 26y + 169 = 0$. [3]

(d) Explain why R(-14, 27) lies outside the circle.

[2]

12 (a) Find $\frac{d}{dx}(x\sin 2x)$.

(b) Hence, find $\int 2x \cos 2x \, dx$.

[1]

[1]

(c) The diagram below shows the curve $y = x^2 \sin 2x$, which passes through the origin and *A*. Differentiate $x^2 \cos 2x$ with respect to *x* and using the result found in part (b), find the area of the shaded region, which is bounded by the curve $y = x^2 \sin 2x$ and the *x*-axis.



13 A motorcycle is rode along a straight horizontal road. As it passes a point A, the brakes are applied and the motorcycle slows down, with its velocity halved as it passes B.For the journey from A to B, the displacement, s m, of the motorcycle from A, t seconds

after passing A, is given by $s = 400 \left(1 - e^{-\frac{t}{10}} \right).$

(a) Find the exact time taken for the journey from A to B.

[4]

(b) Find the average speed of the motorcycle for the journey from A to B. [2]

(c) Find the exact acceleration of the motorcycle at B.

[2]

14 (a) Show that
$$\frac{\cos(A+B) - \cos(A-B)}{\sin(A+B) - \sin(A-B)} = -\tan A$$
. [3]

(b) Given that A and B are both obtuse, and $cos(A+B) = -\frac{7}{25}$, find the value of sin(A+B). [2]

(c)	(i)	Given that $sin(A-B) = \frac{8}{17}$ and $cot(A-B) = \frac{15}{8}$, state the value of	F11
		$\cos(A-B)$.	[1]

(ii) Explain why *A* is larger than *B*.

(d) Using the identity from part (a), show that $\tan A = -\frac{13}{16}$. [2]

[1]

(e) Hence, find the exact value of $\tan B$.

--- End of Paper---