

CATHOLIC JUNIOR COLLEGE JC2 PRELIMINARY EXAMINATIONS Higher 2

CANDIDATE NAME		
CLASS	2T]

CHEMISTRY

Paper 2 Structured Questions

9647/02 Monday 1 September 2014 2 hours

Candidates answer on the Question Paper Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name and class in the spaces provided above.

Write in dark blue or black pen in the spaces provided, on the Question Paper. [PILOT FRIXION ERASABLE PENS ARE NOT ALLOWED]

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

You are reminded of the need for good English and clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part of the question.

	For Examiner's Use	
Paper 1		40
	Q 1	12
	Q 2	15
Paper 2	Q 3	15
	Q 4	15
	Q 5	15 72
	Q 1	20
	Q 2	20
Paper 3	Q 3	20
	Q 4	20
	Q 5	20 80
Total		192

This document consists of **19** printed pages and **1** blank page.

Planning (P)

1 **FA1** is a solution of $1.00 \text{ mol } \text{dm}^{-3}$ sodium hydroxide, NaOH.

FA2 is a solution of 1.00 mol dm⁻³ acid which is either a *monobasic* with the formula HA, or *dibasic* with the formula H_2A .

You are required to determine the *basicity* of an acid (either *monobasic* or *dibasic*), based on the information given below and the planning of the procedure you have to undertake.

Three separate experiments were carried out. Varying volumes of *FA1* were measured and then added to different volumes of *FA2* and the rise in temperatures, as a result of the mixing were recorded. The relevant measurements and results are tabulated below.

	Experiment 1	Experiment 2	Experiment 3
Volume of <i>FA1</i> / cm ³	25	25	50
Volume of <i>FA2</i> / cm ³	25	50	25
Temperature rise, ΔT / °C	6.9	4.6	9.1

(a) (i) Write a balanced equation for the reaction between *FA1* and *FA2* if the acid is *monobasic*. Do not omit spectator ions.

.....

(ii) Write a balanced equation for the reaction between *FA1* and *FA2* if the acid is *dibasic*. Do not omit spectator ions.

.....

[2]

(b) You are provided with the following apparatus and materials:

1 thermometer 1 styrofoam cup Two 50 cm³ measuring cylinders Sufficient *FA1* solution Sufficient *FA2* solution

Plan a simple procedure by describing the steps you would carry out for *Experiment* 1 given above so as to make the relevant measurements and recordings. You are to use *only* the apparatus and materials listed above.

[4]

For Examiner's Use

(c) Use the results of the relevant experiments given in the table above, explain clearly how the *basicity* of the acid is determined. Hence, state the *basicity* of the acid.

(d) (i) Hence, calculate the amount of heat evolved in *Experiment 1*. In your calculations, you are to assume that the specific heat capacity of the reaction mixture is $4.18 \text{ J g}^{-1} \text{ K}^{-1}$ and the density of the reaction mixture is 1 g cm⁻³.

	(ii) Using your answers to (d)(i) and the relevant equation in (a), calculate the enthalpy change of neutralisation, ΔH _{neu} .	For Examiner's Use
(e)	[2] From your calculated ΔH_{neu} value, state whether the acid is strong or weak.	
	[1]	
(f)	Identify one source of error in your procedure and explain how you would minimise the error.	
	[1]	

[Total: 12]

2 Scuba diving is a form of underwater diving, in which a diver uses a self-contained underwater breathing apparatus (scuba) to breathe underwater. Scuba diving is an exhilarating sport, and thanks in part to the gas laws, it is also a safe activity for trained individuals who are in good health.

A typical dive might be 12 to 19 m, but dives to 27 m are not uncommon. As seawater has a slightly higher density than fresh water, the pressure exerted by a column of 10 m of seawater is equivalent to a pressure of 1 atm. Pressure increases with increasing depth, so at a depth of 20 m, the pressure of the water will be 2 atm, and so on. The **total** pressure exerted on the diver is the **sum** of the atmospheric pressure and the pressure due to the column of water.

- (a) Consider a diver rising to the surface of the water from a depth of 6 m without breathing.
 - (i) Calculate the total pressure exerted on a diver at 6 m.

(ii) Show that, when the diver rises to the surface of the water, the volume of air trapped in the diver's lungs would be increased by 1.6 times.

The most important rule in scuba diving is to breathe continuously, especially when ascending, and to avoid holding breath underwater. This is to avoid barotrauma where physical damage to body tissues is caused by a difference in pressure between a gas space inside, or in contact with the body, and the surrounding waters. The sudden expansion of air can fatally rupture the membranes of the lungs. This is different in free diving, which relies on the diver's ability to hold his or her breath until resurfacing, rather than on the use of a breathing apparatus.

(iii) By considering the volume of air in the diver's lungs, explain why barotrauma is not experienced for a free diver ascending at a depth of 6 m.

[5] [Turn over For Examiner's

Use

A scuba set is a breathing set that is carried entirely by an underwater diver and provides the diver with breathing gas at the ambient pressure. A breathing gas is usually a mixture of oxygen and nitrogen used for respiration. For instance, air is the most common and only natural breathing gas. Our bodies function best when oxygen gas has a partial pressure of 0.20 atm, as in the air we breathe. When a diver is submerged, a special valve automatically adjusts the pressure of the air breathed from the scuba tank to ensure that the air pressure equals the water pressure at all times.

6

Although nitrogen gas may seem to be the obvious choice to mix with oxygen gas for use in the scuba tank, there is a serious problem associated with it. When the partial pressure of nitrogen gas exceeds 1 atm, enough of the gas dissolves in the blood to cause *nitrogen narcosis*. The effects on the diver resemble those associated with nitrous oxide ("laughing gas") inhalation. For this reason, helium is often used to dilute oxygen gas.

(b) (i) Show that the oxygen present in air has a partial pressure of 0.20 atm.

(ii) Consider a diver submerged at a depth of 6 m.
 Using your answer in (a)(i), calculate the mole fraction of oxygen that is needed to maintain the partial pressure of 0.20 atm.

7

(iii) Suggest a reason why helium can be used in place of nitrogen gas.

.....

[4]

For

Examiner's Use

Nitrous oxide (N_2O) commonly known as laughing gas, is a colourless and non-flammable gas, with a slightly sweet odour and taste. It is used in surgery and dentistry for its anaesthetic and analgesic effects. It is known as "laughing gas" due to the euphoric effects of inhaling it, a property that has led to its recreational use as a dissociative anaesthetic.

(c) (i) Draw the dot-and-cross diagram of N_2O , given that N is the central atom, and state its shape.

Nitrous oxide decomposes into oxygen and nitrogen according to the following equilibrium:

 $2 N_2O(g) \implies O_2(g) + 2 N_2(g)$

At 25 °C, K_c is 7.30 x 10^{34} mol dm⁻³.

(ii) Suggest how N₂O can be stored to prevent decomposition.

.....

(iii) Suggest the significance of the value of K_c in the reaction.

.....

(iv) The value of $K_{\rm p}$ is related to $K_{\rm c}$ according to the following formula:

$$K_p = K_c(RT)^{\Delta n}$$

where $\Delta n = no$. of moles of gaseous products – no. of moles of gaseous reactants

Using the given formula, calculate the value of K_p for the reaction at 25 °C, indicating clearly its units.

[6]

For Examiner's Use

[Total: 15]

For Examiner's Use

[2]

3 Nickel is a silvery-white metal and shows significant chemical activity with oxygen in powdered form but larger pieces of the metal are slow to react with air at ambient conditions due to the formation of a protective oxide layer.

Cadmium is a soft, bluish-white metal chemically similar to zinc and also displays properties similar to heavy metals. Cadmium occurs as a minor component in zinc ores and is a by-product of zinc manufacture.

Both nickel and cadmium are used in applications such as batteries and electroplating for corrosion resistance.

	nickel	cadmium
melting point / °C	1455	321
density / g cm ⁻³	8.91	8.65

(a) Complete the electronic configuration of nickel(II) and cadmium(II) ions.

₂₈ Ni ²⁺	$1s^2 2s^2 2p^6$
48Cd ²⁺	1s ² 2s ² 2p ⁶

(b) Briefly account for the difference in melting points between nickel and cadmium.

[2]

- (c) The nickel–cadmium battery (NiCad battery) is a type of rechargeable battery using nickel oxide hydroxide and metallic cadmium as electrodes.
 - (i) During electrical discharge, the overall chemical reaction is as shown:

 $2NiO(OH) + Cd + 2H_2O \rightarrow 2Ni(OH)_2 + Cd(OH)_2$

Cadmium reacts with an alkaline electrolyte at one of the electrodes. Write the equations for reactions that take place at the anode and the cathode.

Anode:

[Turn over

(ii) If the cell produces 1.8 A of current, and if the mass of nickel oxide hydroxide contained in a typical NiCad battery is 9.15 g, how long can the battery operate before it stops discharging electrical power?

For Examiner's Use

(iii) List one advantage and one disadvantage of the NiCad battery over the conventional alkaline dry cell battery.

(d) Aqueous nickel(II) chloride, NiC l_2 , is a green solution which undergoes the following reactions:



(i) State the formula of compound **B** given that the coordination number of the cation in **B** is the same as that found in the cation present in NiC $l_2(aq)$.

- For Examiner's Use
- (ii) The following diagram shows how the d-orbitals of a metal ion are split in a tetrahedral and square planar complex.



Both **C** and **D** contain complexes with cations having coordination number of 4. When Cl^{-} ligands in **D** is changed to CN^{-} ligands in **C**, the Ni²⁺ ion changes its geometry and electronic configuration from a 'high spin' state with lower energy gap to a 'low spin' state with higher energy gap.

In a 'high spin' state, the electrons occupy all the d-orbitals singly, before starting to pair up in the lower energy d-orbitals.

In a 'low spin' state, the lower energy d-orbitals are filled first, by pairing up if necessary, before the higher energy d-orbitals are used.

Complete the diagrams showing the electrons distribution of Ni²⁺ ion in **C** and in **D** and hence predict the number of unpaired electrons in each complex.



[Turn over



Number of unpaired electrons:

[5]

[Total: 15]

- 4 Amino acids are critical to life and they serve as the building blocks of proteins.
 - (a) Aspartic acid plays a role in the neuroendocrine system, as a regulator in the synthesis and release of hormones. It also plays an important role in the conversion of carbohydrates into energy. Proper levels of aspartic acid are necessary for the synthesis of other biochemicals.



aspartic acid

The three pK_a values associated with aspartic acid are 2.10, 3.86 and 9.82. The isoelectric point is 2.77.

(i) In the boxes provided, assign the pK_a values for each carboxylic acid group. Suggest, with explanations, the pK_a values associated with these groups.



 For Examiner's Use (ii) Sketch the pH-volume added curve you would expect to obtain when 40 cm³ of 0.10 mol dm⁻³ NaOH is added to 10 cm³ of 0.10 mol dm⁻³ of the **protonated** form of aspartic acid. Show clearly on your curve where the three pK_a values occur and the isoelectric point.



(iii) Write an equation to show how aspartic acid can act as a buffer when a small amount of OH⁻ is added at a pH of 3.86.

(iv) Calculate the pH of the resulting solution when 40 cm³ of NaOH has been added.

For

Examiner's Use (b) Phenylalanine, together with aspartic acid, is also used to make aspartame, an artificial sweetener. The main synthesis steps are shown below.



(c) Hair is made mostly of a protein called keratin, which is also present in nails. The table below shows the R groups of some of the α -amino acids which make up keratin.

Acid	R group
Lys	$-CH_2CH_2CH_2CH_2NH_2$
Ser	-CH ₂ OH
Glu	-CH ₂ CH ₂ CO ₂ H
Cys	-CH ₂ SH
Met	$-CH_2CH_2SCH_3$

15

For Examiner's

Use

Hair can be curly because of the covalent disulfide bonds that can be formed between the amino acid residues. This is brought about by an oxidation process.

(i) Write a chemical equation to show how the cysteine amino acid residues in the keratin molecule can form disulfide bridges.

In order to straighten curly hair permanently, using a heating iron is insufficient. Instead, one will need to go to the hairdresser for rebonding, a chemical hair treatment that makes his or her hair straight.
Ammonium thioglycolate (HSCH₂CO₂NH₄) is a chemical which is used to straighten hair. It can reverse the reaction in (i) and break the disulfide bridges in hair.
Keratin-S-S-keratin + 2 HS-CH₂CO₂NH₄ → 2 HS-keratin + HO₂C-CH₂S-SCH₂CO₂H + 2 NH₃
(ii) Suggest the role of ammonium thioglycolate in the above reaction.
(iii) What aspect of keratin's protein structure is altered during the process of rebonding?

(d) Upon acidic hydrolysis of keratin, a tripeptide, lys-ser-glu is obtained. Draw the structure of the tripeptide formed.

[Total: 15]

For

Examiner's Use **5** Spices of the ginger family such as ginger and turmeric have a long history of culinary and medical use.

17

The following three compounds present in either ginger or turmeric are medically important:



tumerone

The group $-OCH_3$ which is present in zingerone and gingerol can be regarded as inert.

(a) In the boxes below, name one of the three compounds above that will react with each of the following reagents. In each case, give the observations for the reaction undergone.
 Each compound may only be used once.

Reagent	Name of compound	Observations
Neutral FeCl ₃ (aq)		
2, 4-DNPH		
PCl ₅		

[6]

(b) Complete the following diagram with the reagents and conditions required for Steps I-IV, and draw the structure of the product formed from Step III in the box provided.

For Examiner's Use



- 19
- (c) Compound X exhibits two types of stereoisomerism.
 - (i) Circle all chiral centres in the molecule of **Compound X** given in (b).
 - (ii) While there are no C=C bonds in **Compound X**, it exhibits cis-trans isomerism. Explain why this is the case.

.....

.....

(iii) The total number of stereoisomers for a compound is given by 2ⁿ, where n = number of chiral centres + number of pairs of cis-trans isomers

Suggest why, when determining the total number of stereoisomers for **Compound** X, the above statement is not valid.

[4]

[Total: 15]

For Examiner's Use

BLANK PAGE