DARRELL ER (COPYRIGHTED) ©

DARRELL ER (COPYRIGHTED) ©

TOPIC 11.3: ALKENES

DARRELL ER (COPYRIGHTED) ©

THE ABOUT

CHAPTER ANALYSIS

MASTERY

- Important topic
- Take note of alkene's various chemical reactions

EXAM

- Alkenes are always tested
- Understand the difference between 'saturated' & 'unsaturated' compounds

WEIGHTAGE

- Heavy overall weightage
- Entire Organic Chemistry portion accounts for 15-20% of each year's Chemistry paper

DARRELL ER (COPYRIGHTED) ©

KEY CONCEPT

ALKENES
HOMOLOGOUS SERIES
FUNCTIONAL GROUP
GENERAL FORMULA

DARRELL ER (COPYRIGHTED) ©

Name	Carbon atoms	Molecular Formula	Full Structural Formula	Condensed structural formula
Ethene	2	C_2H_4		CH_2CH_2
Propene	3	C_3H_6		CH_2CHCH_3
Butene	4	C_4H_8		$CH_3CH_2CHCH_2$

Alkenes

Alkenes are unsaturated hydrocarbons with a **general formula of C_nH_{2n}.**

As the carbon atoms are **not bonded to the maximum of four atoms**, alkenes are considered to be **'unsaturated'**.

Unsaturated hydrocarbons are hydrocarbons that contain one or more C=C double bond. (For eg: Vegetable oil)

Functional group

Alkenes contain C=C double covalent **bonds.**

DARRELL ER (COPYRIGHTED) ©

KEY CONCEPT

ALKENES
PHYSICAL PROPERTIES
CHEMICAL PROPERTIES
CRACKING



DARRELL ER (COPYRIGHTED) ©

PHYSICAL PROPERTIES

Physical property	Reasoning
Melting and boiling points	**As the number of carbon atoms in the alkene increases**, the **melting and boiling points of alkenes increases as well.** As the **number of carbon atoms in an alkene increases**, **the size of the molecules are bigger and have stronger intermolecular forces of attraction between each other.** As such, more heat energy is needed to overcome the intermolecular forces of attraction between the alkene molecules. Hence, larger alkenes containing more carbon atoms will have higher melting and boiling points.
Volatility	**As the number of carbon atoms in the alkene increases**, the **volatility of alkenes decreases.** (similar to m.p. & b.p.) **With a higher relative molecular mass, there would be stronger intermolecular forces of attraction between the alkene molecules.** As such, more energy is needed to overcome the intermolecular forces of attraction between the alkene molecules. Hence, larger alkene molecules are less likely to evaporate in room temperature.
Density	**As the number of carbon atoms in the alkene increases**, the **density of alkenes increases**.
Viscosity	**As the number of carbon atoms in the alkene increases**, the **viscosity of alkenes decreases**. **(more difficult to flow)** Alkenes with longer hydrocarbon chains flow less easily as they tend to get stuck together.
Flammability	**As the number of carbon atoms in the alkene increases**, the **flammability of alkenes decreases. (more difficult to burn)** The larger alkenes contain a **higher percentage mass of carbon atoms** and would undergo incomplete combustion which results in a smokier flame.
Solubility	All alkenes are **insoluble in water but are soluble in organic solvents** like ethanol.

DARRELL ER (COPYRIGHTED) ©

CHEMICAL REACTIONS

CHEMICAL REACTIONS OF ALKENES

1) **Combustion**
1) **Hydrogenation (add hydrogen gas)**
1) **Bromination (add aqueous bromine)**
1) **Hydration (add water)**

} Addition reactions

1) **Polymerisation (combined alkenes to form long chain)**

1) COMBUSTION

In the presence of excess oxygen, an alkene would undergo **complete combustion**, producing carbon dioxide and water.

$$2\ C_3H_6 + 9\ O_2 \rightarrow 6\ CO_2 + 6\ H_2O$$

If there is insufficient oxygen for complete combustion, the alkene would undergo **incomplete combustion**, producing water and carbon monoxide. (similar to alkanes)

$$C_3H_6 + 3\ O_2 \rightarrow 3\ CO + 3\ H_2O$$

2) HYDROGENATION (addition of hydrogen)



Condition: 200°C and nickel catalyst.

Through the process of hydrogenation, alkenes are converted to alkanes. (E.g. vegetable oils can be converted to margarine.)

Margarine is considered a **saturated compound** as it undergoes hydrogenation process and contains no C=C bonds.

DARRELL ER (COPYRIGHTED) ©

CHEMICAL REACTIONS

CHEMICAL REACTIONS OF ALKENES

1) **Combustion**
1) **Hydrogenation (add hydrogen gas)**
1) **Bromination (add aqueous bromine)**
1) **Hydration (add water)**

} Addition reactions

1) **Polymerisation (combined alkenes to form long chain)**

3) BROMINATION (addition of aqueous bromine)

Conditions: Aqueous bromine, room temperature

When **aqueous bromine reacts with alkenes occurs**, the **reddish-brown solution decolourises to become colourless.**

Aqueous bromine is used as a **test to distinguish alkanes and alkenes** as only alkenes would react with aqueous bromine in the absence of ultraviolet light.

$$C_2H_4\ (g) + Br_2\ (aq) \rightarrow C_2H_4Br_2\ (l)$$

4) HYDRATION (addition of steam)

Addition of steam under the right conditions can cause a reaction with the C=C double bond to produce an alcohol containing the **−OH functional group.**



Conditions:

Temperature of **300°C** and at a pressure of **60 atm**, **Phosphoric(V) acid** as a catalyst.

DARRELL ER (COPYRIGHTED) ©

CHEMICAL REACTIONS

CHEMICAL REACTIONS OF ALKENES

1) **Combustion**
1) **Hydrogenation (add hydrogen gas)**
1) **Bromination (add aqueous bromine)**
1) **Hydration (add water)**

Addition reactions

1) **Polymerisation (combined alkenes to form long chain)**

5) POLYMERISATION

Polymers are formed when **multiple identical alkene molecules (monomers) are joined together to form a large molecule**.

For example, ethene molecules can be joined together through polymerisation to form the polymer: poly(ethene).



DARRELL ER (COPYRIGHTED) ©

CRACKING

Catalytic cracking

Catalytic cracking is a process where a long-chain hydrocarbons from petroleum are broken down into shorter-chain hydrocarbons in a presence of a catalyst.

The process is used for **producing fuels for vehicles, production of alkenes & production of hydrogen.** Smaller chain alkanes and alkenes are more useful than longer chain alkanes as they are in higher demand.

Conditions: **Aluminium oxide (Al_2O_3)** as catalyst and **silicon (IV) oxide (SiO_2)** at a temperature of **600°C**.



At least one of the products formed will be an alkene. Alkanes, hydrogen and even carbon can be produced from the cracking process as well.

Examples:

Long chain alkane → shorter chain alkane + alkene

$$C_{10}H_{22} \rightarrow C_5H_{12} + C_5H_{10}$$

Long chain alkane → shorter chain alkene + shorter chain alkene + hydrogen

$$C_{10}H_{22} \rightarrow C_6H_{12} + C_4H_8 + H_2$$

DARRELL ER (COPYRIGHTED) ©

DARRELL ER (COPYRIGHTED) ©

POLYMER

Condensation Polymerisation
(elimination of water)



LONG CHAIN ALKANE

H_2 gas
(For Haber process)

Catalytic Cracking
(Al_2O_3 & SiO_2, 600 °C)

Addition Polymerisation
(High temp & pressure)

SUGAR

$C_6H_{12}O_6 \rightarrow 2\ C_2H_5OH + 2\ CO_2$

Fermentation
(37°C, yeast & no O_2)

Oxidation
(acidified aqueous potassium manganate(VII) / exposed to air)

Hydration
(300 °C & 60 atm, Phosphoric(V) acid)

ALKANE
C - C

Hydrogenation
(200 °C & nickel)

# ALKENE
C = C

ALCOHOL
-OH

CARBOXYLIC ACID
-COOH

Substitution
(UV light)

Bromination
(Test for C=C bonds)

Esterification
(warm, sulfuric acid)

Prefix

Meth- 1
Eth- 2
Prop- 3
But- 4
Pent- 5
Hex- 6
Hep- 7
Oct- 8
Non- 9
Dec- 10

$CH_4\,(g) + Br_2\,(g) \rightarrow CH_3Br\,(g) + HBr\,(g)$

$C_2H_4\,(g) + Br_2\,(aq) \rightarrow C_2H_4Br_2\,(aq)$



ESTER
-COO-

+ H_2O

ALL ORGANIC COMPOUNDS
Complete Combustion

$X + O_2\,(g) \rightarrow CO_2\,(g) + 2H_2O\,(l)$

Incomplete Combustion

$X + O_2\,(g) \rightarrow CO\,(g) + 2H_2O\,(l)$

Try it yourself! (TYS Question)

28. What is the structure of the product of the reaction between butene, $CH_3–CH_2–CH=CH_2$, and bromine, Br_2? (N2016/P1/Q36)

A $CH_2Br–CH_2–CH_2–CH_2Br$
B $CH_2Br–CH_2–CHBr–CH_3$
C $CH_3–CHBr–CH_2–CH_2Br$
D $CH_3–CH_2–CHBr–CH_2Br$ ( )

Answer:

28. **D**
Bromine adds across the double bond in butene.

$CH_3—CH_2—CH=CH_2 + Br_2 \rightarrow$
$CH_3—CH_2—CHBr—CH_2Br$

Try it yourself! (TYS Question)

51. Which statements about the cracking of hydrocarbons are correct? (N2020/P1/Q36)

1 Cracking involves breaking down hydrocarbon molecules.
2 One of the products of cracking is always unsaturated.
3 Cracking is essential because of the demand for fractions containing smaller molecules.

A 1 and 2 only
B 2 and 3 only
C 1 and 3 only
D 1, 2 and 3

( )

Answer:

51. **D**
Statement 1: Cracking involves breaking down of large, saturated hydrocarbon molecules.
Statement 2: Cracking produces at least one unsaturated hydrocarbon.
Statement 3: One advantage of cracking is that the process produces smaller and more useful molecules which are in demand.

About Us

OVERMUGGED is a learning platform created by tutors, for students.

Our team of specialist tutors offer 1-to-1 private tuition, group tuitions and crash courses.

Follow us on IG and join our Telegram channel to get the latest updates on our free online revision sessions, webinars and giveaways!

If you would want to join Darrell's group tuition, contact him at:
Whatsapp: 8777 0921
Telegram: @DarrellEr
Website: https://www.overmugged.com/darrell

For more free notes & learning materials, visit: www.overmugged.com

OVERMUGGED
By Tutors, For Students

Notes prepared by:

Darrell Er

'O' Levels Chemistry & Physics

OVERMUGGED's Curated Notes

Found the free notes useful? We got something better!

OVERMUGGED's curated notes is a **highly condensed booklet** that **covers all content within the MOE syllabus**.

This booklet consist of **key concept breakdowns**, **worked examples** and **exam tips/ techniques** to required to ace your exams.

Get an **upgraded version** of the free notes and supercharge your revision!

Purchase here.

Crash courses

Check out our upcoming crash courses at:
https://www.overmugged.com/crashcourses

'O' levels subject available:
- Pure Chemistry
- Pure Physics
- Pure Biology
- Combined Science
- E-Math
-A-Math
-English
- History
- Geography
- Combined Humanities
- Principles of Accounts (POA)