

2023 JC1 H2 CHEMISTRY (9729)
CORE IDEA 2: STRUCTURE AND PROPERTIES
TOPIC 3: CHEMICAL BONDING

Name: _____

Civics Group: _____

Students should be able to:

- (a) show understanding that all chemical bonds are electrostatic in nature and describe:
 - (i) ionic bond as the electrostatic attraction between oppositely charged ions
 - (ii) covalent bond as the electrostatic attraction between a shared pair of electrons and positively charged nuclei
 - (iii) metallic bond as the electrostatic attraction between a lattice of positive ions and delocalised electrons
- (b) describe, including the use of 'dot-and-cross' diagrams,
 - (i) ionic bonding as in sodium chloride and magnesium oxide
 - (ii) covalent bonding as in hydrogen; oxygen; nitrogen; chlorine; hydrogen chloride; carbon dioxide; methane; ethene
 - (iii) co-ordinate (dative covalent) bonding, as in formation of the ammonium ion and in the Al_2Cl_6 molecule
- (c) describe covalent bonding in terms of orbital overlap (limited to s and p orbitals only), giving σ and π bonds (see also Section 11.1)
- (d) explain the shapes of, and bond angles in, molecules such as BF_3 (trigonal planar); CO_2 (linear); CH_4 (tetrahedral); NH_3 (trigonal pyramidal); H_2O (bent); SF_6 (octahedral) by using the Valence Shell Electron Pair Repulsion theory
- (e) predict the shapes of, and bond angles in, molecules analogous to those specified in (d)
- (f) explain and deduce bond polarity using the concept of electronegativity
[quantitative treatment of electronegativity is **not** required]
- (g) deduce the polarity of a molecule using bond polarity and its molecular shape (analogous to those specified in (d))
- (h) describe the following forces of attraction (electrostatic in nature):
 - (i) intermolecular forces, based on permanent and induced dipoles, as in $CHCl_3(l)$, $Br_2(l)$ and the liquid noble gases
 - (ii) hydrogen bonding, using ammonia and water as examples of molecules containing $-NH$ and $-OH$ groups
- (i) outline the importance of hydrogen bonding to the physical properties of substances, including ice and water
- (j) explain the terms *bond energy* and *bond length* for covalent bonds
- (k) compare the reactivities of covalent bonds in terms of bond energy, bond length and bond polarity
- (l) describe, in simple terms, the lattice structure of crystalline solid which is:
 - (i) ionic, as in sodium chloride and magnesium oxide
 - (ii) simple molecular, as in iodine
 - (iii) giant molecular, as in graphite and diamond
 - (iv) hydrogen-bonded, as in ice
 - (v) metallic, as in copper
 [the concept of the 'unit cell' is not required]
- (m) describe, interpret and/or predict the effect of different types of structure and bonding on the physical properties of substances
- (n) suggest the type of structure and bonding present in a substance from given information

Lecture Contents

1 Covalent Bonding	4
1.1 Covalent Bond Formation.....	4
1.2 The Overlap of Orbitals: Sigma Bonds and Pi Bonds	5
1.2.1 Sigma (σ) Bonds.....	5
1.2.2 Pi (π) Bonds.....	6
1.3 Types of Covalent Bonds	6
1.3.1 Multiple Bonding – Double and Triple Bond	6
1.3.2 Dative Bonding/Co-ordinate Bonding	7
1.4 Electronegativity.....	8
1.5 Properties of Covalent Bonds.....	9
1.5.1 Bond Polarity	9
1.5.2 Bond Strength.....	10
1.5.3 Bond Length	11
1.6 Representing Covalent Bonds: Dot-And-Cross Diagrams.....	13
1.6.1 Dot-And-Cross for Molecules	14
1.6.2 Dot-And-Cross Diagrams for Cations / Anions	15
1.6.3 Dot-And-Cross Diagrams for Electron Deficient (with Incomplete Octet) Molecules	16
1.6.4 Dot-And-Cross Diagrams for Species with Odd Number (Unpaired) Valence	Electrons.....
1.6.5 Dot-And-Cross Diagrams for Expansion of Octet Structures	17
1.6.6 Dot-And-Cross Diagrams Showing Dative Covalent Bonds.....	18
2 Shape and Polarity.....	20
2.1 The Valence Shell Electron Pair Repulsion (VSEPR) Theory	20
2.2 Polarity of Molecules	24
3 Intermolecular Forces of Attraction.....	27
3.1 Instantaneous Dipole–Induced Dipole Interactions.....	27
3.2 Permanent Dipole–Permanent Dipole Interactions	30
3.3 Hydrogen Bonding.....	31
3.4 Comparing the Three Types of Intermolecular Forces of Attraction	32
4 Covalent Substances and Their Physical Properties	36
4.1 Simple Molecules	36
4.2 Influence of Hydrogen Bonding on Physical Properties of Molecules	38
4.2.1 Unusually high boiling points of ammonia, water and hydrogen fluoride.....	38
4.2.2 Anomalous relative molecular masses of organic acids ($R-CO_2H$)	39
4.2.3 High solubility of ammonia, alcohols ($R-OH$), organic acids and amines ($R-NH_2$)	in water
4.2.4 Different physical properties of isomers	40
4.2.5 Higher density of water compared to ice	41
4.3 Giant Molecular Structures	42

5 Ionic Bonding	45
5.1 Formation of Ionic Bond	45
5.2 Factors affecting Ionic Bond Strength	46
5.3 Giant Ionic Lattice	48
5.4 Physical Properties of Ionic Compounds	49
6 Intermediate Bond Types	51
6.1 Ionic Character in Covalent Bond	51
6.2 Covalent Character in Ionic Bond	52
7 Metallic Bonding	54
7.1 Definition of Metallic Bond	54
7.2 Factors affecting Metallic Bond Strength	55
7.3 Physical Properties of Metals	56
Chemical Bonds Summary	57
Intermolecular Forces of Attraction	58
Shapes of Molecules	59
Appendix 1: A Little Bit of History	60
Appendix 2: Structure of the Atom	62
Appendix 3: Ionic Bonding – Prelude	63
Appendix 4: Covalent Bonding – Prelude	64
Appendix 5: Coordination Number	65

REFERENCES

- 1 Chemistry (for CIE AS & A Level) by Peter Cann & Peter Hughes
- 2 Cambridge International AS and A Level Chemistry Revision Guide by Judith Potter and Peter Cann
- 3 Chemical Bonding and Molecular Geometry by Ronald Gillespie & Paul Popelier

1 Covalent Bonding


1.1 Covalent Bond Formation

- LO (a) show understanding that all chemical bonds are electrostatic in nature and describe:
 (ii) covalent bond as the electrostatic attraction between shared electrons and positively charged nuclei

Covalent bond is the **electrostatic forces of attraction** between **shared electrons** and the **positively charged nuclei**.


A covalent bond is formed when atoms (*usually non-metals*) **share valence electrons** through **overlap of orbitals**, leading to increased electron density between the two nuclei where the shared electrons are most likely to be found.

Using a hydrogen molecule as an example:



- ① As the two hydrogen atoms approach each other from a distance, the electron of one atom will electrostatically repel the electron of the other atom. However, this will be compensated by the additional electrostatic force of attraction between this electron and the nucleus of the other hydrogen atom.
- ② As the hydrogen atoms move closer to each other, the two electrons will encounter an even greater attraction to the opposite nucleus but will also continue to repel each other.
- ③ At the minimum point, the **electrostatic force of attraction** between the two nuclei and shared pair of electrons is the strongest. **This defines the covalent bond length.**
- ④ If the hydrogen atoms continue to move closer to each other, the inter-electronic and inter-nuclear repulsion increase. However, the inter-nuclear repulsion is more significant.

At ③ when a covalent bond is formed, the average nucleus-nucleus distance is known as the **bond length**.




Being attracted to both the hydrogen nuclei, the two electrons spend most of their time in the region between the two nuclei. This is where the **highest electron density** (or electron probability) occurs in a covalent bond. The two electrons are therefore *shared* between the two adjacent atoms.

1.2 The Overlap of Orbitals: Sigma Bonds and Pi Bonds

LO (c) Describe covalent bonding in terms of orbital overlap (limited to s and p orbitals only), giving σ and π bonds.


1.2.1 Sigma (σ) Bonds

A single bond is formed when a pair of electrons is shared between two atoms. If this bond is formed due to the **head-on overlap of an orbital of each atom** (this can involve s orbitals, p orbitals, or a mixture of the two as shown in the diagram below), then this bond is referred to as a **sigma (σ) bond**.



1.2.2 Pi (π) Bonds

It is also possible to form a bond due to the **sideways overlap of a p orbital on each atom**. Such a bond is referred to as a **pi (π) bond**.





In a π bond, there is **less effective overlap of orbitals**. Hence, **a π bond is weaker than a σ bond**.

This sideways overlap of two p orbitals is always preceded by the head-on overlap of two other orbitals on the same atoms *i.e.* **π bonds can only form after a σ bond is formed**.



1.3 Types of Covalent Bonds

1.3.1 Multiple Bonding – Double and Triple Bond

Atoms can share more than one electron pair with their neighbours. Sharing two electron pairs produces a **double bond**, *e.g.* in the oxygen molecule, O_2 , and sharing three electron pairs produces a **triple bond**, *e.g.* in the nitrogen molecule, N_2 .



Carbon can also form multiple bonds, *e.g.* in ethene, C_2H_4 , and ethyne, C_2H_2 .




A **double bond** consists of a **σ bond and a π bond** between the same two atoms.

A **triple bond** consists of a **σ bond and two π bonds** between the same two atoms.

A single bond corresponds to	1 σ bond
A double bond corresponds to	1 σ bond and 1 π bond
A triple bond corresponds to	1 σ bond and 2 π bonds

Example 1A

State the number of σ bonds and π bonds in propene, $CH_3CH=CH_2$.



No. of single bonds = 7 ; No. of double bonds = 1
 No. of σ bonds = 7 + 1 = 8 ; No. of π bonds = 1

1.3.2 Dative Bonding/Co-ordinate Bonding


Usually, a covalent bond consists of each atom contributing one electron for sharing. It is also possible for just one of the atoms involved to provide **both** bonding electrons to form a covalent bond.

The **donor atom** provides two electrons (from a **lone pair of electrons**) to an **acceptor atom**. The acceptor must contain an **empty valence orbital**. This kind of bonding is called **dative bonding** or **co-ordinate bonding**. Dative bonds are represented by arrows “→” showing which way the lone pair is donated.

The dative bond, once formed, is not different from any other covalent bond i.e. has the same bond strength as a normal covalent bond.

Example 1


When gaseous ammonia and gaseous boron trifluoride react with each other, a white solid is formed, with the formula NH_3BF_3 . The nitrogen atom in ammonia has a lone pair, and the boron atom in boron trifluoride has an empty 2p orbital.



Example 2

When gaseous ammonia and gaseous hydrogen chloride react, the white solid ammonium chloride is formed. The lone pair of electrons on the nitrogen atom of ammonia has formed a dative bond with the hydrogen atom of the hydrogen chloride molecule.

At the same time, the $\text{H}-\text{Cl}$ bond breaks and the electrons in the $\text{H}-\text{Cl}$ bond form a fourth lone pair on the chlorine atom, giving rise to a chloride ion.



All four N–H bonds are exactly the same. It is not possible to tell which one is a dative bond.

Example 3

Carbon monoxide has an interesting structure. One of the bonds in the triple bond between carbon and oxygen is a dative bond.



1.4 Electronegativity

LO (f) Explain and deduce bond polarity using the concept of electronegativity [quantitative treatment of electronegativity is not required.]

Electronegativity refers to the ability of an atom to **attract the shared pair(s) of electrons** in a **covalent bond**.

Electronegativity depends on the nuclear charge and the distance of the outer electrons from the nucleus.

In 1960, Linus Pauling developed a relative index of the ability of an atom to attract electrons, referenced to an **arbitrary** value of 2.1 for hydrogen. Numerical values of electronegativity on this scale range from fluorine, with a value of 4.0, to an alkali metal such as caesium, with a value of 0.7. The larger the number, the more electronegative (electron attracting) the atom is.

1																	18
H 2.1																	He –
2												13	14	15	16	17	
Li 1.0	Be 1.5											B 2.0	C 2.5	N 3.0	O 3.5	F 4.0	Ne –
Na 0.9	Mg 1.2	3	4	5	6	7	8	9	10	11	12	Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0	Ar –
K 0.8	Ca 1.0	Sc 1.3	Ti 1.5	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Co 1.8	Ni 1.8	Cu 1.9	Zn 1.6	Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8	Kr 3.0
Rb 0.8	Sr 1.0	Y 1.2	Zr 1.4	Nb 1.6	Mo 1.8	Tc 1.9	Ru 2.2	Rh 2.2	Pd 2.2	Ag 1.9	Cd 1.7	In 1.7	Sn 1.8	Sb 1.9	Te 2.1	I 2.5	Xe 2.6
Cs 0.7	Ba 0.9	lantha- noids	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9	Tl 1.8	Pb 1.8	Bi 1.9	Po 2.0	At 2.2	Rn 2.4
Fr 0.7	Ra 0.9	acti- noids	Rf –	Db –	Sg –	Bh –	Hs –	Mt –	Ds –	Rg –	Cn –		Fl –		Lv –		

lanthanoids	La 1.1	Ce 1.1	Pr 1.1	Nd 1.1	Pm 1.1	Sm 1.2	Eu 1.1	Gd 1.2	Tb 1.1	Dy 1.2	Ho 1.2	Er 1.2	Tm 1.3	Yb 1.1	Lu 1.1
	Ac 1.1	Th 1.3	Pa 1.5	U 1.4	Np 1.4	Pu 1.3	Am 1.3	Cm 1.3	Bk 1.3	Cf 1.3	Es 1.3	Fm 1.3	Md 1.3	No 1.3	Lr –

In general, electronegativity:

- **Increases across the period**, e.g. F is more electronegative than O which is more electronegative than N
- **Decreases down the group**, e.g. Br is less electronegative than Cl

(Useful tip: Remember that **F is the most electronegative element!**)


1.5 Properties of Covalent Bonds

- LO (f) Explain and deduce bond polarity using the concept of electronegativity [quantitative treatment of electronegativity is **not** required.]
- (j) Explain the terms bond energy and bond length for covalent bonds.
- (k) Compare the reactivities of covalent bonds in terms of bond energy, bond length and bond polarity.


1.5.1 Bond Polarity

A **polar covalent bond** is formed if the two atoms involved have **different electronegativities**, resulting in bonding electrons being **unequally shared** between the two atoms. The difference in electronegativities cause a **permanent separation of partial charges (dipoles)** between the two atoms.

For example, in hydrogen chloride, HCl , the more electronegative chlorine atom draws electron density towards itself, resulting in the formation of dipoles.



- $\delta+$: partial positive charge
(i.e. less electronegative / more electropositive)
- $\delta-$: partial negative charge
(i.e. more electronegative / less electropositive)



We sometimes represent the polar HCl molecule in this way:

The 'cross-and-arrow' symbol ($\text{+} \longrightarrow$) shown above is used to show the direction in which the electrons are attracted more strongly i.e. towards the more electronegative atom.


The strength of a dipole is measured by its **dipole moment**, μ , which is the product of the separated electric charge, $\delta+$ or $\delta-$, and the distance between them. The debye unit, D (named after the chemist Peter Debye), has the dimensions of coulomb-metre ($1 \text{ C}\cdot\text{m} = 2.9979 \times 10^{29} \text{ D}$).

The **bigger the difference** in the electronegativities of the atoms, the **larger the dipole moment**, and the **more polar** the covalent bond.

1.5.2 Bond Strength

The strength of a covalent bond is measured by its **bond energy**. The **stronger** the covalent bond, the **higher** the bond energy.

Bond energy is the **amount of energy** required to break **one mole of covalent bonds** in the **gaseous state**.



Energy is **released** during bond **formation** (i.e. this is an **exothermic** reaction)

Energy is **absorbed** during bond **breaking** (i.e. this is an **endothermic** reaction)

The strength of covalent bonds is generally dependent on the following factors:

- **bond order** i.e. number of covalent bonds formed between two atoms
- **effectiveness of orbital overlap**
- **bond polarity**

(i) Bond order

As bond order **increases**, the **number of bonding electrons within the inter-nuclei region increases**. Hence, the **electrostatic forces of attraction for these electrons increases**.

bond order	types of bond
1	single
2	double
3	triple

↓ Bond strength increases

Example: Strength of covalent bond: $\text{C}-\text{C} < \text{C}=\text{C} < \text{C}\equiv\text{C}$

	$\text{C}-\text{C}$	$\text{C}=\text{C}$	$\text{C}\equiv\text{C}$
bond energy / kJ mol^{-1}	350	610	840
bond length / nm	0.154	0.134	0.120

(ii) Effectiveness of orbital overlap

Generally, **larger orbitals are more diffuse**, which results in **less effective orbital overlap**. Hence, bonds formed by **larger atoms** will tend to have **lower bond strength (bond energy)**.

Example: Trend of bond energies of some Group 17 elements

	bond energy / kJ mol^{-1}
$\text{Cl}-\text{Cl}$	244
$\text{Br}-\text{Br}$	193
$\text{I}-\text{I}$	151

(iii) Bond polarity (i.e. unequal sharing of electrons in a covalent bond)

In a polar bond, there are **additional attractive forces between the δ^+ and δ^-** that **strengthens the covalent bond**, leading to **higher bond strength**.

The **more polar** the bond, the **stronger** the bond. This factor of bond polarity only comes into play when bonds have **comparable** bond order and effectiveness of orbital overlap.


Example:

	bond energy / kJ mol^{-1}	bond length / nm
$\text{C}\equiv\text{O}$	1070	0.113
$\text{N}\equiv\text{N}$	994	0.110

CO and N_2 have similar bond lengths. However, the $\text{C}\equiv\text{O}$ bond is polar while the $\text{N}\equiv\text{N}$ is not, leading to the $\text{C}\equiv\text{O}$ bond having higher bond strength.

1.5.3 Bond Length

Bond length is the **average distance** between the **nuclei of two atoms**.

**Example:**

	bond energy / kJ mol^{-1}
H-Cl	431
H-Br	366
H-I	299

Because of the **decrease** in the effectiveness of orbital overlap between the hydrogen atom and halogen atom, the bond length **increases**, and the bond strength (bond energy) **decreases**.

In summary, when


- (1) **bond order increases** or
- (2) **effectiveness of orbital overlap increases** or
- (3) **polarity of bond increases**

bond length decreases and **bond strength (bond energy) increases**.

Example 1B

As F is smaller than Cl, we would expect that F–F should have more effective overlap of orbitals and hence a stronger bond. However, based on the bond energy values, Cl–Cl is in fact a stronger bond. Suggest why this is so.

	bond energy / kJ mol^{-1}	atomic radius / nm
F–F	158	0.072
Cl–Cl	244	0.099



As the F atom is very small, the F–F bond length is very short. This results in the non-bonding electrons on F being in close proximity with those of the other F atom. Thus, repulsion occurs which weakens the covalent bond. This phenomenon happens in small molecules such as F₂ and O₂.

Checkpoint

- ☐ I know that all chemical bonds are basically electrostatic forces of attraction.
- ☐ I know the definition of a covalent bond.
- ☐ I know how sigma and pi bonds are formed, and how they lead to the formation of single, double and triple bonds.
- ☐ I know what dative bonding is, identify when it is formed, and how to represent it.
- ☐ I am able to identify if a bond is polar or non-polar.
- ☐ I am able to use the 3 factors (bond order, effectiveness of orbital overlap or bond polarity) to explain differences in bond strength/bond energy.


1.6 Representing Covalent Bonds: Dot-And-Cross Diagrams

LO (b) Describe, including the use of 'dot-and-cross' diagrams:

- (ii) covalent bonding, as in hydrogen; oxygen; nitrogen; chlorine; hydrogen chloride; carbon dioxide; methane; ethene
- (iii) co-ordinate (dative covalent) bonding, as in formation of the ammonium ion and in the Al_2Cl_6 molecule.

When an atom forms bonds with other atoms, normally only electrons in the **valence shell** are involved. When looking at bonding in *simple terms*, we can ignore the distinction between the various types of orbitals (s, p, d, etc.) in a shell and consider all electrons in the valence shell as a group.

If an atom has more than one electron in its valence shell, it can form more than one covalent bond with other atoms. For example, the beryllium atom ($1s^2 2s^2$) has a pair of 2s electrons. When forming a compound such as beryllium hydride, BeH_2 , these electrons can unpair themselves and form two bonds with two hydrogen atoms:



The boron atom ($1s^2 2s^2 2p^1$) can form three bonds as exemplified by boron hydride, BH_3 , while the carbon atom ($1s^2 2s^2 2p^2$) with four electrons in its valence shell can form four bonds as in methane, CH_4 , shown above.

Bonding diagrams often only show the electrons in the valence shell, omitting the inner shells.

How to draw dot-and-cross diagrams for species containing covalent bonds

Step 1: Identify the **central atom** in the **chemical formula**. The central atom is *usually* the atom with the **lowest electronegativity (except H)**.

Step 2: Determine the **number of valence electrons (by group number)** for each atom.

For an **anion**, add **electron(s)** to the **more electronegative** atom (F, O, Cl, N).
For a **cation**, remove **electron(s)** from the **less electronegative** atom.

Step 3: Determine the **type of bond formed** (single, double, triple or dative) between the central atom and its neighbouring atoms through sharing of valence electrons to achieve **noble gas electronic configurations**. Use **dots** and **crosses** on **alternate atoms**.

Step 4: Arrange the **remaining valence electrons (not involved in bonding) in pairs**. These electrons are known as **lone pairs of electrons** (non-bonding electrons).

Step 5: Check that each atom achieves a **stable noble gas configuration** (*ideal case*, but sometimes, the central atom does not achieve this). **Elements from period 3 onwards can expand octet** i.e. have more than 8 electrons around the atom (see Note on next page for the reason why). If the central atom cannot expand octet, consider changing a double bond to a dative bond.

Note:

- * Octet Rule (which was taught in secondary school) is based on empirical observation and it applies strictly to the **period 2 elements** (e.g. C, N, O, F) due to the availability of only the 2s and 2p orbitals in the valence shell.
- * Unlike elements in the second row of the Periodic Table, those in the third and subsequent rows can use their **d orbitals** in bonding, in addition to their s and p orbitals. Hence, they can **expand octet due to their vacant and energetically accessible d orbitals**. As a result, they are able to form more than four covalent bonds to other atoms (e.g. PCl_5 , SF_6).
- * For *many* elements, the number of bonds they can form in a molecule is fixed and is termed the **covalency** of the element. For example, oxygen has a covalency of 2, and carbon has a covalency of 4.
- * Elements in Groups 15 to 17 in the third and subsequent rows of the Periodic Table (Period 3 and higher) can display more than one covalency. For example, phosphorus has a covalency of 3 or 5, and chlorine has a covalency of 1, 3 or 5.

1.6.1 Dot-And-Cross for Molecules

	Valence Electrons / Bond Types	Dot-and-Cross Diagram
1) F_2O	F (Gp 17): 7 valence electrons O (Gp 16): 6 valence electrons \Rightarrow forms single bond with each F	
2) CO_2	C (Gp 14): 4 valence electrons O (Gp 16): 6 valence electrons \Rightarrow forms double bond with C	
3) HCN	C (Gp 14) : 4 valence electrons H: 1 valence electron \Rightarrow forms single bond with C N (Gp 15): 5 valence electrons \Rightarrow forms triple bond with C	
4) H_2O_2	*Note: TWO O central atoms O (Gp 16): 6 valence electrons \Rightarrow forms single bond with O and H H: 1 valence electron \Rightarrow forms single bond with O	

1.6.2 Dot-And-Cross Diagrams for Cations / Anions

	Valence Electrons / Bond Types	Dot-and-Cross Diagram
5) NO_2^+	Electronegativity: N < O Charge = +1 (remove 1 electron from N) N (Gp 15): $5 - 1 = 4$ valence electrons O (Gp 16): 6 valence electrons \Rightarrow forms double bond with N	
6) OCl^-	Electronegativity: O > Cl Charge = -1 (add 1 electron to O) O (Gp 16): $6 + 1 = 7$ valence electrons Cl (Gp 17): 7 valence electrons \Rightarrow forms single bond with O	
7) CO_3^{2-}	Electronegativity: O > C Charge = -2 (shared by 2 O atoms, add 1 electron to each O) C (Gp 14): 4 valence electrons 2 O (Gp 16) <u>each</u> : $6 + 1 = 7$ valence electrons <u>each</u> \Rightarrow forms single bond with C 1 O: 6 valence electrons \Rightarrow forms double bond with C	


1.6.3 Dot-And-Cross Diagrams for Electron Deficient (with Incomplete Octet) Molecules

		Dot-and-Cross Diagram
8) BeCl_2	Central atom from Group 2 and 13 (e.g. Be, B, Al) may have less than 8 valence electrons due to them having fewer valence electrons.	$\begin{array}{c} \times \times \\ \times \text{Cl} \times \bullet \text{Be} \bullet \times \text{Cl} \times \\ \times \times \end{array}$
9) BH_3		$\begin{array}{c} & & \text{H} \\ & & \times \\ \text{H} \times & \bullet & \text{B} \\ & \bullet & \\ & \times & \text{H} \end{array}$
10) AlCl_3		$\begin{array}{c} \times \times \\ \times \text{Cl} \times \\ \times \\ \times \text{Cl} \times \bullet \text{Al} \bullet \times \text{Cl} \times \\ \times \times \end{array}$

Electron deficient molecules may achieve a noble gas electronic configuration through **addition reactions**, resulting in the formation of **dative bonds**.


For example:

☼ Reaction between ammonia and boron hydride





☼ Dimerisation of aluminium chloride

The electron-deficient aluminium chloride may achieve a noble gas electronic configuration through the formation of a dimer.





1.6.4 Dot-And-Cross Diagrams for Species with Odd Number (Unpaired) Valence Electrons

Note: Central atom may have **fewer than 8** valence electrons.

		Dot-and-Cross Diagram
11) <u>NO</u>	Central atom may have odd number of valence electrons .	
12) <u>NO₂</u>	<p>Substances with an unpaired electron are called radicals and are usually very reactive.</p> <p>Radicals may achieve a noble gas electronic configuration through the formation of a dimer. e.g. $2\text{NO}_2 \rightarrow \text{N}_2\text{O}_4$</p>	

1.6.5 Dot-And-Cross Diagrams for Expansion of Octet Structures





		Dot-and-Cross Diagram
13) <u>PCl₅</u>	<p>Only elements from period 3 onwards have <u>vacant and energetically accessible d orbitals</u> to expand beyond octet structure (i.e. having more than eight valence electrons).</p>	
14) <u>SO₄²⁻</u>		

1.6.6 Dot-And-Cross Diagrams Showing Dative Covalent Bonds

	Valence Electrons/ Bond Types	Dot-and-Cross Diagram
15) $\underline{\text{N}}\text{O}_2$	<p>N (Gp 15): 5 valence electrons O (Gp 16): 6 valence electrons</p> <p>1 O forms double bond with N 1 O forms dative bond with N</p>	
16) $\text{H}_3\underline{\text{O}}^+$	<p>O (Gp 16): 6 valence electrons Electronegativity: H < O Charge = +1 (remove 1 electron from H) 1 H: $1 - 1 = 0$ \Rightarrow forms dative bond with O</p> <p>Remaining 2 H: 1 valence electron \Rightarrow each forms single bond with O</p>	

Self Check 1B

Draw the dot-and-cross diagrams for the following substances:

**Checkpoint**

- ☐ I know the steps to drawing dot-and-cross diagrams.
- ☐ I am able to draw dot-and cross-diagrams for molecules and polyatomic ions, knowing when they can have with odd electrons, incomplete octet, expanded octet and dative bonds.

2 Shape and Polarity






In a polyatomic species, the valence electron pairs of each atom, both bond pairs and lone pairs, repel one another to be as far as possible from each other. Hence, covalent molecules and polyatomic ions possess a definite, unique, three-dimensional shape which is determined by the angles between the bonds within them.



- LO (d) Explain the shapes of, and bond angles in, molecules such as BF_3 (trigonal planar); CO_2 (linear); CH_4 (tetrahedral); NH_3 (trigonal pyramidal); H_2O (bent); SF_6 (octahedral) by using the Valence Shell Electron Pair Repulsion theory.
- (e) Predict the shapes of, and bond angles in, molecules analogous to those specified in (d)

2.1 The Valence Shell Electron Pair Repulsion (VSEPR) Theory

The Valence Shell Electron Pair Repulsion (VSEPR) theory is used to predict the shape of a molecule or ion. The two general principles of VSEPR theory are:

Principle 1 **Electron pairs** (bond pairs and lone pairs) around a central atom are arranged **as far apart as possible** so as to **minimise repulsion**.

no. of electron pairs	shape	electron pairs are directed to...
2		opposite sides of each other
3		corners of an equilateral triangle
4		corners of a tetrahedron
5		corners of a trigonal bipyramid
6		corners of an octahedron


no. of electron pairs	2	3	4
example	$\text{O}=\text{C}=\text{O}$		
bond angle	180°	120°	109.5°

no. of electron regions	electron-pair geometry	no. of bond pairs	no. of lone pairs	molecular geometry (actual shape)	examples	bond angle
2	linear —A—	2	0	linear —A—	CO ₂ O=C=O	180°
3	trigonal planar 	3	0	trigonal planar 	BF ₃	120°
		2	1	bent 	SnCl ₂	<120° (e.g. 119°)
4	tetrahedral 	4	0	tetrahedral 	CH ₄	109.5°
		3	1	trigonal pyramidal 	NH ₃	107°
		2	2	bent 	H ₂ O	104.5°
5	trigonal bipyramidal 	5	0	trigonal bipyramidal 	PCl ₅	120° (equatorial) 90°(axial)
		4	1	see-saw / distorted tetrahedral 	SF ₄	<120° (equatorial) <90°(axial)
		3	2	T-shaped 	ClF ₃	<90°
		2	3	linear 	[ICl ₂] ⁻	180°
6	octahedral 	6	0	octahedral 	SF ₆	90° (equatorial & axial)
		5	1	square pyramidal 	IF ₅	<90° (equatorial & axial)
		4	2	square planar 	XeF ₄	90°


Note: Use wedge and dash notation to represent tetrahedral, trigonal pyramidal and see-saw. Indicate the plane clearly with dotted lines for trigonal bipyramidal, octahedral and square pyramidal.

Example 2A


1. Draw the structure of the following compounds and predict their shapes and bond angles.
 (a) PCl_4^+ (b) IF_4^+ (c) ICl_4^-



tetrahedral
 109.5°



see-saw
 $<90^\circ, <120^\circ$



square planar
 90°

2. Which one of the following pairs do the molecules have similar shapes?

A BeCl_2 and H_2O

A: BeCl_2 – linear, H_2O – bent

B BF_3 and NH_3

B: BF_3 – trigonal planar, NH_3 – trigonal pyramidal

C AlCl_3 and BCl_3


C: AlCl_3 and BCl_3 – trigonal planar

D CO_2 and SO_2

D: CO_2 – linear, SO_2 – bent

Answer: C

3. Predict the shape and bond angle about the nitrogen atoms in di-imine, $\text{HN}=\text{NH}$.




About each N atom, there are **2 bond pairs and 1 lone pair**.

To minimise repulsion, these three electron pairs are directed to the corners of an equilateral triangle.

Since **lone pair–bond pair repulsion > bond pair–bond pair repulsion**, bond angle of H–N=N is **less than 120° (e.g. 119°)** and di-imine is **bent** about N.


Special Cases: While bond angle typically is determined by the number of lone pair and bond pairs around the central atom, the additional factor that can affect the bond angles of molecules is the **electronegativity of the central atom or the terminal atoms**.

The **repulsion between the neighbouring electron pairs is decreased when the central atom has a lower electronegativity or when the terminal atoms have greater electronegativity**. In both cases, the electron density around the central atom decreases, leading to less repulsion between the bonding electron pairs.



Both H_2O and OF_2 have two bond pairs and two lone pairs, but OF_2 has a smaller bond angle than H_2O . Why?


- In OF_2 , the more electronegative terminal F atoms draw electron density away from the central O atom.
- The two bond pairs of electrons are further away from the nucleus of the central O atom.
- Thus, the two bond pairs of electrons experience less repulsion between them as compared to the bond pair of electrons in H_2O , resulting in a smaller bond angle in OF_2 .



2.2 Polarity of Molecules


LO (g) Deduce the polarity of a molecule using bond polarity and its molecular shape (analogous to those specified in (d)).

Each of the polar bonds in a molecule has its own dipole associated with it.




The **net dipole moment** of the **molecule** depends on its **shape**. Depending on the relative angles between the bonds, the individual bond dipoles can:

either **reinforce each other**




- If the bond dipoles reinforce each other, molecules with very large dipoles can be formed.

or **cancel each other**



If **cancellation is complete**, the resulting molecule will have **no net dipole moment**, and so the molecule will be **non-polar**.



Other examples (with the same terminal atoms):





structure	shape	(bond) dipole moments...	polarity
$\text{O}=\text{C}=\text{O}$	linear	cancel off	non-polar
	trigonal planar	cancel off	non-polar
	bent	do not cancel off	polar
	tetrahedral	cancel off	non-polar
	trigonal pyramidal	do not cancel off	polar
	bent	do not cancel off	polar
	trigonal bipyramidal	cancel off	non-polar
	See-saw (distorted tetrahedral)	do not cancel off	polar
	T-shaped	do not cancel off	polar
	linear	cancel off	non-polar
	octahedral	cancel off	non-polar
	square pyramidal	do not cancel off	polar
	square planar	cancel off	non-polar

 To determine if a molecule is **polar** or **non-polar**

- determine the actual shape of the molecule
- consider the electronegativity of the various atoms present in the molecule
- determine the net dipole moment by resolving the individual dipole moments

Example 2B

1. Which molecule has the largest dipole?



Answer: B

A: C–H bonds are generally considered to be non-polar.

B: Dipole moment due to C=O bond.

C: Dipole moments due to C=O and C–Cl bonds cancel out to a certain extent, hence the net dipole moment is smaller than that in option B.


D: Dipole moments of C–Cl bonds cancel out.

Comparing the Polarities of NH_3 and NF_3 (for enrichment)

Despite the differences in electronegativities between N and H, and between N and F, being similar, NH_3 has a much higher dipole moment compared to NF_3 .

This is because dipoles are induced not just in polar bonds, but also in lone pairs of electrons. The lone pair of electrons is an area of negative charge, so it forms the δ^- end of a dipole which points away from the positively charged nucleus of the atom.

In NH_3 , the dipole due to the lone pair helps reinforce the dipoles from the N–H bonds, resulting in a highly polar molecule. However, in the case of NF_3 , this dipole partially offsets the dipole due to the N–F bonds, resulting in a smaller resultant dipole.

**Checkpoint**

- ☐ I am able to predict the shape of the molecule and its bond angle.
- ☐ I am able to represent the shape of molecules and polyatomic ions clearly, using dash and wedge where necessary, and drawing the bonds at appropriate angles to each other.
- ☐ I am able to predict if a molecule is polar or non-polar.

3 Intermolecular Forces of Attraction

The covalently bonded molecules we have looked at so far have **simple molecular structures**.

Substances made up of such molecules are often gases or liquids at room temperature. This is because, although the **intramolecular** covalent bonds *within* each molecule are very strong, the **intermolecular forces of attraction** *between* one molecule and another are comparatively weaker and require less energy to overcome.

There are three main categories of intermolecular forces of attraction:

- **instantaneous dipole–induced dipole interactions**
- **permanent dipole–permanent dipole interactions**
- **hydrogen bonding**

LO (h) Describe the following forces of attraction (electrostatic in nature):


- intermolecular forces, based on permanent and induced dipoles, as in $\text{CHCl}_3(l)$; $\text{Br}_2(l)$ and the liquid noble gases
- hydrogen bonding, using ammonia and water as examples of molecules containing $-\text{NH}$ and $-\text{OH}$ groups

3.1 Instantaneous Dipole–Induced Dipole Interactions

Temporary fluctuations in the electron distributions within atoms and non-polar molecules could result in the formation of short-lived **instantaneous dipoles**, which produce attractive forces between otherwise non-polar substances. These attractive interactions are weak and fall off rapidly with increasing distance.


Consider a pair of adjacent He atoms:

- On average, the two electrons in each He atom are uniformly distributed within the atom. However, because the electrons are in constant motion, their distribution in one atom is likely to be *asymmetrical* at any given instant, resulting in an **instantaneous dipole** that lasts in the order of 10^{-16} seconds:



- As shown above, the instantaneous dipole on one atom can interact with the electrons in an adjacent atom, pulling them towards the positive end of the instantaneous dipole or repelling them from the negative end. The net effect is that the first atom causes the **temporary** formation of a dipole, called an **induced dipole**, in the second. Interactions between these temporary dipoles cause atoms to be attracted to one another.
- An instant later, the first atom might change its dipole through another movement of electrons within it, so the original dipole attraction might be destroyed. This changed dipole will induce new dipoles in another neighbouring atom, however, and so new intermolecular forces of attraction will be formed.

Similar interactions are formed between molecules such as hydrogen gas.







The **strength of instantaneous dipole–induced dipole interactions** is related to the ease of polarisation of the electron cloud in an atom or molecule (i.e. their **polarisability**)

The polarisability of the electron cloud of a substance depends on:

(a) Number of electrons


- When a molecule has **more electrons**, it will have a **bigger and more polarisable electron cloud**.
- This leads to **stronger** instantaneous dipole–induced dipole interactions, which requires **more energy** to overcome.

molecule	boiling point / °C	reasons
CH ₄	−164	These molecules have the same shape (tetrahedral), but the number of electrons increases going down the group, from 10 in CH ₄ to 18 in SiH ₄ to 36 in GeH ₄ and 54 in SnH ₄ .
SiH ₄	−112	
GeH ₄	−89	
SnH ₄	−52	
F ₂	−188	All of these are linear, diatomic molecules. Down the group, the number of electrons increases, from 18 in F ₂ , to 34 in Cl ₂ , to 70 in Br ₂ and 106 in I ₂ .
Cl ₂	−35	
Br ₂	59	
I ₂	184	
CH ₄	−164	Along the alkane series, the molecules contain more electrons (an extra 8 for each CH ₂ group).
CH ₃ CH ₃	−89	
CH ₃ CH ₂ CH ₃	−42	
CH ₃ CH ₂ CH ₂ CH ₃	−1	
<div><div><p>Methane</p></div><div><p>Ethane</p></div><div><p>Propane</p></div><div><p><i>n</i>-Butane</p></div></div>		


(b) **Shape of molecule** (refers to **elongated** versus **spherical** shape)

- Compared to spherical molecules with similar M_r , **elongated molecules have greater surface area of contact.**
- This leads to **stronger** instantaneous dipole-induced dipole interactions, which requires **more energy** to overcome.

For example, 2,2-dimethylpropane and pentane are **isomers** i.e. both have the same chemical formula, C_5H_{12} , but different arrangement of atoms.




Since they have the same chemical formula, they will have the same number of electrons. However, 2,2-dimethylpropane has a lower boiling point of $9.5\text{ }^{\circ}\text{C}$, while pentane boils at a higher temperature of $36.1\text{ }^{\circ}\text{C}$. This is due to the difference in the shapes of the molecules.




3.2 Permanent Dipole–Permanent Dipole Interactions


Uneven distribution of electronic charge within **polar molecules** results in a permanent dipole. The positive end of one molecule's dipole can attract the negative end of another polar molecule.



Similarly, repulsive dipole-dipole interactions can occur if two positive or two negative ends are adjacent to each other.




Since molecules in liquid and gas phase move freely and continuously, polar molecules always experience both attractive and repulsive dipole-dipole interactions simultaneously.



On average, however, the attractive interactions dominate, resulting in an intermolecular attraction called **permanent dipole–permanent dipole interactions**.

An example of permanent dipole–permanent dipole interactions is the attractions between HCl molecules.



Within a series of compounds of similar molar mass, the **strength** of the permanent dipole–permanent dipole interactions **increases** as the **dipole moment** of the **molecules** increases, *i.e.* as the molecules become **more polar**.

compound	molar mass /g mol ⁻¹	dipole moment /D	boiling point /K
(CH ₃) ₃ CH (methylpropane)	58	0.132	261
C ₂ H ₅ OCH ₃ (methoxyethane)	60	1.17	281
CH ₃ CH ₂ CHO (propanal)	58	2.72	322
CH ₃ COCH ₃ (propanone)	58	2.88	329

3.3 Hydrogen Bonding

When a **hydrogen atom is bonded to a very electronegative atom such as nitrogen, oxygen, or fluorine**, it will have a significant partial positive charge (δ^+). Hence, this hydrogen atom with a large δ^+ can experience a particularly strong attraction to a **lone pair of electrons on a nitrogen, oxygen, or fluorine atom** in an adjacent molecule. An intermolecular attractive force called a **hydrogen bond** is thus created.

Hydrogen bond is defined as the **electrostatic forces of attraction** between the **partial positive (δ^+) hydrogen atom that is directly bonded to a very electronegative F, O or N atom** and a **lone pair of electrons** of another **F, O or N atom**.

While the name “hydrogen bond” makes such an electrostatic force of interaction sound like it is a type of covalent bond, it is not. In fact, hydrogen bonds are about 10 times weaker than a typical covalent bond.

Hydrogen bonds are represented by dotted lines between the δ^+ hydrogen and the lone pair of electrons on a very electronegative element (F, O, N).



$\cdots \cdots \cdots$ hydrogen bond
X, Y = F, O, N

Note: Always include the **lone pair** and **partial charges** in a diagram showing hydrogen bonds.

The **strength of hydrogen bonds increases** when the **electronegativity of the atom bonded to the δ^+ hydrogen atom increases**. For example, HF has higher boiling point than NH_3 because the F atom in HF is more electronegative than the N atom in NH_3 . Hence, HF has stronger hydrogen bonding than NH_3 .

However, if we compare the boiling point of H_2O to that of HF, we realise that the above explanation does not work. O atom in H_2O is less electronegative than the F atom in HF, but H_2O has a higher boiling point.


molecule (boiling point)	HF (20 °C)	H_2O (100 °C)	NH_3 (-33 °C)
diagram			
no. of lone pairs (on each F/O/N)	3	2	1
no. of H atoms	1	2	3
no. of H bonds (per molecule)	1	2	1
electronegativity	F	> O	> N

In this case, we need to consider the number of hydrogen bonds formed per molecule, or the **extensiveness of hydrogen bonding**. H_2O has the highest boiling point because H_2O has **more extensive** hydrogen bonding than HF and NH_3 . It can form **two hydrogen bonds per water molecule** whereas both NH_3 and HF can **only form one hydrogen bond per molecule**.

3.4 Comparing the Three Types of Intermolecular Forces of Attraction

The most significant intermolecular forces of attraction that exist in a species depends on the polarity of the molecule. The flow chart below shows an easy way to help you determine the intermolecular forces of attraction in a given covalent compound.

Note that instantaneous dipole-induced dipole interactions exist in **all** molecules, not just in non-polar molecules. However, due to the relative strengths of the intermolecular forces of attraction, it may not be the most significant in polar molecules.






Generally, for molecules with **similar** number of electrons, the relative strengths of intermolecular forces of attraction are in the order of:

hydrogen bonds	>	permanent dipole– permanent dipole interactions	>	instantaneous dipole– induced dipole interactions
---------------------------	-------------	--	-------------	--

This has implications on physical properties, such as boiling points of covalent compounds.


In the examples below, the compounds are grouped together by similar number of electrons. This implies that the instantaneous dipole–induced dipole interactions will be similar in strength within each grouping.

compound	formula	polarity	type of IMF	no. of electrons	boiling point / °C
ethane	CH ₃ –CH ₃	non-polar	id–id	18	–89
methanal		polar	pd–pd	16	–19
methanol	CH ₃ –OH	polar	hydrogen bonding	18	+65
propane	CH ₃ –CH ₂ –CH ₃	non-polar	id–id	26	–42
ethanal		polar	pd–pd	24	+20
ethanol	CH ₃ –CH ₂ –OH	polar	hydrogen bonding	26	+79
butane	CH ₃ –CH ₂ –CH ₂ –CH ₃	non-polar	id–id	34	0
propanal		polar	pd–pd	32	+49
propan-1-ol	CH ₃ –CH ₂ –CH ₂ –OH	polar	hydrogen bonding	34	+97

The first compound in each grouping is a hydrocarbon which is essentially non-polar. Molecules of these compounds are held together by instantaneous dipole–induced dipole interactions *only*. The second compound in each grouping is an aldehyde with a polar –CHO group (refer to propanal). Molecules of these compounds are held together by stronger permanent dipole–permanent dipole interactions, on top of instantaneous dipole–induced dipole interactions, leading to higher boiling points. The last compound in each grouping is an alcohol with a polar –OH group, which is capable of forming hydrogen bonds. These molecules are held together by even stronger hydrogen bonding, on top of the permanent dipole–permanent dipole interactions and instantaneous dipole–induced dipole interactions. Hence, they have the highest boiling points compared to the rest in the group.

Example 3A

1. Consider the following three compounds:



Explain why compound 3 has a higher boiling point than compound 1.

Both compounds have simple molecular structure.

Compound 1 is a branched molecule while compound 3 is a straight-chain molecule.

Compound 3 has greater surface area of contact compared to compound 1.



Hence, Compound 3 has stronger instantaneous dipole–induced dipole interactions than compound 1 which requires more energy to overcome.

Explain why compound 2 has a higher boiling point than compound 3.

Compound 2 is a **simple molecular** compound with **stronger hydrogen bonds**, while compound 3 is a **simple molecular** compound with **weaker instantaneous dipole–induced dipole interactions**.

More energy is required to overcome the **stronger hydrogen bonds** of compound 2.

2. Which of the following statements best explains why the boiling point of butanone, $\text{C}_4\text{H}_8\text{O}$ (79.6 °C) is higher than that of pentane, C_5H_{12} (36.1 °C)?



- A The relative molecular mass of butanone is higher than that of pentane.
- B The butanone molecule has a larger surface area than the pentane molecule.
- C There are hydrogen bonds between butanone molecules, but not between pentane molecules.
- D There are permanent dipole–permanent dipole interactions between butanone molecules, but only instantaneous dipole–induced dipole interactions between pentane molecules.

Answer: D

Pentane is non-polar while butanone is polar. Thus, butanone will have permanent dipole–permanent dipole interactions which are stronger and requires more energy to overcome than the instantaneous dipole–induced dipole interactions that pentane has. Thus, butanone would have a higher boiling point than pentane.

Self Check 3A

1. The gecko, a small lizard, can climb up a smooth glass window. The gecko has millions of microscopic hairs on its toes and each hair has thousands of pads on its tip. The result is that the molecules in the pads are extremely close to the glass surface on which the gecko is climbing.

What is the main attractive force between the gecko's toe pads and the glass surface?

- A co-ordinate bonds
 - B covalent bonds
 - C ionic bonds
 - D instantaneous dipole–induced dipole interactions
2. The boiling point of water (100 °C) is greater than that of HF (20 °C)? Which statement is a correct explanation of this?
- A Each hydrogen bond formed between water molecules is stronger than that formed between HF molecules.
 - B There are more atoms in a water molecule than there are in a HF molecule, so the permanent dipole–permanent dipole forces are stronger in water.
 - C There are, on average, more hydrogen bonds between water molecules than there are between HF molecules.
 - D The water molecule has a greater permanent dipole than the HF molecule.

Checkpoint





- ☐ I know what the 3 types of intermolecular forces are.
- ☐ I know the factors that affect the strengths (and extensiveness for hydrogen bonds) of the various intermolecular forces of attraction.
- ☐ I am able to identify the type(s) of intermolecular forces of attraction present in any given substance.
- ☐ I am able to explain melting and boiling point differences based on the strengths of the different intermolecular forces of attraction.

4 Covalent Substances and Their Physical Properties

- LO (i) Outline the importance of hydrogen bonding to the physical properties of substances including ice and water.
- (I) Describe, in simple terms, the lattice structure of a crystalline solid which is:
- (ii) simple molecular, as in iodine
 - (iii) giant molecular, as in graphite and diamond
 - (iv) hydrogen-bonded, as in ice

4.1 Simple Molecules

In substances with simple molecular structures (extends to monoatomic substances as well), there are intermolecular (or interatomic) forces of attraction, such as **instantaneous dipole-induced dipole interactions**, **permanent dipole-permanent dipole interactions**, and **hydrogen bonding**, which are **weak** compared to intramolecular covalent bonds.



In general, substances with simple molecular structures exhibit the following physical properties:

Property	Reasons
Low melting/ boiling point	<p>Small amount of energy is needed to overcome either the weak instantaneous dipole-induced dipole interactions/ permanent dipole-permanent dipole interactions/ hydrogen bonds.</p> <p>Note: When a substance consisting of simple covalent molecules melts or boils, covalent bonds within the molecules are NOT broken (unless the substance decomposes).</p>
Non-conductors of electricity in any state	Electrons are localised in covalent bonds and are not mobile to conduct electricity.

Property	Reasons				
Generally soluble in non-polar solvents (e.g. benzene, CCl_4 , hexane); insoluble in polar solvents (e.g. water)	In general, substances with similar types of intermolecular forces of attraction dissolve in each other; “ like dissolves like ”. Dissolution can be a physical (e.g. dissolve ethanol in water) or a chemical process (e.g. dissolve HCl in water). To predict solubility we need to consider the relative strengths of solute–solvent interactions, solute-solute interactions and solvent–solvent interactions. For a solute to be soluble, solute–solvent interactions must be comparable in strength as solute–solute and solvent–solvent interactions.				
	example	I_2 in water	I_2 in hexane	$\text{CH}_3\text{CH}_2\text{OH}$ in water	HCl in water
	interaction	solute–solute	instantaneous dipole–induced dipole	instantaneous dipole–induced dipole	hydrogen bonding
		solvent–solvent	hydrogen bonding	instantaneous dipole–induced dipole	hydrogen bonding
		compare	stronger than	approximately the same as	approximately the same as
		solute–solvent	instantaneous dipole–induced dipole	instantaneous dipole–induced dipole	hydrogen bonding
	solubility	insoluble	soluble	soluble	soluble
* HCl ionises in water : $\text{HCl}(\text{aq}) \rightarrow \text{H}^+(\text{aq}) + \text{Cl}^-(\text{aq})$ (refer to pg. 49 for more information about ion-dipole interactions)					

Note: The mnemonic “like dissolves like” is coined to help you predict the solubility of solutes in specific solvents. It is NOT a scientific explanation!

Example 4A


Explain why $\text{CH}_3\text{CH}_2\text{OH}$ is soluble in CH_3COCH_3 .

The energy released in forming hydrogen bonds between the $\text{CH}_3\text{CH}_2\text{OH}$ and CH_3COCH_3 molecules (i.e. solute-solvent interactions) is sufficient to overcome the hydrogen bonds between the $\text{CH}_3\text{CH}_2\text{OH}$ molecules (i.e. solute-solute interactions) and permanent dipole–permanent dipole interactions between the CH_3COCH_3 molecules (i.e. solvent–solvent interactions). Thus, $\text{CH}_3\text{CH}_2\text{OH}$ is soluble in CH_3COCH_3 .

4.2 Influence of Hydrogen Bonding on Physical Properties of Molecules

4.2.1 Unusually high boiling points of ammonia, water and hydrogen fluoride

when compared to other hydrides in the same group:



All hydrides in each group have **simple molecular** structures.

As **number of electrons increases** down each group, **strength of instantaneous dipole-induced dipole interactions increases**. A **larger amount of energy** is required to overcome the stronger instantaneous dipole-induced dipole interactions. Hence, boiling point of hydrides increases down the group.


However, boiling points of ammonia, water and hydrogen fluoride are **higher than expected**. This is because **more** energy is needed to overcome the **stronger hydrogen bonds** in **ammonia, water** and **hydrogen fluoride** compared to the **weaker instantaneous dipole-induced dipole interactions** in other hydrides.

(See Section 3.3 for the comparison between the boiling points of ammonia, water, and hydrogen fluoride.)

4.2.2 Anomalous relative molecular masses of organic acids ($R-CO_2H$)

Ethanoic acid, CH_3CO_2H , has an **apparent M_r of 120.0** which is **double** the expected M_r of 60.0 in the gas phase or when dissolved in benzene.


This is due to each acid molecule pairing up to form a **cyclic (closed ring) dimer** *via hydrogen bonds*.




4.2.3 High solubility of ammonia, alcohols ($R-OH$), organic acids and amines ($R-NH_2$) in water

These substances can form **hydrogen bonds** with water molecules as shown below:

CH_3NH_2 soluble in H_2O



CH_3OH (methanol) soluble in H_2O




Note: When illustrating solubility in water via hydrogen bonding, show:

- 1 at least 1 solute molecule and 1 water molecule
- 2 hydrogen bonds,
- 3 lone pair of electrons,
- 4 partial charges (dipoles)
- 5 labels for hydrogen bonds



4.2.4 Different physical properties of isomers

4-nitrophenol has a boiling point of 279 °C, while that of 2-nitrophenol is **lower** at 216 °C. Also, 4-nitrophenol is **more soluble in water** compared to 2-nitrophenol.




☞ **Intermolecular** hydrogen bond: formed **BETWEEN** different molecules.

☞ **Intramolecular** hydrogen bond: formed **WITHIN** the **same** molecule.

	4-nitrophenol	2-nitrophenol
type of interactions	 <p>intermolecular hydrogen bonding</p> <p>Hydrogen bonds formed between two 4-nitrophenol molecules.</p> <p>⇒ more extensive intermolecular hydrogen bonding</p>	 <p>intramolecular hydrogen bonding</p> <p>Hydrogen bond formed within one 2-nitrophenol molecule due to the -NO₂ and -OH groups being close together in 2-nitrophenol.</p> <p>⇒ more extensive intramolecular hydrogen bonding</p> <p>⇒ less extensive intermolecular hydrogen bonding</p>
boiling point	<p>More energy is needed to overcome the more extensive intermolecular hydrogen bonds in 4-nitrophenol.</p> <p>Hence, 4-nitrophenol has a higher boiling point</p>	<p>Less energy is needed to overcome the less extensive intermolecular hydrogen bonds in 2-nitrophenol.</p> <p>Hence, 2-nitrophenol has a lower boiling point</p>
solubility	<p>More extensive intermolecular hydrogen bonds can be formed between molecules of 4-nitrophenol and water.</p> <p>Hence, 4-nitrophenol is more soluble in water.</p>	<p>Less extensive intermolecular hydrogen bonds can be formed between molecules of 2-nitrophenol and water.</p> <p>Hence, 2-nitrophenol is less soluble in water.</p>

4.2.5 Higher density of water compared to ice

Hydrogen bonding holds ice molecules in place in an **orderly open structure** with **large amount of empty spaces** in between.




In this extended network, each water molecule forms **two** hydrogen bonds with its neighbouring water molecules such that each oxygen atom is surrounded by **four** hydrogen atoms.

Water molecules in the liquid state are able to occupy the spaces in between when ice melts. More molecules present in a fixed volume accounts for the higher density of water.

4.3 Giant Molecular Structures

Giant molecules can have **giant covalent** or **giant molecular structures**. Such structures consist of **extensive three-dimensional arrays of atoms**, which can either be all of the same type, as in the elements carbon (diamond and graphite) and silicon, or of two different elements such as in silicon(IV) oxide, SiO_2 .




Think about this: If each Si atom is bonded to 4 O atoms and each O atom is bonded to 2 Si atoms, why is the chemical formula of silicon(IV) oxide SiO_2 , instead of Si_2O_4 ?

The **atoms are all joined to one another by strong covalent bonds**. So a single crystal of diamond or quartz is in fact a single molecule. For atoms to become free from the lattice, these strong bonds must be broken. Therefore, substances containing this type of lattice have very high melting and boiling points:

substance	formula	type of interatomic attraction	melting point/ $^{\circ}\text{C}$	boiling point/ $^{\circ}\text{C}$
silicon	Si	covalent	1412	3265
silicon(IV) oxide	SiO_2	covalent	1610	2959
diamond	C	covalent	3500	3930

Property	Reasons
High melting / boiling point	Large amount of energy is needed to overcome the strong covalent bonds between atoms in a giant molecular structure .
Non-conductors of electricity in any state	Electrons are localised in covalent bonds and are not mobile to conduct electricity.
Insoluble in both water and non-polar solvents	No favourable solute-solvent interactions between solute particles and water / solvent molecules can be formed to break down the giant molecular structure for hydration / solvation.
Hard, strong and non-malleable	Atoms are held closely together by strong covalent bonds in a giant molecular structure . (Can be used as abrasives due to their hardness and high melting points , e.g. diamond-tipped tools is used to cut rocks)

Giant molecular structure of graphite consists of **atoms** held together by **strong covalent bonds** in **giant extensive planar layers** with the **adjacent layers** held together by **weak instantaneous dipole–induced dipole interactions**.



substance	formula	type of interatomic attraction	melting point/°C	boiling point/°C
boron(III) nitride	BN	covalent	2967	2500 (sublime)
graphite	C	covalent	–	3915 (sublime)

Property of graphite	Reasons
High melting/boiling point	Large amount of energy is needed to overcome the strong covalent bonds between atoms in the giant molecular layered structure .
Good conductors of electricity parallel to the layers but non-conductors of electricity perpendicular to the layers	3 of the 4 valence electrons are used to form covalent bonds, the 4th electron is delocalised parallel to the whole layer . The presence of delocalised electrons act as mobile charge carriers to conduct electricity along the plane . Electrons cannot delocalise between the layers.
Insoluble in both water and non-polar solvents	No favourable solute–solvent interactions between solute particles and water / solvent molecules can be formed to break down the giant molecular layered structure for hydration / solvation.
Soft and slippery	Adjacent layers are held together by weak instantaneous dipole–induced dipole interactions and so the layers can easily slide over one another. (Can be used as a lubricant because of this property, as well as its ability to withstand high temperatures due to its high melting point).

Self Check 4A

1. Silicon carbide (carborundum) is a shiny, hard, chemically inert material with a very high melting point. It can be used to sharpen knives and make crucibles. Which type of structure explains these properties?
- A** a giant structure with covalent bonds between silicon and carbon atoms
 - B** a giant structure containing metallic bonding
 - C** a giant layered structure with covalent bonds between atoms and instantaneous dipole-induced dipole interactions between the layers
 - D** a simple molecular structure with covalent bonds between the atoms of silicon and carbon
2. Which of the following statements correctly describe the graphite lattice?
- 1** The lattice contains delocalised electrons.
 - 2** Each carbon atom in the lattice has three closest neighbours.
 - 3** The layers of carbon atoms are held together by strong covalent bonds.
- A** 1, 2 and 3 are correct
 - B** 1 and 2 only are correct
 - C** 1 and 3 only are correct
 - D** 2 and 3 only are correct

Checkpoint

- ☐ I am able to explain the physical properties observed in substances with simple molecular structure by relating to the structure and bonding of these substances.
- ☐ I am able to use hydrogen bonding to explain certain unusual physical properties e.g. boiling points, anomalous relative molecular mass, solubility and density differences.
- ☐ I am able to explain the physical properties observed in substances with giant molecular structure by relating to the structure and bonding of these substances.

5 Ionic Bonding


- LO (a) Show understanding that all chemical bonds are electrostatic in nature and describe:
- (i) ionic bond as the electrostatic attraction between oppositely charged ions;
- (b) Describe, including the use of 'dot-and-cross' diagrams:
- (ii) ionic bonding, as in sodium chloride and magnesium oxide;
- (l) Describe, in simple terms, the lattice structure of a crystalline solid which is:
- (i) ionic, as in sodium chloride and magnesium oxide

5.1 Formation of Ionic Bond

Ionic bonds are **strong electrostatic forces of attraction** between **oppositely charged ions** in a giant ionic lattice structure

Ionic bonds are typically formed between a metallic element and a non-metallic element, when one atom (the metal) totally gives away one or more electrons to the non-metallic atom. This results in **ions** being formed.

Atoms with *low ionisation energies* can lose valence electrons to become positive ions, known as **cations**. The electrons are donated to atoms to form negative ions, known as **anions**. This can be represented using a '**dot-and-cross**' diagram:



Dots and crosses distinguish where the electron comes from, although in reality there is no difference between them. The charge on the ion should be shown as a superscript outside the square bracket.

The ions that make up ionic compounds are not always monatomic, *i.e.*, they do not always contain *only one atom*. Several common ions contain *groups* of atoms which are **covalently bonded** together, and which have an overall positive or negative charge e.g. SO_4^{2-} and NH_4^+ .

5.2 Factors affecting Ionic Bond Strength

The strength of an **ionic bond** is measured by the **magnitude of lattice energy** (ΔH_{latt}) of the ionic compound.

Lattice energy, ΔH_{latt} of an ionic crystal is the **energy released** when **one mole** of the **solid ionic compound** is **formed from its constituent gaseous ions**.



$$|\Delta H_{\text{latt}}| \propto \left| \frac{q_+ \times q_-}{r_+ + r_-} \right|$$

where q_+ and q_- : charges on the cation and anion respectively

r_+ and r_- : ionic radii of cation and anion respectively

Generally, the **greater** the **magnitude of lattice energy**, the **stronger** the ionic bonds. Also, generally the **difference in charge is more important than the difference in radius**.

Example 5A

In terms of structure and bonding, explain why MgO (m.p. 2850 °C) has a higher melting point than Na₂O (m.p. 1130 °C).

Both MgO and Na₂O have giant ionic lattice structures held together by strong electrostatic forces of attraction between oppositely charged ions.

$$|\Delta H_{\text{latt}}| \propto \left| \frac{q_+ \times q_-}{r_+ + r_-} \right|$$


Both compounds have the same anionic charge and radius.

However, Mg²⁺ has a higher ionic charge and smaller ionic radius than Na⁺.

More energy is required to overcome the stronger ionic bonds in MgO. Hence, MgO has a higher melting point.

For Your Info

Ionic bonding involves a separation of charge, which is usually an unfavourable endothermic process. The reason why it becomes more favourable with metals is their comparatively low ionisation energies. In sodium chloride, NaCl, for example, the outer electron is fairly easily lost from the sodium atom and becomes more attracted to the chlorine nucleus than to its original sodium nucleus.



Therefore, in the gas phase:



The fact that ΔH^\ominus is positive means that this is an energetically unfavourable process.

But when the ions form a solid lattice, the ions attract one another, and much energy is released:



Hence, the overall energy change, from gas-phase atoms to solid compound, is:




The complete process is seen to be highly exothermic, and therefore favourable. The benefits of total electron transfer are completed by the resulting electrostatic forces of attraction between the cation and anion that are formed.

5.3 Giant Ionic Lattice




In an ionic lattice of a solid, ions are arranged in a **fixed, orderly** manner in a **giant ionic lattice** structure.

- Each cation is surrounded by a number of anions as its closest neighbours.
- Each anion is surrounded by a number of cations as its closest neighbours.
- No two cations or anions will be adjacent to each other.
- The electrostatic forces of attraction exists between ions of opposite charge. Hence, an ionic bond is said to be **non-directional**.

In the case of sodium chloride, NaCl , each Na^+ ion is surrounded by six of the larger Cl^- ions while each Cl^- ion is surrounded by six Na^+ ions. (Refer to Appendix 5 for information on Coordination Number for ionic compounds)



5.4 Physical Properties of Ionic Compounds

Physical property	Reasons
High melting/ boiling point	Large amount of energy is needed to overcome the strong electrostatic forces of attraction between oppositely charged ions .
Strong, hard but brittle	<p>Hard: oppositely charged ions are held closely together by strong electrostatic forces of attraction in a giant ionic lattice structure.</p>  <p>Brittle: stress applied on the ionic lattice with regular pattern causes sliding of layers of ions resulting in ions of similar charges coming together and resultant repulsion shatters the ionic lattice structure.</p>
Non electrical conductors in solid state	Ions can only vibrate about fixed positions and hence are not mobile to conduct electricity.
Good electrical conductors in molten/aqueous state	<p>Presence of ions as mobile charge carriers to conduct electricity under the influence of an electric current/ field.</p> <p>Electrical conductivity increases with increasing concentration of the ions in the solution.</p>
Generally, soluble in polar solvent (e.g. water) i.e. solute particles surrounded by solvent (water) molecules	<p>Favourable ion-dipole interactions form between ions and water molecules. Hydration, whereby the ions are surrounded by water molecules, has occurred.</p> <p>The interactions formed will release energy to overcome the strong electrostatic forces of attraction between oppositely charged ions in the giant ionic lattice structure.</p> <p>The giant ionic lattice structure breaks down, and the solid dissolves.</p>
<p align="center"><u>Diagrammatic representation of ion-dipole interactions</u></p> <p>δ^- end of water attracted to positive ions and δ^+ end of water attracted to negative ions forming ion-dipole interactions:</p> <div style="display: flex; align-items: center; justify-content: space-around;">  <div style="border: 1px solid black; padding: 5px; text-align: center;"> <p>The "δ^-" oxygen ends in H_2O are attracted to the cations</p> </div>  <div style="border: 1px solid black; padding: 5px; text-align: center;"> <p>The "δ^+" hydrogen ends in H_2O are attracted to the anions</p> </div> </div>	
Insoluble in non-polar solvents (e.g. benzene, CCl_4 , hexane)	No favourable ion-dipole interactions between ions and non-polar solvent molecules can be formed. The giant ionic lattice structure does not break down.


Self Check 5B

Electric cable, as used in houses, is made of copper wire surrounded by poly(chloroethene) also called polyvinylchloride or PVC. However, electric cable used in fire alarm systems has copper wire surrounded by magnesium oxide which acts as an insulator. The whole cable is encased in thin copper tubing.

- (i) Describe the bonding in magnesium oxide and explain why it acts as an insulator.
- (ii) Suggest a reason why copper is suitable for encasing the magnesium oxide.

6 Intermediate Bond Types

Many covalent and ionic bonds have **intermediate bond character** – some covalent bonds have **ionic character** and some ionic bonds have **covalent character**



6.1 Ionic Character in Covalent Bond

Ionic character in a covalent bond arises due to sharing of electrons between atoms with a **considerable electronegativity difference**.

The **more electronegative** atom acquires a **partial negative charge** (denoted by δ^-), the **less electronegative** atom acquires a **partial positive charge** (denoted by δ^+), forming a **dipole moment** in a **polar** covalent bond

The extent of **ionic character** in a covalent bond **increases** with an **increase in electronegativity difference** between the covalently bonded atoms.


If we compare HBr and HI, both have simple molecular structures with weak permanent dipole-permanent dipole interactions between molecules.

As electronegativity of iodine is lower than that of bromine (as electronegativity decreases down a group), the electronegativity difference between the atoms in HI is smaller than that between HBr. Hence, HI has a smaller degree of ionic character.

6.2 Covalent Character in Ionic Bond

In a pure ionic bond, there is complete transfer of electrons and the electron clouds of both the cation and anion remain intact.

Covalent character in an ionic bond arises due to **polarisation**, which refers to the **distortion of anion electron cloud by a neighbouring cation** in an ionic compound. Due to this, there is some form of “electron sharing” between oppositely charged ions, leading to some covalent character being induced in the ionic bond



Fajans' rules can be used to predict the extent of polarisation, which depends on the **charge on the cation** and the **relative sizes of the cation and anion**:

Low ← Polarisation → High	
Low positive charge	High positive charge
Large cation	Small cation
Small anion	Large anion

The concept of **charge density** is useful for explaining the influence that an ion (usually a cation) has on the electrons in adjacent ions or molecules. The charge density of an ion can be calculated

by dividing its charge by its ionic radius, i.e. $\frac{q_+}{r_+}$ or $\frac{q_-}{r_-}$

where q_+ is the charge of the cation, q_- is the charge of the anion, r_+ is the cationic radius, r_- is the anionic radius.

The extent of polarisation (and hence attraction) of the electron cloud in adjacent anions depends on the charge density and, in turn, the **polarising power of the cation**.

- Small, highly charged ions (e.g. Al^{3+}) have **greater charge densities** and hence **higher polarising powers** than large, singly charged ions (e.g. Cs^+).
- If two ions have similar charge densities, their compounds often have similar properties (e.g. AlCl_3 and BeCl_2).

The extent of polarisation also depends on the **polarisability of the anion**. The **larger the anionic radius**, the **more polarisable** the anion is.

Hence, the extent of covalent character in an ionic bond is larger when:

- The cation has a high charge and a small radius i.e. high charge density (highly polarising).
- The anion has a large radius (highly polarisable).

The idea of ion polarisation is important in explaining the thermal decomposition of Group 2 carbonates under the topic of 'Periodic Table'.

Example 6A

In terms of structure and bonding, account for the following observation:

compound	melting point / °C
AlF_3	1090
AlCl_3	193

Understanding the context:

Due to the larger size of Cl^- ($r_{\text{Cl}^-} = 0.181 \text{ nm}$) compared to F^- ($r_{\text{F}^-} = 0.136 \text{ nm}$), the Cl^- ion is more polarisable than F^- , leading to substantial polarisation of the $\text{Al}-\text{Cl}$ bond in AlCl_3 .

Since Al^{3+} has high charge density and is highly polarising, this results in significant covalent character to the $\text{Al}-\text{Cl}$ bond. Hence AlCl_3 has a simple molecular structure while AlF_3 is essentially ionic.

AlF_3 has a giant ionic lattice structure while AlCl_3 has a simple molecular structure. More energy is required to break the stronger electrostatic forces of attraction between the oppositely charged ions in AlF_3 than the weaker instantaneous dipole-induced dipole interactions between AlCl_3 molecules. Thus, AlF_3 has a higher melting point than AlCl_3 .

7 Metallic Bonding


- LO (a) Show understanding that all chemical bonds are electrostatic in nature and describe:
- (ii) metallic bond as the electrostatic attraction between a lattice of positive ions and delocalised electrons.
 - (l) Describe, in simple terms, the lattice structure of a crystalline solid which is:
 - (v) metallic, as in copper

7.1 Definition of Metallic Bond

Metallic bonds are **strong electrostatic forces of attraction** between **metal cations** and **sea of delocalised electrons** in a **giant metallic lattice** structure

The metal atoms contribute their valence electrons to form cations in a **sea of delocalised electrons**. The **valence electrons** are **delocalised** due to the weak electrostatic forces of attraction between them and the nucleus.

In any given piece of metal, the **metallic lattice** is **electrically neutral** as the number of positive and negative charges balance out.



7.2 Factors affecting Metallic Bond Strength

The strength of metallic bonds is:

- **proportional to number of delocalised valence electrons** contributed per atom
- **proportional to the charge density of metal cation**, $\frac{q_+}{r_+}$

where q_+ is the charge of the cation, r_+ is the cationic radius.

In comparing two metals, the **difference in cationic charge tends to have a greater effect on metallic bond strength than the difference in cationic radius**.

The metallic bond strength is reflected in the boiling point of the metal.


E.g. Comparing the metallic bond strength of sodium, magnesium and aluminium

metal	sodium (Na)	magnesium (Mg)	aluminium (Al)
boiling point / °C	883	1091	2519
no. of delocalised valence electrons	1 (Group 1)	2 (Group 2)	3 (Group 13)
cationic charge, q_+	+1	+2	+3
cation radius, r_+ / nm	0.095	0.065	0.050
metallic bond strength	Na	< Mg	< Al

Note:

Transition metals tend to have particularly high melting and boiling points. The reason is that they can involve the **3d** electrons in the delocalisation as well as the **4s**. We will talk about this again in the topic of 'Chemistry of Transition Elements'.
















7.3 Physical Properties of Metals

Property / use	Reasons
High melting/boiling point	Large amount of energy is needed to overcome the strong electrostatic forces of attraction between metal cations and sea of delocalised electrons during melting / boiling.
Good conductors of electricity in solid and liquid states	Presence of sea of delocalised electrons as mobile charge carriers to conduct electricity under the influence of an electric current/field.
Good conductors of heat	The sea of delocalised electrons transfers the heat energy quickly throughout the metallic structure.
Malleable (can be beaten into sheets) Ductile (can be pulled out into wires)	Stress applied on a metallic lattice causes sliding of layers of cations over one another without causing the metallic structure to break down as the sea of delocalised electrons are holding the cations together to prevent repulsion between the cations . 
Alloy formation, e.g. brass (copper and zinc) steel (iron and carbon)	Presence of cations (metals)/atoms (non-metals) of different sizes in the metallic lattice inhibits sliding of layers , hence tensile strength of alloy is increased .

Checkpoint

- ☐ I am able to explain the physical properties observed in ionic compounds by relating to the structure and bonding of these substances.
- ☐ I understand the concept that covalent and ionic bonds can have intermediate bond character.
- ☐ I am able to compare the extent of covalent character between two ionic compounds based on the charge and size of the ions.
- ☐ I am able to explain the physical properties observed in metals by relating to the structure and bonding of these substances.




Chemical Bonds Summary

		ionic bonding	covalent bonding		metallic bonding
written description		<u>electrostatic force of attraction</u> between <u>oppositely charged ions</u>	<u>electrostatic force of attraction</u> between the <u>shared electrons</u> and <u>positively charged nuclei</u> between <u>2 atoms</u> . <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div style="text-align: center;"> ↓ <i>covalent</i> ↓ overlapping of 2 <i>half-filled orbitals</i> </div> <div style="text-align: center;"> ↓ <i>dative / coordinate</i> ↓ overlapping of 1 <i>filled</i> and 1 <i>empty orbital</i> </div> </div>		<u>electrostatic force of attraction</u> between <u>metal cation</u> and <u>sea of delocalised electrons</u> .
dot-and-cross diagram		e.g. KCl $[K]^{\oplus} [\overset{\cdot\cdot}{\underset{\cdot\cdot}{\text{Cl}}}]^{\ominus}$	e.g. BCl ₃ 	e.g. H ₃ O ⁺ 	
directionality		non-directional	directional		non-directional
structures		giant ionic lattice	simple molecular	giant covalent / giant molecular	giant metallic lattice
physical properties	m.p. / b.p.	high / non-volatile	low / volatile	high / non-volatile	high / non-volatile
	hardness	hard and brittle	soft	hard	hard but malleable & ductile
	electrical conductivity	conducts in liquid and aqueous states (<i>mobile ions</i>)	non-conductor	non-conductor (except those with layers e.g. graphite)	conducts in solid and liquid states (<i>mobile electrons</i>)
	solubility	<i>generally</i> soluble in polar solvents but insoluble in non-polar solvents	<i>generally</i> insoluble in polar solvents but soluble in non-polar solvents	<i>generally</i> insoluble in all solvents	<i>generally</i> insoluble in all solvents (Note that some metals can react with solvents e.g. H ₂ O)
pictorial representation	solid		 (This diagram is for diatomic molecules)	 e.g. C or Si	 • electron
	liquid				
	gas				


Intermolecular Forces of Attraction

(for simple molecular substances)


















(3 types)

		instantaneous dipole– induced dipole interactions <i>(present in <u>all</u> molecules)</i>	permanent dipole– permanent dipole interactions	hydrogen bonding
strength (similar M_r)		Usually very weak small M_r < Large M_r branched molecule < straight molecule.	weak	moderate
molecules involved		non-polar	polar <i>(atoms of different electronegativities and molecule with a net dipole moment)</i>	Molecules consist of H–F bonds H–O bonds H–N bonds and an electronegative atom with at least one lone pair of electrons, e.g. alcohols, RCOOH, amines, NH ₃ , H ₂ O
diagrammatic representation				
effects on physical properties	m.p./ b.p.	very low	low	moderate
	solubility	soluble in non-polar solvents <i>generally</i> insoluble in polar solvents	intermediate solubilities	<i>generally</i> soluble in water

Flowchart:




Shapes of Molecules

VALENCE-SHELL ELECTRON-PAIR REPULSION (VSEPR) THEORY					
electron pairs	arrangement	bond pairs	lone pairs	molecular shape	
2		2	0	linear e.g. BeCl_2 , CO_2	
3		3	0	trigonal planar e.g. BF_3 , SO_3	
		2	1	bent e.g. SO_2	
4		4	0	tetrahedral e.g. CH_4	
		3	1	trigonal pyramidal e.g. NH_3	
		2	2	bent e.g. H_2O	
5		5	0	trigonal bipyramidal e.g. PCl_5	
		4	1	see-saw e.g. SF_4	
		3	2	T-shaped e.g. ClF_3	
6		6	0	octahedral e.g. SF_6	
		5	1	square pyramidal e.g. IF_5 , SF_5^{2-}	
		4	2	square planar e.g. XeF_4 , ICl_4^-	

Strength of electron pair repulsion: lone pair-lone pair > lone pair-bond pair > bond pair-bond pair

Appendix 1: A Little Bit of History.....

In the early 20th century, the American chemist Gilbert N. Lewis began to use dots in lecture, while teaching undergraduates at Harvard, to represent the electrons around atoms. His students favoured these drawings, which stimulated him in this direction. From these lectures, Lewis noted that elements with a certain number of electrons seemed to have a special stability: it appeared that once a core of eight electrons has formed around a nucleus, the layer is filled, and a new layer is started. Lewis also noted that various ions with eight electrons also seemed to have a special stability. On these views, he proposed the rule of eight or octet rule: ions or atoms with a filled layer of eight electrons have a special stability.



Moreover, noting that a cube has eight corners Lewis envisioned an atom as having eight sides available for electrons, like the corner of a cube. Subsequently, in 1902 he devised a conception in which cubic atoms can bond on their sides to form cubic-structured molecules.

In 1913, while working as the chair of the department of chemistry at the University of California, Berkeley, Lewis read a preliminary outline of paper by an English graduate student, Alfred Lauck Parson, who was visiting Berkeley for a year. In this paper, Parson suggested that the electron is not merely an electric charge but is also a small magnet (or “magneton” as he called it) and furthermore that a chemical bond results from two electrons being shared between two atoms.

Moreover, he proposed that an atom tended to form an ion by gaining or losing the number of electrons needed to complete a cube. Thus, Lewis structures show each atom in the structure of the molecule using its chemical symbol. Lines are drawn between atoms that are bonded to one another; occasionally, pairs of dots are used instead of lines. Excess electrons that form lone pairs are represented as pair of dots, and are placed next to the atoms on which they reside.


To summarise his views on his new bonding model, Lewis states:

Two atoms may conform to the rule of eight, or the octet rule, not only by the transfer of electrons from one atom to another, but also by sharing one or more pairs of electrons...Two electrons thus coupled together, when lying between two atomic centres, and held jointly in the shells of the two atoms, I have considered to be the chemical bond. We thus have a concrete picture of that physical entity, that “hook and eye” which is part of the creed of the organic chemist.

The following year, in 1917, an unknown American undergraduate chemical engineer named Linus Pauling was learning the Dalton hook-and-eye bonding method at the Oregon Agricultural College, which was the vogue description of bonds between atoms at the time. Each atom had a certain number of hooks that allowed it to attach to other atoms, and a certain number of eyes that allowed other atoms to attach to it. A chemical bond resulted when a hook and eye connected. Pauling, however, wasn't satisfied with this archaic method and looked to the newly emerging field of quantum physics for a new method.

In 1927, the physicists Fritz London and Walter Heitler applied the new quantum mechanics to the deal with the saturable, non-dynamic forces of attraction and repulsion, *i.e.*, exchange forces, of the hydrogen molecule. Their valence bond treatment of this problem, in their joint paper, was a landmark in that it *brought chemistry under quantum mechanics*. Their work was an influence on Pauling, who had just received his doctorate and visited Heitler and London in Zürich on a Guggenheim Fellowship.

Subsequently, in 1931, building on the work of Heitler and London and on theories found in Lewis' famous article, Pauling published his ground-breaking article "The Nature of the Chemical Bond" in which he used quantum mechanics to calculate properties and structures of molecules, such as angles between bonds and rotation about bonds. On these concepts, Pauling developed hybridisation theory to account for bonds in molecules such as CH_4 , in which four sp^3 hybridised orbitals are overlapped by hydrogen's 1s orbital, yielding four sigma (σ) bonds.



(Taken from wikipedia : https://en.wikipedia.org/wiki/History_of_molecular_theory)


Why are We Studying This?

Molecular structure is the foundation on which chemistry, the study of matter and the changes it undergoes, rests. Much of chemistry is concerned with changes at the molecular level as structures are elucidated and chemical reactions occur.

Appendix 2: Structure of the Atom

Understanding the nature of the chemical bond relies on knowledge of the composition and structure of the atom. The model of the atom that emerged from the early work of Thomson, Rutherford, Moseley, and Bohr was of a central, very small, positively charged nucleus composed of positively charged protons and neutral neutrons, surrounded by one or more negatively charged electrons moving at high speed and effectively occupying a volume much larger than that of the nucleus.

The lack of reactivity of the noble gases – no known compounds at that time – led both Walther Kossel and Lewis to conclude in 1916 that these substances have a particularly stable arrangement of electrons. This in turn led to the development of the shell model of the atom. In the shell model, the electrons in an atom are arranged in successive spherical layers or shells surrounding the nucleus. The outer shell is never found to contain more than the number of electrons in the valence shell of a noble gas. The outer shell is called the valence shell because it is these electrons that are involved in bond formation and give the atom its valence.



Experimental support for the shell model has been provided by the determination of the ionisation energies of free atoms in the gas phase and by the analysis of the spectra of such atoms. These measurements have given a picture of the arrangement of the electrons in an atom in terms of their energies, which can also be deduced from the quantum mechanical description of an atom. Quantum mechanics also shows us that electrons do not have fixed positions in space but are in constant motion, following paths that cannot be determined. Hence, it is strictly not correct to talk about the *arrangement* of the electrons. It is their energy, not their positions, which can be determined.

On the basis of the shell model, two apparently different models of the chemical bond were proposed, the ionic model and the covalent model.

Appendix 3: Ionic Bonding – Prelude





In 1916, W. Kossel noted that the loss of an electron by an alkali metal gives a positive ion, such as Na^+ (2,8) or K^+ (2,8,8), where the numbers in parentheses represent the number of electrons in successive shells. Hence, these ions have the same electron arrangement as a noble gas. Similarly, the gain of an electron by a halogen gives a negative ion, such as a fluoride ion, F^- , (2,8) or a chloride ion, Cl^- , (2,8,8), also with the electron arrangement of a noble gas: that is, an outer shell containing eight electrons.

Kossel proposed that these ions are formed because their valence shell electrons have the same stable arrangements as a noble gas. He considered solid sodium chloride to consist of positive sodium ions (cations) and negative chloride ions (anions) held together in a regular pattern by electrostatic attraction. Each crystal of solid sodium chloride can be regarded as a single giant molecule, in which a very large number of ions are arranged in a regular manner that continues through the crystal.

Misconception

Misconception: Atoms form stable ions by losing or gaining one or more electron to have the same stable arrangements as a noble gas.

Correction: The formation of ions does not necessarily lead to greater stability. Metal atoms take in energy (ionisation energy) to free the electrons from attraction by the nucleus, which means that the resulting cations (and isolated electrons) are less stable than the atoms.




Evidence that solids such as NaCl do consist of ions was provided by the observation that these materials are conducting in the molten state and in solution such as water. In these states the ions are free to move independently of each other under the action of an applied electric field. Sodium chloride is a non-conductor in the solid state, because the ions are fixed in position.

Sodium chloride and many similar compounds are said to be ionic compounds held together by ionic bonds. However, even though the term “ionic bond” is widely used, it is a vague and ill-defined concept. Electrostatic forces act in all directions and through relatively long distances so that the *attractive forces are not confined to just two neighbouring oppositely charge ions*. Moreover, there are also *repulsive forces between ions of like charge*.


Appendix 4: Covalent Bonding – Prelude

Clearly the explanation of the chemical bond given by Kossel cannot apply to homonuclear molecules (*i.e.* molecules comprising of the same type of atoms) such as Cl_2 . Almost simultaneously with the publication of Kossel's theory, G. N. Lewis published a theory that could account for such molecules. Like Kossel, Lewis was impressed with the lack of reactivity of the noble gases. But he was also impressed by the observation that the vast majority of molecules have an even number of electrons, which led him to suggest that in molecules, electrons are usually present in pairs. In particular, he proposed that in a molecule such as Cl_2 the two atoms are held together by sharing a pair of electrons because in this way each atom can obtain a noble gas electron arrangement, as in the following examples:



Diagrams of this type are called Lewis diagrams or Lewis structures. The bond between the two atoms could be called a shared-electron-pair bond but it is now universally called a covalent bond – a term introduced by Irving Langmuir (1919). In a Lewis diagram, the pairs of electrons that are not forming bonds are called nonbonding pairs or, more usually, lone pairs.

Although Lewis had no clear idea of why electrons are found in molecules as pairs, or how a shared pair of electrons holds two atoms together, the ideas of the shared electron pair, the covalent bond and the octet rule enable us to understand the formulas of a vast number of molecules and their relationship to the positions of the elements in the periodic table.



Because the formation of electron pairs seemed to contradict Coulomb's law, according to which electrons repel each other so that they should keep as far apart as possible, Lewis even suggested that Coulomb's law is not obeyed over the very short distances between electrons in atoms and molecules. Although we now know that Coulomb's law is obeyed for all distances between charges, in making the assumption about the importance of electron pairs, Lewis displayed remarkable *intuition*: electrons do indeed form pairs in most molecules, despite their mutual electrostatic repulsion. Nonetheless Lewis diagrams showing bonding pairs and lone pairs are still widely used today, and the electron pair remains a central concept in chemistry.

Lewis called the apparent tendency of atoms to acquire a noble gas electron arrangement, either by forming ions or by sharing electron pairs, the rule of eight. Later Langmuir called it the octet rule, and this is the term that is now generally used. Lewis did not regard the rule of eight as being as important as the rule of two, according to which electrons are present in molecules in pairs, because he found more exceptions to the octet rule than to the rule of two. There are only a few exceptions to the rule of two, such as molecules with an odd number of electrons (free radicals), whereas there are a large number of exceptions to the octet rule.

Appendix 5: Coordination Number

- ☼ Coordination number, with regards to an ionic lattice, refers to the number of oppositely charged ions that surrounds the ion of concern.

The factors that affect coordination number are:

- Relative size of the ions

For any given ionic compound,

- Typical coordination numbers to be considered are 4, 6 and 8.
- There are 3 possible scenarios as shown below:

scenarios	consequence	coordination number
one ion is very much smaller in size than the other	region of space around the small ion is limited, hence reducing the number of oppositely charged ions around it	4
one ion is slightly smaller in size than the other	intermediate between the other two scenarios.	6
both cations and anions are similar in size	region of space around both ions are optimised, hence maximising the number of oppositely charged ions around it.	8

- Relative charge







To achieve electrical neutrality,

- Cation with +2 charge will need twice as many anions with –1 charge, when compared to a cation with +1 charge

compound	cationic charge	cationic radius/ nm	anionic charge	anionic radius/ nm	$\frac{\text{anionic radius}}{\text{cationic radius}}$	coordination number of cation	coordination number of anion
ZnS	+2	0.08	–2	0.19	2.4	4	4
NaCl	+1	0.10	–1	0.18	1.8	6	6
MgO	+2	0.07	–2	0.14	2.0	6	6
CsCl	+1	0.17	–1	0.18	1.1	8	8
CaF ₂	+2	0.10	–1	0.13	1.3	8	4

Self Check Answers:

Self Check 1B

(a) BF_3 (b) Cl_2O (c) SO_3 (d) O_3 (e) H_2F^+ (f) AlH_4^- (g) ICl_4^- 

Self Check 3A

Question 1:

Answer: D

The millions of hairs and thousands of pads on each hair would lead to significant instantaneous dipole – induced dipole interactions.

Coordinate bonds, covalent bonds and ionic bonds are very strong and thus it would be too difficult to break (not feasible for the gecko).

Question 2:

Answer: C

Both H_2O and HF have intermolecular hydrogen bonds. As the formation of hydrogen bonds requires a lone pair on F, O or N, **AND** a H atom bonded to F, O or N, H_2O is able to form two hydrogen bonds per molecule while HF can only form one (limited by the number of H atoms).

Self Check 4A

Question 1:

Answer: A

“Very high melting point” – suggests giant structure

“hard” – suggests giant molecular structure (NOT layered) or giant ionic structure

The properties suggest silicon carbide has a giant molecular structure similar to that of diamond. Hence, it has covalent bonds between silicon and carbon atoms.

Question 2:

Answer: B

- 1 True: Only 3 electrons of the C atoms in graphite are involved in bonding. The last electron is delocalised.
- 2 True: The C atoms in graphite form 3 bonds with neighbouring C atoms.
- 3 False: Valency is the number of extra electrons that the atom will need in order to attain a stable structure.

Self Check 5B

- (i) In MgO , ionic bonding occurs due to **electrostatic forces of attraction between oppositely charged Mg^{2+} and O^{2-}** . At room temperature, the **ions** can only **vibrate about fixed positions** and hence are **not mobile** to conduct electricity.
- (ii) MgO has **very high melting point** and so would not melt in a fire.
Unlike PVC, MgO **upon heating does not give off poisonous fumes** such as HCl .
- (iii) Copper is **ductile and malleable** and hence can be bent into shape and pulled into tube to encase MgO .