

PIONEER JUNIOR COLLEGE

JC2 PRELIMINARY EXAMINATION HIGHER 2

CANDIDATE NAME			
CT GROUP	I 2	INDEX NUMBER	
CHEMISTRY			9647/02
Paper 2 Structured C	Questions		18 September 2013
Candidates answer of	on the Question Paper.		2 hours
Additional Materials:	Data Booklet		

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name on all work you hand in. Write in dark blue or black pen on both sides of the paper. You may use a soft pencil for any diagrams, graphs or rough workings. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

FOR EXAMINER'S USE						
Paper 2						
1	/ 12	5		/5		
2	/ 10	6		/ 7		
3	/ 14	Penalty	sf	units		
4	/ 24	Total		/ 72		

This document consists of **20** printed pages.

1 Planning

Heat is absorbed when potassium iodide dissolves in water. Sucrose, on the other hand, does not produce a temperature change when added to water.

964

By studying the temperature changes when known masses of potassium iodide dissolve, the percentage by mass of potassium iodide in a mixture of potassium iodide and sucrose, **FA 5** can be determined.

You are provided with

- a test tube of FA 1 which contains 2.00 g of potassium iodide
- a test tube of **FA 2** which contains 4.00 g of potassium iodide
- a test tube of **FA 3** which contains 6.00 g of potassium iodide
- a test tube of **FA 4** which contains 8.00 g of potassium iodide
- an empty test tube labelled FA 5
- **FA 5**, a mixture of potassium iodide and sucrose
- a thermometer
- a polystyrene cup
- other common apparatus in the laboratory
- (a) The percentage composition of potassium iodide in the mixture, **FA5**, is estimated to be 50%. Suggest a suitable value for mass of **FA 5** used.

.....[1]

(b) Describe, in a series of numbered steps, how you would measure the temperature changes by dissolving separate samples of FA 1 to FA 5 in water. Hence, determine the mass of potassium iodide in a mixture of FA 5.

Your plan should include the following:

- the apparatus for measurements
- how you would take and record the various measurements
- the tables for recording the data measured
- the graph to be plotted

(c) Sketch the graph you suggested in (b). Show clearly how the percentage by mass of potassium iodide in **FA 5** can be determined.

[2]

(d) A second student conducted a similar experiment using the same apparatus and the same volume of water as in your plan in (b) but using twice the mass of FA 1 to FA 5. Suggest whether your experiment or the second student's experiment would result in a more reliable determination of the percentage by mass of potassium iodide in FA 5.

[2]

(e) (i) A third student measured a temperature change of 3 °C when he dissolved 12 g of potassium iodide in 100 cm³ of water. Assume that the specific heat capacity of the final solution is 4.18 J g⁻¹ K⁻¹, and that its density is 1.00 g cm⁻³, calculate the enthalpy change of solution of potassium iodide.

(ii) The published value for the enthalpy change of solution of potassium iodide is +20.3 kJ mol⁻¹. Suggest an explanation for the difference between the published value and the value calculated in (i).

[2]

[Total: 12]

2 In a typical iodine clock reaction, solutions of hydrogen peroxide, sulfuric acid, potassium iodide, starch and sodium thiosulfate are mixed. There are two reactions occurring in the reaction mixture.

6

In the first reaction, iodide ions are oxidised slowly by acidified hydrogen peroxide to form iodine.

$$H_2O_2(aq) + 2I^-(aq) + 2H^+(aq) \rightarrow I_2(aq) + 2H_2O(l)$$
 reaction I

In the second reaction, iodine is reduced back to iodide ions as they are generated.

 $2S_2O_3^{2-}(aq) + I_2(aq) \rightarrow S_4O_6^{2-}(aq) + 2I^{-}(aq) \qquad \mbox{reaction II}$

When all the sodium thiosulfate present has reacted, iodine will appear in the solution producing an immediate deep blue colour with starch indicator.

A series of experiments was carried out using different volumes of the five reagents, each solution being made up to the same total volume with water. The following results were obtained.

expt	volume of	volume	time for the				
no.	0.10	0.100	0.100	0.050	starch	of	appearance
	mol dm⁻³	mol dm⁻³	mol dm⁻³	mol dm⁻³	solution	water	of deep
	H_2O_2	KI	HC/	$Na_2S_2O_3$	/ cm ³	/ cm ³	blue colour
	/ cm ³	/ cm ³	/ cm ³	/ cm ³			/ s
1	18	12	6	2	3	59	23
2	6	12	12	2	3	65	70
3	6	18	12	2	3	59	47
4	12	12	12	2	3	59	35

(a) Calculate the relative initial rate of reaction I for each of these experiments.

[1]

(b) Given that the order of reaction with respect to hydrogen peroxide is found to be one, use the results in (a) to deduce the order of reaction with respect to I⁻ and H⁺ in reaction I. Explain your reasoning. (c) Hence write an overall rate equation for reaction I, stating the units of the rate constant.

[1]

(d) Suggest, with reasons, which of the following two mechanisms, **A** or **B**, fits the observed kinetics data.

Mechanism A	
$\overline{2H^+(aq) + 2I^-(aq)} \rightarrow 2HI (aq)$	(fast)
$2HI(aq) + H_2O_2(aq) \rightarrow I_2(aq) + 2H_2O(l)$	(slow)
Mechanism B	
$H_2O_2(aq) + I^-(aq) \rightarrow H_2O(l) + OI^-(aq)$	(slow)
$H^+(aq) + OI^-(aq) \rightarrow HIO(aq)$	(fast)
$HIO(aq) + I^{-}(aq) + H^{+}(aq) \rightarrow I_{2}(aq) + H_{2}O(l)$	(fast)
	••••••
	[2]

- 8
- (e) With the aid of a sketch of the Boltzmann distribution, explain how an increase in temperature increases the rate of the reaction.

 • • • •	 • • • •	• • • •	•••	•••	•••	•••	•••	•••	•••	•••	• • •	•••	•••	•••	•••	•••	•••	• • •	•••	•••	•••	••	•••	•••	•••	•••	•••	•••	•••	••	• • •	•••	•••	• • •	•
 	 ••••						•••					• • •	•••	•••							• • •	•••			•••					•••			•••	•••	•
 	 						•••						•••	•••											•••					•••			•••		•
 	 ••••						•••					•••	•••	•••					•••						•••					•••			•••	•••	•
 	 •••	••••										• • •	•••	•••					• • •		• • •	•••			•••					••			•••	•••	•
 	 ••••						•••					• • •	•••	•••					• • •			••			•••					••			•••	 [4	-]
																														[Го	ota	l: ⁻	10]

9
(a) The following diagram shows how the d-orbitals of iron in [Fe(H₂O)₆]²⁺ complex are split in an octahedral environment.



The arrangement of electrons in the d-orbitals depends on the *spin states* of complexes.

In a 'high spin' state, the electrons occupy all the d-orbitals singly, before starting to pair up in the lower energy d-orbitals.

In a 'low spin' state, the lower energy d-orbitals are filled first, by pairing if necessary, before the higher energy d-orbitals are used.

When the H₂O ligands in $[Fe(H_2O)_6]^{2+}$ are changed to CN^- ligands, the Fe²⁺ ion changes its electronic configuration from a 'high spin' state to a 'low spin' state.

(i) Briefly explain why the spin state changes to low spin, when the ligands coordinated to Fe^{2+} are changed from H₂O to CN⁻.

(ii) Use diagrams to show the electronic distribution in the d-orbitals in the complexes $[Fe(H_2O)_6]^{2+}$ and $[Fe(CN)_6]^{4-}$ respectively.

(iii) Many transition elements and their compounds are paramagnetic, which means that they are attracted to a magnetic field. This property is due to the presence of **unpaired electron(s)** in the d–orbitals.

The table below compares the relative paramagnetism of $[V(H_2O)_6]^{2+}$, $[Fe(H_2O)_6]^{2+}$ and $[Fe(CN)_6]^{4-}$.

972

formula of complex	relative paramagnetism
$[V(H_2O)_6]^{2+}$	3
[Fe(H ₂ O) ₆] ²⁺	4
[Fe(CN) ₆] ^{4–}	x

With reference to (ii), explain the relative paramagnetism for $[V(H_2O)_6]^{2+}$ and $[Fe(H_2O)_6]^{2+}$. Hence, suggest a value of **x** in the table above, giving your reasoning.

[6]

(b) A student was given a solution containing iron(II) and zinc(II) ions and was tasked to separate the two cations such that each cation is present in its **solid** state. He proposed and carried out the following reactions.

To the solution in a test-tube, NaOH(aq) was added in excess. A green precipitate was obtained. Filtering the mixture gave a green residue and a colourless filtrate. When $HNO_3(aq)$ was added to the colourless filtrate, a white precipitate was obtained.

(i) Suggest the identity of the green precipitate and state the formula of the compound in the colourless filtrate.

identity of green precipitate:

formula of the compound in the colourless filtrate:

- 11
- (ii) Give the identity of the white precipitate. With the aid of an appropriate equation, explain why the addition of $HNO_3(aq)$ to the filtrate causes the white precipitate to form.

[4]

(c) Anodising is the process to increase the thickness of the oxide layer on the surface of metal artefacts so as to increase the resistance of the metal to corrosion and to allow the application of dyes.

An aluminium mobile phone casing with a surface area of 200 cm² is to be anodised. Given that the density of Al_2O_3 is 4.0 g cm⁻³, calculate the total quantity (in coulombs) of electricity that will be required to increase the thickness of the oxide layer on the casing by 0.2 mm.

974

(a) On heating, a solid sample of phosphorus pentachloride is completely vaporised in a 2 dm³ vessel. At 227 °C, the following equilibrium is set up:

 $PCl_5(g) \Rightarrow PCl_3(g) + Cl_2(g)$

The partial pressure of Cl_2 at equilibrium is found to be 2.00 atm. Given that the equilibrium constant, K_p , is 1.01 atm,

(i) write an expression for the equilibrium constant, K_{p} , for this reaction,

.....

(ii) calculate the total pressure of the system at equilibrium,

(iii) calculate the mass of the solid PCl_5 placed in the vessel initially.

- (b) Phosphorus pentachloride is a useful chlorinating agent for organic synthesis. Using phosphorus pentachloride in one of the steps, propose synthetic routes starting from propan-1-ol to the following compounds.
 - (i) propanamide

(ii) butanoic acid

- (c) For **each** of the following pairs of compounds, describe a chemical test which would enable you to distinguish between them. State clearly the observations for each compound.
 - (i) $CH_3CH_2CONH_2$ and $CH_3CH_2CO_2^-NH_4^+$



(e) Glutamine, H₂NCH(CH₂CH₂CONH₂)CO₂H, is a non-essential amino acid as it can be synthesised in the body. The intake of glutamine supplements can help athletes to recover at fast speed from excess of physical stress.

The pH change when 1.0 mol dm^{-3} of NaOH is added to 25 cm³ of a 1.0 mol dm^{-3} solution of **protonated** form of glutamine is shown below.

977



(i) State the pK_1 and pK_2 values of glutamine.

(ii) Use the structure of glutamine, and its pK_1 and pK_2 values to suggest the species present at points **G** and **J** in the graph above.

(iii) Write ionic equations to show how the solution at point **H** is able to maintain an almost constant pH upon addition of a small amount of

Ι	H⁺(aq) ions	
II	OH⁻(aq) ions	
		[5]
	[Total:	24]

[2]

[1]

5 *Terfenadine* is an antihistamine which was originally used for the treatment of allergies. However, it was withdrawn from the market in 1997 due to its side effect of affecting the heart's rhythm which may result in cardiac arrest.

The structure of the drug is given below.



- (a) Draw the structural formulae of the **organic products** when *Terfenadine* undergoes each of the following reactions.
 - (i) heated over aluminium oxide.

(ii) heated under reflux with hot acidified dichromate(VI) ions.

(b) Name the type of reaction that occurs in the reaction in (a)(i).

.....

(c) State the type of isomerism exhibited by *Terfenadine,* and give one further piece of relevant information about it.

[2]

[Total: 5]

6 Compound **C** has the molecular formula $C_8H_6O_4$. It is an aromatic compound which contains three functional groups.

19

Data	about the	reactions	of C	are	given	in	the table	Э.
------	-----------	-----------	------	-----	-------	----	-----------	----

reaction	reagent	result
1	neutral FeCl ₃ (aq)	violet complex formed
2	Br ₂ (aq) in an excess	Br ₂ is decolourised; white solid formed with $M_r = 323.8$
3	2,4-dinitrophenylhydrazine	orange precipitate formed
4	PCl ₅	white fumes and a neutral cyclic compound D formed with $M_{\rm r} = 148$
5	Na	colourless gas evolved. When 0.1 mol of C is reacted with excess sodium metal, 2.4 dm ³ hydrogen is formed, measured at room temperature and pressure.

- (a) Name the functional group that reaction 1 shows to be present in C. [1]
 (b) Name the functional group that reaction 3 shows to be present in C. [1]
- (c) Based on reaction 5 and your answer in (a) and (b), deduce the identity of the third functional group present in C.

[2]

[1]

(d) Deduce the molecular formula of the product formed in reaction 2.

(e) Draw the fully displayed structures of **C** and **D**.

[2]

[Total: 7]

End of Paper