

YISHUN JUNIOR COLLEGE

JC2 PRELIMINARY EXAMINATION 2014

CHEMISTRY

9647/02

HIGHER 2

Paper 2 Structured Questions

2 hours

Candidates answer on the Question Paper.

Additional Materials: Data Booklet



READ THESE INSTRUCTIONS FIRST

Write your name and CTG on all the work that you hand in.

Write in dark blue or black pen on both sides of the paper.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.

A Data Booklet is provided.


For Examiner's Use	
Paper 1	
Total	/ 40
Paper 2	
1	/ 12
2	/ 23
3	/ 9
4	/ 10
5	/ 18
Total	/ 72
Paper 3	
Total	/ 80
Overall	/ 192

Answer **all** the questions.

Planning (P)

- 1** Fermentation is a natural process. Man applied fermentation to make products such as wine, mead, cheese and beer long before the biochemical process was understood. In the 1850s and 1860s, Louis Pasteur became the first scientist to be known to have studied fermentation when he demonstrated fermentation was caused by living cells.

The fermentation of carbohydrates into alcohol can be represented as:



This reaction is catalyzed by yeast enzymes called zymases. A balanced chemical equation for this process using table sugar or sucrose is:



This type of yeast fermentation can be studied through its CO_2 production. Using the information above, you are required to write a plan to determine the rate of CO_2 production in a 30-minute fermentation process.

You are provided with the following materials:

- 0.100 mol dm⁻³ sucrose solution
- yeast suspension
- apparatus normally found in a school laboratory

- (a)** Write a step-by-step plan on how you would carry out this experiment.

Your plan should include the following:

- show how the quantity of sucrose used is calculated
- all essential experimental details
- a diagram of your experimental set up
- a table with appropriate headings to show the data you would record when carrying out your experiment

.....

.....

.....

.....

.....


.....

.....

[6]

Alcohol produced by this fermentation is concentrated or enriched by distillation.

A quantity of reaction mixture, after the fermentation process, is placed in the following set-up.



(b) What is meant by distillation?

.....
 [1]

(c) Why is this simple distillation suitable for the fermentation mixture? Give **two** reasons.

.....

 [2]

(d) What is a disadvantage of this type of distillation?

.....
 [1]

- (e) Alcohol content in a distillate is often determined by measuring its density, which depends heavily on the percentage of alcohol. A measured density is compared with tabulated values of the density of known mixtures of alcohol and water to determine its alcohol content; typically given as a volume percentage.

Most manufacturers of liquor report the alcohol content by its Proof. The Proof is double the volume percentage alcohol:

$$\text{Proof} = 2 \times \text{volume \%}$$

You are provided with the data on a distillate obtained from the distillation process:

1. Mass of sample = 97.0 g
2. Volume of sample = 100.0 cm³
3. Data of alcohol % volume versus density (g/cm³).

Use these data below to determine the Proof of this sample.

% by Volume	Density [g/cm³]
10.0	0.98569
15.0	0.98024
20.0	0.97518
25.0	0.97008
30.0	0.96452
35.0	0.95821
40.0	0.95097
45.0	0.94277
50.0	0.93350

[2]


[Total : 12]

- 2** “Hard water” is water that has high mineral content such as calcium ions. In domestic settings, hard water is often indicated by a lack of suds formation when soap is agitated in water.

A typical sample of ‘hard water’ has a concentration of calcium ions of $2.50 \times 10^{-4} \text{ mol dm}^{-3}$.

- (a)** In order for a detergent to be used in ‘hard water’, sodium tripolyphosphate, $\text{Na}_5\text{P}_3\text{O}_{10}$, is added as a water softening agent.

The sodium tripolyphosphate ‘softens’ water by complexing with calcium ions. The complexation reaction is as follows:



For this reaction with calcium ions, the equilibrium constant is $7.7 \times 10^8 \text{ mol}^{-1} \text{ dm}^3$

- (i)** Write the K_c expression for the reaction.
- (ii)** Hence, calculate the concentration of tripolyphosphate ion required to reduce the calcium ion concentration in a typical sample of ‘hard water’ to $1.0 \times 10^{-6} \text{ mol dm}^{-3}$.

(b) 'Hard water' also contains magnesium ions which can form a precipitate with the detergent. For example, magnesium ions form magnesium iodide, MgI_2 , in the presence of potassium iodide.

(i) The lattice energy of MgI_2 is $-2327 \text{ kJ mol}^{-1}$ while the values of the enthalpy change of hydration are listed below:

Ions	$\Delta H_{\text{hyd}} / \text{kJ mol}^{-1}$
Mg^{2+}	-1920
I^-	-295

Calculate the enthalpy change of solution of magnesium iodide.

(ii) Using your answer from (b)(i) and the fact that entropy change of solution of magnesium iodide is positive, predict whether magnesium iodide is soluble in water at room temperature. Give your reasoning.

.....

.....

.....

.....

[4]

- (c) Construct a Born-Haber cycle in the grid below to calculate the enthalpy change of formation of MgCl_3 . Use the relevant data given below as well as from the *Data Booklet*.

Enthalpy change of atomisation of Mg	+146 kJ mol^{-1}
1 st electron affinity of Cl	-364 kJ mol^{-1}
ΔH_f (MgCl_2)	-652 kJ mol^{-1}
Lattice energy of $\text{MgCl}_3(\text{s})$	-5010 kJ mol^{-1}

- (i) Energy / kJ mol^{-1}



- (ii) Comparing ΔH_f (MgCl_2) with your answer in (c)(i), comment on the relative stabilities of MgCl_2 and MgCl_3 .

.....

.....

[4]

- (d) Calcium is a fairly soft, silvery-grey metal which quickly tarnishes in air. Vanadium is a hard grey metal which is resistant to corrosion at ordinary temperatures. Metallic calcium has no commercial uses; vanadium is widely used as an alloying element in steels.

Data about calcium and vanadium are given below:

	Calcium	Vanadium
Electronic configuration	[Ar] 4s ²	[Ar] 3d ³ 4s ²
Atomic radius / nm	0.197	0.122
Melting point / °C	843	1710
Density / g cm ⁻³	1.54	6.07
Oxidation states in aqueous solution	+2	+2, +3, +4, +5

- (i) Although the vanadium atom has more electrons than the calcium atom, the atomic radius of vanadium is smaller than that of calcium. Suggest an explanation for this.

.....

.....

.....

.....

- (ii) The melting point of vanadium is significantly higher than that of calcium. Account for the difference in their melting points.

.....

.....

.....

.....

.....

- (iii) Compounds of calcium contain Ca^{2+} while the corresponding compounds of vanadium contain V^{3+} . Use the *Data Booklet* to explain in thermodynamic terms why Ca^{3+} compounds do not exist and V^{3+} compounds do.

.....

.....

.....

.....

.....


[8]

- (e) Fluorspar is an important mineral composed of calcium and fluorine. It is used in a wide variety of chemical; metallurgical and ceramic processes.

Fluorspar, CaF_2 , is an ionic compound. Draw a 'dot-cross' diagram to show the chemical bonding in fluorspar.

[1]

- (f) Fluorspar reacts with concentrated sulfuric acid to form hydrogen fluoride gas.



Data about HF, HCl, HBr and HI are given below.

	HF	HCl	HBr	HI
Boiling point / °C	+20	−85	−67	−35
Bond energy / kJ mol ^{−1}	562	431	366	299
ΔH_f^θ / kJ mol ^{−1}	−269	−92	−36	+26

- (i) Explain why the boiling points of HCl, HBr and HI increases down the group.

.....

.....

.....

- (ii) Suggest why the boiling point of HF is much higher than those of the other three.

.....

.....

.....


.....

[3]


[Total: 23]

- 3** Being the most reactive of all elements, fluorine is commonly found in organic compounds.

Carbonyl fluoride, COF_2 (structure as shown below) exists as a toxic gas which reacts rapidly with water to evolve two different acidic gases.



- (a)(i)** The reaction of carbonyl fluoride with water is found to proceed via the formation of an intermediate as shown:



State the reaction that has occurred to produce the above intermediate.

.....

- (ii)** The above intermediate is found to dissociate rapidly to give two different acidic gases. One of these gases is carbon dioxide. State the identity of the other gas.

.....

- (iii)** Write a balanced equation of the reaction between carbonyl fluoride and water.

.....

[3]

- (b)** An unknown amount of carbonyl fluoride was reacted with water (in excess). The two acidic gases released are then reacted with NaOH (aq). 25.00 cm^3 of $0.100 \text{ mol dm}^{-3}$ of NaOH (aq) was required for complete neutralisation.

- (i)** Calculate the number of moles of NaOH used.

(ii) Calculate the number of moles of CO_2 produced.

(iii) Using your answer in **(b)(ii)**, calculate the volume of carbon dioxide in m^3 produced under the conditions of 30°C and 1 atm (assuming ideal gas behavior).

(iv) Calculate the mass of carbonyl fluoride used.

[6]

[Total: 9]

- 4 The acidity of hydrogen-containing compounds varies remarkably from one compound to another.

The South African plant *Dichapetalum cymosum* contains fluoroethanoic acid, FCH_2COOH . Data about ethanoic acid, fluoroethanoic acid and chloroethanoic acid are given below:

	$\text{CH}_3\text{CO}_2\text{H}$	$\text{FCH}_2\text{CO}_2\text{H}$	$\text{ClCH}_2\text{CO}_2\text{H}$
$\text{p}K_{\text{a}}$	4.76	2.57	2.87

- (a)(i) Define the term $\text{p}K_{\text{a}}$.

.....

- (ii) Calculate K_{a} for fluoroethanoic acid.

[2]

- (b)(i) Comparing the three acids given above, state the strongest acid and the weakest acid.

Strongest acid:

Weakest acid:

- (ii) Suggest an explanation for the differences in acid strengths of the three acids in terms of their structure and bonding.

.....


.....

.....

.....

[4]

- (c) Procaine is a local anaesthetic drug used primarily to reduce the pain of intramuscular injection of penicillin.



Procaine

There are two basic functional groups in the procaine molecule, with pK_b values of 4.19 and 9.37.

- (i) Name the two basic functional groups present in the procaine molecule.

.....


- (ii) Match the given pK_b values to the corresponding functional groups.

pK_b	Functional Group
4.19	
9.37	

[4]

[Total: 10]

- 5 Dengue fever and dengue haemorrhagic fever are the most common mosquito-borne viral disease in the world. There was no drug to treat dengue until 2014. According to The Straits Times' report dated 12 July 2014, the first safe and effective dengue vaccine in the world could be out commercially by July 2015. Before the announcement on the vaccine, cinnamon oil made the most effective natural mosquito repellent known to man. And, each of the following compounds, which are present in cinnamon, appears to be effective as pesticide.



The group $-\text{OCH}_3$ which is present in compounds **C** and **D** can be regarded as inert.

- (a) Only **one** of the compounds **A**, **B**, **C** or **D** will react with **each** of the following reagents. In each case, identify the compound concerned and draw the structural formula of the organic product formed.

Each compound may be used once, more than once, or not at all.

reagent	compound A, B, C or D	structural formula of the organic product(s)
CH_3COCl		
Hot dilute H_2SO_4		

reagent	compound A, B, C or D	structural formula of the organic product(s)
2,4- dinitrophenylhydrazine		

[7]

(b) Compound **A** may be converted into compound **B** in a two-stage process.

(i) What is the structural formula of the intermediate in this conversion?

(ii) For each stage, in the reaction sequence, give reagents and conditions.

Stage I: Reagents




Conditions.....

Stage II: Reagents

Conditions.....

[4]

Another similar aromatic flavouring agent is vanillin. Vanillin is a phenolic aldehyde with the molecular formula $C_8H_8O_3$:



Eugenol was used in the production of isoeugenol for the manufacture of vanillin.

- (c) What type of isomerism do eugenol and isoeugenol exhibit?

.....
[1]

- (d) Isoeugenol exists as a pair of stereoisomers. Draw the structures of the isomers and state the type of isomerism shown.

Type of isomerism:

[2]

- (e) 16.4 g of eugenol yield 13.0 g of vanillin. Calculate the percentage yield of vanillin.

[2]

- (f) Name a reaction in which these three compounds (vanillin, eugenol and isoeugenol) react similarly, and give the structure of the major organic product formed from eugenol.

Reaction:

Structure of major organic product:

[2]

[Total: 18]

END OF PAPER