



ST ANDREW'S JUNIOR COLLEGE  
PRELIMINARY EXAMINATIONS  
HIGHER 1

CANDIDATE  
NAME

--

CLASS

2	3			
---	---	--	--	--

**H1 CHEMISTRY**

**Paper 2 Structured Questions**

**8873 / 02**

**27 August 2024**  
**2 hours**

Candidates answer on the Question Paper.  
Additional Materials: Data Booklet

**READ THESE INSTRUCTIONS FIRST**

Write your centre number, index number, name and class at the top of this page.

Write in dark blue or black pen.

You may use a HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** questions in the **spaces provided** on the Question Paper.

A Data Booklet is provided.

**Section A**

Answer **all** the questions.

**Section B**

Answer **one** question.

Question Number	Total Marks	Marks Obtained
SECTION A		
1	14	
2	7	
3	9	
4	30	
SECTION B		
5	20	
6	20	
TOTAL	80	

The use of an approved scientific calculator is expected, where appropriate.

The number of marks is given in brackets [ ] at the end of each question or part question.

## Section A

Answer **all** the questions in this section in the spaces provided.

1 This is a question about Group 14 elements and its compounds.

(a) (i) A sample of silicon contains three isotopes, as shown in Table 1.1. [2]

Table 1.1

Isotope	Relative Isotopic mass	Percentage abundance. %
$^{28}\text{Si}$	27.976	92.22
$^{29}\text{Si}$	28.976	4.69
$^{30}\text{Si}$	29.973	3.09

Calculate the relative atomic mass of silicon in this sample, giving your answer to **two decimal places**.

(ii) Complete the electronic configuration for silicon. [1]

**1s<sup>2</sup>** .....

(iii) Explain why second ionisation energy of silicon is lower than that of aluminium. [2]

.....  
 .....  
 .....  
 .....

[TURN OVER]

- 1 (b) When carbon tetrachloride,  $\text{CCl}_4$ , and silicon tetrachloride,  $\text{SiCl}_4$ , were added to water, the following observations were seen.

carbon tetrachloride	Forms two immiscible layers
silicon tetrachloride	White solid and steamy fumes

- (i) Write a balanced equation for the reaction of  $\text{SiCl}_4$  with water. [1]

.....

- (ii)  $\text{CCl}_4$  has no reaction with water. Suggest an explanation for the inertness of  $\text{CCl}_4$  to water. [1]

.....

.....

- (iii) Explain why  $\text{CCl}_4$  forms two immiscible layers with water when they are mixed. [2]

.....

.....

.....

.....

- 1 (c) Due to carbon's ability to form strong multiple bonds, it can form different allotropes like graphene and fullerene.

Graphene is an allotrope of carbon that occurs as two-dimensional sheets while fullerenes are molecules of carbon atoms with hollow shapes. An example of a spherical fullerene is buckminsterfullerene,  $C_{60}$ , while carbon nanotube (CNT) is a type of cylindrical fullerene.

Buckminsterfullerene has low melting point and is slippery while CNT, like graphene, has high tensile strength and is an excellent conductor of electricity.

- (i) Graphene is a *nanomaterial*.

Define what is meant by the term *nanomaterial*.

[1]

.....

.....

.....

- (ii) Draw a diagram to illustrate the structure of graphene.

[1]

[TURN OVER]

- 1 (c) (iii) Explain how carbon nanotube is an excellent conductor of electricity and has high tensile strength. [2]

.....  
.....

- (iv) Suggest why buckminsterfullerene is a good lubricant. [1]

.....  
.....

[Total: 14]

[TURN OVER

- 2 Propanone,  $\text{CH}_3\text{COCH}_3$ , reacts with iodine,  $\text{I}_2$ , in the presence of an acid catalyst.

An experiment is performed using  $1.00 \text{ mol dm}^{-3}$  of  $\text{I}_2$  and  $1.00 \text{ mol dm}^{-3} \text{H}^+(\text{aq})$  where concentration of  $\text{CH}_3\text{COCH}_3$  is monitored over time. The same experiment was then repeated using  $0.500 \text{ mol dm}^{-3} \text{H}^+(\text{aq})$  instead.

The results obtained are used to plot the graph below.




Fig 2.1

- 2 (a) It was found that the reaction is zero order with respect to  $[I_2]$ .

Define the term *order of reaction*.

[1]

.....  
.....

- (b) (i) Use the graph, Fig 2.1, to determine the order of reaction with respect to  $[CH_3COCH_3]$  and  $[H^+]$  respectively.

[2]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

- (ii) Write the rate equation for this reaction. Hence, calculate the value of rate constant,  $k$ . Include the units.

[2]

.....  
.....

[TURN OVER]

- 2 (c) Other halogens, like chlorine and bromine, can also react with propanone in similar reactions as iodine.

Describe and explain how the reactivity of halogens varies down the group. Include relevant data from the *Data Booklet* in your answer. [2]

.....

.....

.....

.....

.....

.....

.....

.....

[Total: 7]

[TURN OVER]



3 A sample of sodium nitrite,  $\text{NaNO}_2$ , was analysed by titrating with yellow aqueous cerium(IV) ions,  $\text{Ce}^{4+}$ . The experimental procedures for this analysis were as follows.

- 0.140 g of impure  $\text{NaNO}_2$  was dissolved in deionised water to form a colourless solution. It was then transferred to a  $100 \text{ cm}^3$  volumetric flask and top up to the mark with more deionised water.
- $25.0 \text{ cm}^3$  of this solution required  $22.05 \text{ cm}^3$  of  $0.0400 \text{ mol dm}^{-3} \text{ Ce}^{4+}(\text{aq})$  to reach the end-point. During the titration,  $\text{Ce}^{4+}$  was reduced to colourless  $\text{Ce}^{3+}$  while nitrite ion was oxidised to form colourless nitrate ion,  $\text{NO}_3^-$ .

(a) (i) Draw a dot-and-cross diagram of  $\text{NaNO}_2$ . [2]

(ii) Construct the ionic equation for the reaction between sodium nitrite and cerium(IV) ions. [1]

.....

(iii) Calculate the percentage purity of this sample of  $\text{NaNO}_2$ . [3]

[TURN OVER]

- 3 (a) (iv) Predict the colour change at the end-point of this titration. [1]

.....





- (b) Suggest and explain the difference in the ionic radii of  $\text{Ce}^{4+}$  and  $\text{Ce}^{3+}$ . [2]

.....  
.....  
.....  
.....  
.....  
.....  
.....

[Total: 9]

[TURN OVER

4 Diol is an organic compound with 2 –OH groups. The following are some examples of diol.

Compound **A**Compound **B**Compound **C**Compound **D**

(a) (i) Compounds **A** to **D** are constitutional isomers.

Define the term *constitutional isomers*.

[1]

.....  
 .....  
 .....

(ii) Give the systematic name of compound **A**.

[1]

.....

(iii) Both compound **A** and butane are made up of four carbon atoms. Despite this, butane is less reactive than **A**.

Explain briefly why this is so.

[1]

.....  
 .....  
 .....

[TURN OVER]

- 4 (a) (iv) **Table 4.1** shows the boiling points of compounds **A** to **D**.

**Table 4.1**

Compound	Boiling point / °C
<b>A</b>	210
<b>B</b>	192
<b>C</b>	230
<b>D</b>	176

With reference to structure and bonding, explain why compound **C** has a higher boiling point than compound **B**.

[2]

.....

.....

.....

.....

- (v) With reference to (a)(iv), explain why compound **D** has the lowest boiling point.

[2]

.....

.....

.....

.....

[TURN OVER

- 4 (b) (i) In the spaces provided below, draw the structure of the organic product obtained when compound **D** is reacted with the following reagents and conditions. State the type of reaction for each reagent and condition.

I)  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{H}_2\text{SO}_4$ , heat

II) Concentrated  $\text{H}_2\text{SO}_4$ ,  $170^\circ\text{C}$

III) excess  $\text{CH}_3\text{COOH}$ , concentrated  $\text{H}_2\text{SO}_4$ , heat

[6]

Reagent and condition	I	II	III
Structure of organic product			
Type of Reaction			

- (ii) With reference to (b)(i), state the role of concentrated  $\text{H}_2\text{SO}_4$  in reaction III.

[1]

.....

- (iii) Compound **B** can be synthesised from a bromoalkane.


Draw the **displayed** formula of this bromoalkane. State the reagents and conditions for this synthesis.

[2]

Reagents and conditions: .....

[TURN OVER]

- 4 (c) When 2 –OH groups are bonded to the same carbon atom, the compound is known as geminal diol. Geminal diol is unstable and will spontaneously lose water to form a carbonyl compound. An equation to represent this is as shown.




- (i) Explain what is meant by a *reversible reaction*. [1]

.....  
 .....  
 .....

- (ii) For the conversion between geminal diol and carbonyl compound, suggest whether the ratio of  $\frac{[\text{products}]}{[\text{reactants}]}$  at equilibrium will be less than, equal to or greater than 1. [1]

.....


- (iii) For the following compound, draw the structure of the carbonyl compound that it will form.




[1]

[TURN OVER

- 4 (d) Diol can react with diisocyanate to form polyurethane, which is a polymer. The equation representing this reaction is as shown.



Spandex is a fabric that contains polyurethane formed from compound **E** and  $\text{HOCH}_2\text{CH}_2\text{OH}$ .



Compound **E**

Unlike nylon which is derived from coal and petroleum, all the constituent parts of polyurethane in spandex are synthesised in the laboratory. Spandex is non-biodegradable. When polyurethane in spandex is heated, a variety of harmful products can be produced, such as carbon monoxide,  $\text{CO}$ , and hydrogen cyanide,  $\text{HCN}$ .

The properties of spandex and nylon are shown in **Table 4.2**.

**Table 4.2**

Property	Spandex	Nylon
Stretchability	High	Moderate
Durability	High	High
Retention	High	Low to moderate
Breathability	High	High

- (i) State the molecular formula of **E**.

[1]

.....

[TURN OVER

- 4 (d) (ii) Spandex fibres are strong due to the intermolecular forces between the polymer chains.

State the two types of intermolecular forces responsible for this property and the group(s) of atoms involved. [3]

.....

.....

.....

.....

- (iii) Explain why spandex fabric dissolves slowly when placed in aqueous alkali. [1]

.....

.....

.....

- (iv) Deduce whether spandex is classified as a thermoplastic or thermoset. Explain your answer. Hence, state whether it can be recycled. [2]

.....

.....

.....

.....



- 4 (d) (v) Suggest a preferred method to dispose spandex. State a disadvantage associated with this method. [1]

.....

.....

.....

.....

- (vi) In high performance sports, such as running, strenuous movements are involved and therefore, a large amount of sweat is produced. It is important for the sportsperson to feel comfortable when wearing the sportswear.

Between spandex and nylon, state which polymer will be more suitable to make the sportswear for high performance sports. Explain your choice by considering the properties listed in **Table 4.2**. [3]

.....

.....

.....

.....

.....

.....

[Total: 30]

[TURN OVER]

**Section B**

Answer one question from this section in the spaces provided.

- 5 (a) New car exhaust systems are fitted with a catalytic converter where it converts harmful gases like carbon monoxide and nitrogen dioxide to harmless gases like carbon dioxide and nitrogen before the exhaust gases are being released into the atmosphere.

- (i) Write an equation to represent the chemical reaction taking place in the catalytic converter as described in (a). [1]

.....

- (ii) State the type of catalyst involved and explain the mode of action of this type of catalyst. [4]

.....  
.....  
.....  
.....  
.....  
.....  
.....

**[TURN OVER]**

- 5 (a) (iii) Explain, with the aid of a Boltzmann distribution diagram, on how this catalyst affects the rate of reaction.

[3]

.....


.....

.....

.....

.....

- (b) An industrial process in the production of a fuel,  $\text{N}_2\text{O}_3$  involved the following equilibrium system.



In one of such processes, 4 mol of dinitrogen tetroxide,  $\text{N}_2\text{O}_4$ , and 9 mol of nitrogen monoxide,  $\text{NO}$ , was heated in a  $3.0 \text{ dm}^3$  vessel at  $450^\circ\text{C}$ . The two gases reacted slowly to produce  $\text{N}_2\text{O}_3$ . The amount of  $\text{N}_2\text{O}_3$  at equilibrium was found to be 4.8 mol.

- (i) Write an expression for the equilibrium constant,  $K_c$  for the reaction, stating its units.

[2]

[TURN OVER]

5 (b) (ii) Determine the equilibrium concentration of the gases and hence the value of  $K_c$ . [3]

(iii) Define the term *dynamic equilibrium*. [1]

.....  
.....

(iv) Suggest and explain the temperature and pressure conditions which could help to increase the yield of  $N_2O_3$ . [4]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

[TURN OVER

5 (b) (v) The actual industrial temperature and pressure conditions were as followed:

- Temperature: 500 °C
- Pressure: 150 atm

Explain why these conditions were applied instead of what was suggested in

(b)(iv).

[2]

.....

.....

.....

.....

.....

.....

[Total: 20]

[TURN OVER

- 6 In this current environmental landscape, it is important to search for alternative energy sources. Methanol and methane have been investigated to serve as alternative sources of fuels for motor cars.

(a) (i) Write an equation to represent the standard enthalpy change of combustion of methanol. [1]


.....

(ii) Using relevant data from the *Data Booklet*, calculate another value for the standard enthalpy change of combustion of methanol. [2]

[TURN OVER

- 6 (a) (iii) *Use of Data Booklet is relevant to this question.*

The standard enthalpy change of combustion of methanol can be determined in the laboratory using the apparatus shown in the diagram below.



The temperature of water rose from 25 °C to 90 °C when all the methanol was completely burnt. The density of methanol is 0.792 g cm<sup>-3</sup> and the standard enthalpy change of combustion of methanol is -715 kJ mol<sup>-1</sup>.

Calculate the percentage efficiency of this heating process.

[4]

- (iv) Suggest why there was a discrepancy between your answer in (a) (ii) and that quoted in (a) (ii).

[1]

.....

.....

[TURN OVER

- 6 (b) Methanoic acid, commonly known as formic acid, is used as a food preservative. It can be formed from oxidising methanol.

(i) Write a balanced equation for the formation of methanoic acid from methanol. You can use [O] to represent the oxidising agent.

[1]

.....


(ii) Using methanoic acid as an example, explain what is meant by an acid – conjugate base pair.

[1]

.....

.....

- (c) 25.0 cm<sup>3</sup> of 0.100 mol dm<sup>-3</sup> methanoic acid, HCOOH, was titrated with 0.100 mol dm<sup>-3</sup> NaOH. The titration curve is shown below.



[TURN OVER]



6 (c) (i) Justify, with relevant calculations, that methanoic acid is a weak acid. [2]

(ii) Write a balanced chemical equation for the reaction between methanoic acid and sodium hydroxide. [1]

.....

(iii) Determine the value of X. [1]

(iv) Explain, with the aid of a chemical equation to explain why pH is greater than 7 [2]  
when X cm<sup>3</sup> of NaOH was added.

.....

.....

.....

(v) Suggest a suitable indicator for this titration and explain your choice. [2]

.....

.....

.....

[TURN OVER]

- 6 (c) (vi) It was found that a buffer mixture was obtained when  $\frac{x}{2}$  cm<sup>3</sup> of NaOH was added. Write equations to show how the mixture resisted changes in pH when small amounts of H<sup>+</sup> and OH<sup>-</sup> was added to a separate solution of this buffer mixture. [2]

.....

.....

.....

[Total: 20]

END OF PAPER

[TURN OVER



