Class:	Register No:	Name:



CRESCENT GIRLS' SCHOOL SECONDARY FOUR PRELIMINARY EXAMINATION 2024

CHEMISTRY

Paper 1 Multiple Choice 28 August 024
1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, highlighters, glue or correction fluids.

Write your name, index number and class on the Answer Sheet in the spaces provided.

DO NOT WRITE ON ANY BARCODES.

There are **forty** questions on this paper. Answer **all** questions. For each question, there are four possible answers, **A**, **B**, **C** and **D**.

Choose the one you consider correct and record your choice in soft pencil on the OTAS sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet

A copy of the Periodic Table is printed on page 19.


The use of an approved scientific calculator is expected, where appropriate.

This booklet consists of 19 printed pages, including the cover page.

1 A student measures the rate of the reaction between marble chips, CaCO₃, and dilute hydrochloric acid.

$$CaCO_3 + 2HCI \rightarrow CaCl_2 + CO_2 + H_2O$$

Which diagrams show the apparatus that are suitable for this experiment?



marble chips


- **A** 1 and 2
- **B** 1, 2 and 4

marble chips

® € 30


- **C** 2 and 3
- **D** 2, 3 and 4

2 Chromatograms of a urine sample using two different solvents are shown below.



Based on the two chromatograms, which drug(s) is/are present in the urine sample?

- Α drug **P** only
- В drug **Q** only
- C drugs P and Q only
- D drugs P, Q and R
- 3 Since the discovery of graphite, scientists have been able to extract a single layer of carbon atoms (known as graphene) and convert it to another material known as graphane by attaching one hydrogen atom to each carbon atom as shown below.



Which property of graphene is **not** likely to be shared by graphane?

- Α It is insoluble in water.
- В It is very strong.
- С It has a high melting point.
- D It is an electrical conductor.

- An isotope of element **Z** has 20 neutrons and 17 protons. Which is the correct symbol for an ion of the isotope of element **Z**?
 - A $\frac{18}{17}$ Z +


B $\frac{18}{17}$ Z -

 $C = \frac{37}{17} Z$

- D $\frac{37}{18}$ Z -
- 5 Hydrogen is able to form compounds with metals and non-metals. The formulae of some of these compounds are shown below.
 - CH₄
- HCl
- MgH_2
- KH

What is the order of melting point of these compounds?


- lowest melting point highest melting point Α CH₄ HCl KΗ MgH_2 В CH₄ HCl MgH_2 ΚH C HCl CH_4 MgH_2 KΗ D KΗ MgH_2 HClCH₄
- 6 The heating curves (not drawn to scale) of nitrogen and oxygen over a period of time are shown in the graph.



At which temperature will there be two **different** states of matter co-existing at the same time, in a mixture of nitrogen and oxygen under similar conditions?

- **A** 180 °C
- **B** 200 °C
- **C** 215 °C
- **D** 220 °C

7 The diagram below shows the bonding between $\bf P$ and $\bf Q$ in the covalent molecule, $\bf PQ_2$.



What are the electronic structures of atoms ${\bf P}$ and ${\bf Q}$ before combining together to form the above molecule?

	Р	Q
Α	2.8	2.8.8
В	2.6	2.8.7
С	2.6	2.8.6
D	2.4	2.8.7

8 0.1 mole of a chloride XCl_2 combines with 10.8 g of water to form the hydrated salt, $XCl_2.nH_2O$.

What is the value of **n**?


9 Ammonia reacts with chlorine according to the equation shown below:

$$2NH_3(g) + 3Cl_2(g) \rightarrow N_2(g) + 6HCl(g)$$

If 90 cm 3 of ammonia is mixed with 60 cm 3 of Cl_2 and all the volumes were measured at room temperature and pressure, what is the total volume of gases at the end of the reaction?

- **A** 20 cm³
- **B** 120 cm³
- **C** 140 cm³
- **D** 190 cm³

When 4.8 g of magnesium is heated in a crucible, 5.9 g of magnesium oxide is formed.



What is the percentage yield of magnesium oxide?

- **A** 53%
- **B** 74%
- **C** 80%
- **D** 81%

A student is given two samples, one of which is aluminium oxide and the other is magnesium carbonate. He needs to find a method to identify the two samples.

Which of the following show(s) the correct method(s) and observation(s)?

	method	observation(s)
1	add nitric acid	only aluminium oxide dissolves
2	add nitric acid	both samples dissolve. Effervescence is observed in the reaction with magnesium carbonate
3	add sodium hydroxide	only aluminium oxide dissolves
4	add sodium hydroxide	both samples dissolve. Effervescence is observed in both the reactions

- A 1 and 4 only
- B 2 only
- C 2 and 3 only
- **D** 3 only

12 Butterfly pea flower extract is commonly used in drinks nowadays and it changes colour according to different pH values.

The table below shows the colours of butterfly pea flower extract at different pH values.

pH range	colour
0 – 3	violet
4 – 8	blue
9 – 11	green
12 – 14	yellow

Which pair of substances can be distinguished by adding butterfly pea flower extract to each substance separately?

- A acid rain and aqueous sodium chloride
- B aqueous ammonia and limewater
- **C** aqueous sodium sulfate and aqueous sodium chloride
- **D** dilute hydrochloric acid and dilute sulfuric acid
- Which reaction will produce the least volume of carbon dioxide?
 - A sodium carbonate and hydrochloric acid
 - B copper(II) carbonate and hydrochloric acid
 - C magnesium carbonate and sulfuric acid
 - **D** lead(II) carbonate and sulfuric acid
- 14 The table below shows the results of some tests carried out on separate portions of a solution **M**.

test	observation
aqueous sodium hydroxide added	test-tube feels warm and no precipitate forms
acidified aqueous silver nitrate added	white precipitate forms


What could be the identity of solution M?

- A hydrochloric acid
- B potassium sulfate
- C sodium chloride
- **D** zinc sulfate

A student stated that since low temperatures produce a greater yield of ammonia, the reaction should be carried out at 50 °C instead of 450 °C.

Which statement best explains why the reaction is not carried out at 50 °C?

- A Ammonia is unstable at 50 °C.
- **B** The reactants are unstable at 50 °C.
- **C** The reaction is too slow at 50 °C.
- **D** The reaction mixture is easily separated at higher temperatures.
- Ammonium chloride is heated strongly in a boiling tube. Damp blue and red litmus papers were placed at the mouth of the boiling tube for the gases produced.



Which row shows the correct sequence of observations that would be made?

	first observed colour change	final colour of both litmus papers
Α	The damp blue litmus paper turns red.	red
В	The damp blue litmus paper turns red then bleaches.	white
С	The damp red litmus paper turns blue.	blue
D	The damp red litmus paper turns blue.	red


- 17 Which are redox reactions?
 - 1 $HCl + NaOH \rightarrow NaCl + H_2O$
 - 2 $Zn + 2HNO_3 \rightarrow Zn(NO_3)_2 + H_2$
 - 3 Ag₂SO₄ + 2NaC $l \rightarrow 2$ AgCl + Na₂SO₄
 - 4 $2Fe^{2+} + Cl_2 \rightarrow 2Fe^{3+} + 2Cl^{-}$
 - **A** 1, 2 and 3

B 1 and 3

C 2 and 4

D 3 and 4


Dry hydrogen gas is passed over a heated brown powdered solid and then through a cooled U-tube before the excess of hydrogen is burned in air.



A colourless liquid collects in the U-tube. What could the brown powdered solid be?

- A aluminium oxide
- B copper(II) oxide
- c iron(III) oxide
- D magnesium oxide
- 19 Equal masses of different metals 1 to 4 are placed in the test tubes containing an equal volume of hydrochloric acid of equal concentration.


The thermometers show the maximum temperature recorded for the reaction. (The room temperature is $25\,^{\circ}\text{C.}$)



Which statements are most likely to be true?

- I Metal 3 will displace metals 2 and 4 from their aqueous salt solutions.
- II Metal 2 can likely be extracted by chemical reduction of its oxide by carbon.
- III Metal 1 is likely to be obtained by electrolysing its molten chloride.
- **A** I, and II only
- B I and III only
- C II and III only
- **D** I, II and III

The energy profile diagram represents a chemical reaction carried out with a catalyst and without a catalyst.




- What is the enthalpy change for the catalysed reaction?
- **A** 125 kJ/mol
- **B** 50 kJ/mol
- **C** + 75 kJ/mol
- **D** + 100 kJ/mol
- 21 Hydrogen peroxide reacts with potassium iodide in the presence of dilute acid to produce iodine molecules as shown in the equation below.

$$H_2O_2$$
 (aq) + $2I^-$ (aq) + $2H^+$ (aq) $\rightarrow I_2$ (aq) + $2H_2O$ (l)

- Which factor would **not** affect the rate of this reaction?
- A Concentration of hydrogen peroxide
- **B** Concentration of potassium iodide
- C Pressure of the reacting vessel
- **D** Temperature of the reacting vessel and its surroundings

In the graph shown, curve 1 was obtained by the decomposition of 100 cm³ of 1.0 mol/dm³ hydrogen peroxide solution with manganese(IV) oxide as the catalyst. The equation for this reaction is shown.


$$2H_2O_2 \ \rightarrow \ 2H_2O + O_2$$




Which change to the original experimental conditions would produce curve 2?

- A adding some 0.1 mol/dm³ hydrogen peroxide solution
- **B** lowering the temperature
- **C** using a different catalyst
- **D** using less manganese(IV) oxide
- 23 In which set-up will the bulb light up?







В



C



D



24 During an electrolysis, X⁺ and Y⁻ ions are selectively discharged as shown in the equations below:

Cathode :
$$2X^+ + 2e^- \rightarrow X_2$$

Anode :
$$2Y^- \rightarrow Y_2 + 2e^-$$

What can the electrolyte be?

- A aqueous magnesium chloride
- B aqueous sodium sulfate
- C concentrated magnesium chloride
- D molten potassium chloride
- In which electrolysis experiment would there be no change in the concentration of the solution?

	<u>electrodes</u>	<u>electrolyte</u>
Α	carbon	aqueous copper(II) sulfate
В	copper	aqueous copper(II) sulfate
С	carbon	concentrated potassium chloride
D	platinum	dilute sulfuric acid

Methane reacts very slowly with air at room temperature. However, if a transition metal T is added to the methane-air mixture, the methane ignites quickly.

A student made some statements about the observation.

- I Addition of T reduces the activation energy.
- II Addition of T increases the enthalpy change.
- III Addition of T increases the rate of reaction.
- IV Addition of T reduces the energy of the reactants.

Which statements are correct?

- A I and II only
- **B** I and III only
- C II and III only
- **D** All of the above


27 The positions of the elements W, X, Y and Z are shown in part of the periodic table.

										22.2	
										z	
w									Υ	** · · · *****************************	
	Х		Π		ALL AND ALL AN						

Which statement is **not** correct?

- A All the elements are reactive except for element Z.
- **B** Element W and element Y can form ionic bonds.
- **C** Element X will react with element Z in the ratio 1:2.
- **D** Element Y and element Z will form a compound by sharing electrons.
- Which statement is **not** true when chlorine gas is bubbled into potassium iodide solution?
 - A Chlorine is more reactive than iodine and hence displaces iodine form potassium iodide solution.
 - **B** Potassium iodide is the reducing agent.
 - **C** The ionic equation for the reaction is $Cl_2(g) + 2I(aq) \rightarrow 2Cl(aq) + I_2(aq)$
 - **D** The solution turns from brown to colourless.
- 29 Bioethanol can be obtained from the fermentation of the sugar in sugarcane. Which statement best explains why burning of bioethanol is considered more environmentally sustainable compared to the use of fossil fuels?
 - A sugarcane grows, it absorbs carbon dioxide produced during photosynthesis.
 - **B** Cabon dioxide and water are formed during burning of bioethanol.
 - **C** Sugarcane plants can be regrown and replaced within a short period of time.
 - **D** Sugarcane plants need to be planted and transported for treatment.

30 Figure below shows the fractional distillation of petroleum.



Which statement best describes the fractions at X and Y?

- A The molecules in fraction X contain more carbon atoms than the molecules in fraction Y.
- **B** The molecules in fraction X are more flammable than the molecules in fraction Y.
- **C** The molecules in fraction X are larger than the molecules in fraction Y.
- **D** The molecules in fraction X have higher boiling points than the molecules in fraction Y.
- 31 An unsaturated hydrocarbon, C_4H_6 reacts with 0.10 mole of hydrogen gas to form the corresponding alkane.

What is the mass of C₄H₆ that is required to react with the hydrogen gas completely?

- **A** 0.90 g
- **B** 1.80 g
- **C** 2.70 g
- **D** 3.60 g
- Which one of the following shows the correct structural formula and name of the ester formed when methanoic acid reacts with propanol?

	structural formula	<u>name</u>
Α	CH ₃ CH ₂ COOCH ₃	methyl propanoate
В	CH ₃ CH ₂ COOCH ₃	propyl methanoate
С	HCOOCH ₂ CH ₂ CH ₃	methyl propanoate
D	HCOOCH ₂ CH ₂ CH ₂	propyl methanoate

33 Which structures are isomers?

A I, II and IV

- B I, II and V
- **C** I, III and IV

- **D** II, III and V
- 34 The diagram below shows an organic compound, cysteine.

Which statement about cysteine is true?

- A Effervescence is observed when magnesium metal is added to cysteine.
- **B** It decolourises acidified potassium manganate(VII).
- **C** It forms a polymer with the same linkage as terylene.
- **D** It forms an addition polymer with other units of cysteine.

An organic compound M undergoes a 2-stage process to form a compound N of chemical formula: CH₃CH₂COOH.

The reagents and conditions of the 2 reactions are as follows:

stage number	reagent(s)	conditions
1	steam	300 °C, 65 atm,
		Phosphoric(V) acid
2	acidified potassium manganate(VII)	heat

Which can be a possible identity of compound M?

- A butane
- **B** butene
- **C** propane
- **D** propene
- **36** Which statement is true about addition polymers and condensation polymers?
 - A Addition polymers are formed from alkenes while condensation polymers are formed from alkanes.
 - **B** Addition polymers produce water as a by-product whereas condensation polymers do not produce any by-products.
 - **C** Condensation polymers could produce water as a by-product whereas addition polymers do not produce any by-product.
 - **D** Nylon is an example of an addition polymer where terylene is an example of a condensation polymer.

37 Kevlar is a polymer with high tensile strength, which is five times greater than steel. It is a lightweight and strong fibre with many applications ranging from being used in bulletproof vests to tires. It has the structure below.

Which could be the monomer(s) for Kevlar?

A
$$H_2N$$
 CO_2H


$$c$$
 H_2N and HO_2C CO_2H

To reduce atmospheric pollution, the waste gases from a coal-burning power station are passed through powdered calcium carbonate.

Which waste gas will **not** be removed by the calcium carbonate?

- A carbon dioxide
- B nitrogen monoxide
- C phosphorus(V) oxide
- **D** sulfur dioxide

- **39** Which statements are always true of methane and carbon dioxide?
 - 1 Both gases can be produced by cattle.
 - 2 Both gases cause acid rain.
 - 3 Methane burns in limited oxygen to produce carbon dioxide.
 - 4 They are both greenhouse gases.
 - A 1 and 2 only
 - **B** 1 and 4 only
 - C 2 and 3 only
 - **D** 3 and 4 only
- The figure below shows the reaction of zinc in air. When all the grey solid has turned yellow, the source of heat was removed. Upon cooling, the yellow solid turned white.



During the reaction, a sample of 250 cm³ of air was used.

What is volume of the remaining air left after the experiment?

A 52.5 cm³

B 105 cm³

C 197.5 cm³

D 395 cm³

The Periodic Table of Elements

	18	2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton	84	24	×	xenon 131	98	~	radon	1	118	ő	oganesson	ı
	17			6	ш	fluorine 10	17	C	chlorine 35.5	35	ፙ	bromine	200	23	ч	iodine 127	82	¥	astatine	1	11/	<u>s</u>	tennessine	1
	16			8	0	oxygen 16	16	S	sulfur 32	34	Se	selenium	6/	25	<u>e</u>	tellurium 128	84	S.	polonium	1 5	116	_	Iivermorium	1
	15			7	z	nitrogen 1.4	15	۵	phosphorus 31	33	As	arsenic	ç/	51	Sp	antimony 122	83	ä	bismuth	507	115	Mc	moscovium	ı
	14			9	ပ	carbon 12	14	Si	silicon 28	32	Ge	germanium	/3	20	S	119 119	82	Pp	lead 2007	707	114	Εl	flerovium	1
	13			5	В	boron 11	13	Αl	aluminium 27	31	Ga	gallium	0/	49	п	indium 115	81	11	thallium	204	113	욷	nihonium	1
									12	30	Zu	zinc	ÇQ	48	ၓ	cadmium 112	80	Hg	mercury	107	112	ပ်	copernicium	ı
									£	59	ಪ	copper	64	47	Ag	silver 108	62	Αn	gold 40.7	181	111	ß	roentgenium	ı
Group									10	28	Z	nickel	60	46	Pd	palladium 106	8/	₫	platinum	CAL	110	Ds	darmstadtium	1
ž									6	27	ပိ	cobalt	60	45	돈	rhodium 103	22	1	mipin	761	109	Ħ	meitnerium	ı
		- I	hydrogen 1						80	56	Fe	io C	20	44	쮼	ruthenium 101	9/	ő	osmium	130	108	£	hassium	ı
							_		7	52	M	manganese	çç	43	ပု	technetium I	75	æ	rhenium	180	107	듐	bohrium	ı
				number	pol	mass			9	24	ර්	chromium	75	42	№	molybdenum 96	74	≥	tungsten	184	106	Sg	seaborgium	ı
			Key	proton (atomic) number	atomic symbol	name relative atomic mass			2	23	>	vanadium	LG!	4	g	niobium 93	73	<u>a</u>	tantalum	181	105	<u>ප</u>	dubnium	ı
				proton	ato	relati			4	22	j	titanium	48	40	ZŁ	zirconium 91	72	Ξ	hafnium	8/1	104	፟ጅ	rutherfordium	ı
				_					3	21	တိ	scandium	42	39	>	yttrium 89	57-71	lanthanoids		00,	89-103	actinoids		
	2			4	Be	benyllium Q	12	Mg	magnesium 24	20	ပ္ပ	calcium	40	38	Š	strontium 88	26	Ba	barium 407	13/	88	Ra	radium	1
	1			3	=	lithium 7	=	Na	sodium 23	19	×	potassium	33	37	윤	nubidium 85	55	ర	caesium	133	8/	<u></u>	francium	ı

_				_			
71	3	Intetium	175	103	ځ	lawrencium	ļ
20	χp	ytterbium	173	102	å	nobelium	1
69	T	thulium	169	101	Md	mendelevium	1
89	ш	erbium	167	100	Fm	fermium	1
29	운	holmium	165	66	Es	einsteinium	1
99	ò	dysprosium	163	86	ರ	californium	ı
65	Д	terbium	159	26	쓢	berkelium	ı
64	g	gadolinium	157	96	S	curium	ļ
63	Ш	europium	152	92	Am	americium	1
62	Sm	samarium	150	94	Pu	plutonium	ı
	Pm	_					
9	P	neodymium	144	92	_	uranium	238
	ፈ	praseodymium	141	91	Pa	protactinium	231
28	ဝီ	cerium	140	90	드	thorium	232
25	Гa	lanthanum	139	89	Ac	actinium	ı
	lanthanoids			actinoids			

The volume of one mole of any gas is $24 \, dm^3$ at room temperature and pressure (r.t.p.). The Avogadro constant, $L = 6.02 \times 10^{23} \, \text{mol}^{-1}$.

19