



**PU3 H2 Mathematics Paper 1 2012 Prelim II**

1(i)	$\sqrt{(4p)^2 + (7p)^2 + (-4p)^2} = 1$ $p^2 = \frac{1}{81}$ $p = \frac{1}{9} \quad (\text{reject } -ve)$
1(ii)	Length of projection of a on b.
1(iii)	$a \times b = \begin{pmatrix} 3 \\ -2 \\ 6 \end{pmatrix} \times \begin{pmatrix} \frac{4}{9} \\ \frac{7}{9} \\ -\frac{4}{9} \end{pmatrix} = \begin{pmatrix} -\frac{34}{9} \\ 4 \\ \frac{29}{9} \end{pmatrix}$
2	$\frac{2x^2 + 2x + 1}{x^2 - 2x + 1} \geq 1$ $\frac{2x^2 + 2x + 1}{(x-1)^2} \geq 1$ $x^2 + 4x \geq 0$ $x \leq -4 \text{ and } x \geq 0$ $x \neq 1$ $\frac{2(\ln x)^2 + 2(\ln x) + 1}{(\ln x)^2 - 2(\ln x) + 1} \geq 1$ $\ln x \leq -4, \ln x \geq 0$ $x \leq e^{-4}, x \geq 1$ $x \neq 1 \Rightarrow x \neq e$
3(i)	$a - b + c = 6$ $1.21a + 1.1b + c = 0.12$ $2.25a + 1.5b + c = 1$ $a = 2, b = -3, c = 1$
3(ii)	 <p><math>f(x)</math> is increasing for <math>x \geq 0.75</math>.</p>

3(iii)	
4	$(4)^{\frac{1}{2}} \left(1 + \frac{x}{4}\right)^{\frac{1}{2}} = 2 \left[ 1 + \frac{1}{2} \left(\frac{x}{4}\right) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!} \left(\frac{x}{4}\right)^2 + \dots \right]$ $= 2 + \frac{x}{4} - \frac{x^2}{64} + \dots$ $\left \frac{x}{4}\right  < 1 \Rightarrow  x  < 4$ $x = \frac{1}{2}$ $\sqrt{\frac{1}{2} + 4} = \sqrt{\frac{9}{2}} = \frac{3}{\sqrt{2}}$ $\frac{3}{\sqrt{2}} = 2 + \frac{\frac{1}{2}}{4} - \frac{\left(\frac{1}{2}\right)^2}{64} + \dots$ $\frac{3}{\sqrt{2}} \approx \frac{543}{256}$ $2\sqrt{2} = \frac{512}{181}$
5(a)	$a = 2000$ $r = 1.02$ $d = -50$ $T_{n+1} = 2000(1.02)^n - 50[1.02^{n-1} + 1.02^{n-2} + \dots + 1]$ $= 2000(1.02)^n - 50 \left( \frac{1 - 1.02^n}{1 - 1.02} \right)$ $= 2000(1.02)^n + 2500(1 - 1.02^n)$ $= 2000(1.02)^n - 500(1.02^n)$ $= 500(5 - 1.02^n)$ $500(5 - 1.02^n) \leq 0$ $\ln 5 \leq n \ln 1.02$ $n \geq 81.27$ <p>It can last 81 years.</p> $500(5 - 1.02^{30}) = \$1594.32$

5(b)	$a + 8d = 50$ $\frac{15}{2}(2a + 14d) = 570$ $a = -46, \quad d = 12$ $S_n > 500$ $\frac{n}{2}[-92 + (n-1)12] > 500$ $12n^2 - 104n - 1000 > 0$ $n < -9.59 \text{ (rej)} \text{ or } n > 9.59$ $\therefore \text{least } n \text{ is } 10.$
6(i)	$x^2 - y^2 = 2xy - 1$ $2x - 2y\left(\frac{dy}{dx}\right) = 2\left[y + x\left(\frac{dy}{dx}\right)\right]$ $2x - 2y = 2x\left(\frac{dy}{dx}\right) + 2y\left(\frac{dy}{dx}\right)$ $\frac{dy}{dx} = \frac{2x - 2y}{2x + 2y}$ $\frac{dy}{dx} = \frac{x - y}{x + y}$
6(ii)	$\frac{x - y}{x + y} = 0$ $x = y$ $x^2 - x^2 = 2x^2 - 1$ $x^2 = \frac{1}{2}$ $x = \pm\sqrt{\frac{1}{2}}$ $x = \frac{1}{\sqrt{2}}, \quad y = \frac{1}{\sqrt{2}}$ $x = -\frac{1}{\sqrt{2}}, \quad y = -\frac{1}{\sqrt{2}}$
7	$\frac{dy}{dx} = \frac{1}{2}(x + y)^2$ $\frac{d^2y}{dx^2} = (x + y)\left(\frac{dy}{dx} + 1\right)$ $\frac{d^3y}{dx^3} = \left(1 + \frac{dy}{dx}\right)^2 + (x + y)\left(\frac{d^2y}{dx^2}\right)$ $\frac{d^3y}{dx^3} - \left(1 + \frac{dy}{dx}\right)^2 - (x + y)\left(\frac{d^2y}{dx^2}\right) = 0$

When  $x = 0$ ,  $y = 1$

$$\frac{dy}{dx} = \frac{1}{2}(0+1)^2 = \frac{1}{2}$$


$$\frac{d^2y}{dx^2} = (0+1)\left(\frac{1}{2}+1\right) = \frac{3}{2}$$

$$\frac{d^3y}{dx^3} = \left(1+\frac{1}{2}\right)^2 + (0+1)\left(\frac{3}{2}\right) = \frac{15}{4}$$

$$y = 1 + \frac{1}{2}x + \frac{\frac{3}{2}x^2}{2!} + \frac{\frac{15}{4}x^3}{3!} + \dots$$

$$y = 1 + \frac{1}{2}x + \frac{3}{4}x^2 + \frac{5}{8}x^3 + \dots$$

8



$$V = \pi \int_0^{\pi/3} (\tan x)^2 dx$$

$$= \pi \int_0^{\pi/3} (\sec^2 x - 1) dx$$

$$= \pi [\tan x - x]_0^{\pi/3}$$

$$= \pi \left[ \sqrt{3} - \frac{\pi}{3} \right]$$

$$= \sqrt{3}\pi - \frac{\pi^2}{3} \text{ units}^3$$

9(a)

$$(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})^5 (\cos \frac{\pi}{6} - i \sin \frac{\pi}{6})^9$$

$$= (\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})^5 (\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6}))^9$$

$$= (\cos \frac{5\pi}{2} + i \sin \frac{5\pi}{2})(\cos(-\frac{9\pi}{6}) + i \sin(-\frac{9\pi}{6}))$$

$$= \left(e^{i\frac{5\pi}{2}}\right) \left(e^{-i\frac{9\pi}{6}}\right)$$

$$= e^{i\pi}$$

$$= -1$$

9(b)	$z^3 = 27e^{i\frac{\pi}{2}}$ $z^3 = 27e^{i(\frac{\pi}{2} + 2k\pi)}$ $z = 3e^{i(\frac{\pi}{6} + \frac{2k\pi}{3})}, \quad k = -1, 0, 1$ $z_1 = 3(0 - i) = -3i$ $z_2 = 3\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{3\sqrt{3}}{2} + \frac{3}{2}i$ $z_3 = 3\left(-\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = -\frac{3\sqrt{3}}{2} + \frac{3}{2}i$
10(i)	<p>GC:</p> $\alpha = 1.052$ $\beta = 5.505$
10(ii)	$x_{n+1} = \ln(x_n - 1) + 4$ <p>If a sequence converges, as <math>n \rightarrow \infty</math>, <math>x_n \rightarrow L</math> and <math>x_{n+1} \rightarrow L</math>.</p> $L = \ln(L - 1) + 4$ $\ln(L - 1) - L + 4 = 0$ <p>Hence, <math>L = \alpha</math> or <math>L = \beta</math>.</p>
10(iii)	<p>Maximum point: <math>(2, 2)</math></p> $x_{n+1} = \ln(x_n - 1) + 4$ $x_{n+1} - x_n = \ln(x_n - 1) - x_n + 4$ <p>Since <math>\ln(x - 1) - x + 4 \leq 2</math>,</p> $x_{n+1} - x_n \leq 2$ $x_{n+1} \leq x_n + 2$
11	$\frac{dx}{dt} = m - px$ <p>When <math>x = 0.5</math>, <math>\frac{dx}{dt} = 0</math></p> $0 = m - p(0.5)$ $m = 0.5p$ $\frac{dx}{dt} = (0.5p) - px$ $= -\frac{1}{2}p(2x - 1)$ $= -k(2x - 1) \text{ where } k > 0.$

When  $t = 0$ ,  $x = 1.5$ ,  $\frac{dx}{dt} = -0.02$ ,

$$\frac{dx}{dt} = -k(2x - 1)$$

$$\frac{1}{2x-1} \left( \frac{dx}{dt} \right) = -k$$

$$\int \frac{1}{2x-1} dx = -k \int 1 dt$$

$$\frac{1}{2} \ln |2x-1| = -kt + c$$

$$c = \frac{1}{2} \ln 2$$

$$-0.02 = -k(3-1)$$

$$k = 0.01$$


$$\frac{1}{2} \ln |2x-1| = -0.01t + \frac{1}{2} \ln 2$$

When  $x = 1.01$ ,


$$\frac{1}{2} \ln |2(1.01) - 1| = -0.01t + \frac{1}{2} \ln 2$$

$$t = 50 \ln \left( \frac{100}{51} \right) \approx 33.7 \text{ sec}$$

12(i)



12(ii)



13(i)	$\overrightarrow{BA} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ $\overrightarrow{CB} = \begin{pmatrix} 2 \\ -8 \\ 4 \end{pmatrix}$ $\overrightarrow{BA} \times \overrightarrow{CB} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \times \begin{pmatrix} 2 \\ -8 \\ 4 \end{pmatrix} = 2 \begin{pmatrix} 2 \\ 5 \\ 9 \end{pmatrix}$ $\begin{pmatrix} 0 \\ -2 \\ 1 \end{pmatrix} \bullet \begin{pmatrix} 2 \\ 5 \\ 9 \end{pmatrix} = 0 - 10 + 9 = -1$ $\therefore p:r \cdot \begin{pmatrix} 2 \\ 5 \\ 9 \end{pmatrix} = -1$
13(ii)	$l_1 : r = \begin{pmatrix} 1 \\ -4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix}$ $l_2 : r = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} k \\ 2 \\ 3 \end{pmatrix}$ $-4 + \lambda = 1 + 2\mu$ $-1 + 4\lambda = -1 + 3\mu$ $\lambda = -3, \quad \mu = -4$ $1 + 2(-3) = 0 + k(-4)$ $k = \frac{5}{4}$
13(iii)	$\begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix} \bullet \begin{pmatrix} 2 \\ 5 \\ 9 \end{pmatrix} = 45 \neq 0$ <p>Since <math>l_1</math> not perpendicular to plane's normal, <math>l_1</math> does not lie in the plane.</p>

13(iv)	$\left[ \begin{pmatrix} 1 \\ -4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix} \right] \begin{pmatrix} 2 \\ 5 \\ 9 \end{pmatrix} = -1$ $45\lambda = 26$ $\lambda = \frac{26}{45}$ $r = \begin{pmatrix} 1 \\ -4 \\ -1 \end{pmatrix} + \left( \frac{26}{45} \right) \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix} = \begin{pmatrix} \frac{97}{45} \\ -\frac{154}{45} \\ \frac{59}{45} \end{pmatrix}$
13(v)	$\cos \theta = \frac{\frac{5}{2} + 10 + 27}{\left(\sqrt{\frac{233}{16}}\right)\left(\sqrt{110}\right)}$ $\theta = 0.16191 \text{ rad}$ $\square = \frac{\pi}{2} - 0.16191 = 1.41 \text{ rad } (80.7^\circ)$