



Catholic Junior College
JC2 Preliminary Examinations
Higher 2

CANDIDATE
NAME

CLASS

2T

INDEX NUMBER

CHEMISTRY

9729/03

Paper 3 Free Response

9 September 2024

WORKED SOLUTIONS

This document consists of **29** printed pages and **1** blank page.

Section A


Answer **all** the questions in this section.

- 1 (a) Electrocatalysts are a specific form of catalysts that participate in electrochemical reactions. An example is the direct-methanol fuel cells (DMFCs) where methanol is oxidised over an electrocatalyst consisting of a mixture of transition metals such as platinum and rhodium to carbon dioxide.

The main reaction occurring at the anode in the DMFCs is shown below:



However, methanoic acid can also be produced at the anode and it can be further oxidised to carbon dioxide.



Some standard electrode potentials are shown in Table 1.1.

Table 1.1

	$E^\ominus / \text{V}$
$\text{HCO}_2\text{H} / \text{CH}_3\text{OH}$	+0.03
$\text{CO}_2 / \text{HCO}_2\text{H}$	+0.61

- (i) Deduce the type of catalysis in DMFCs. Explain your answer. [1]
- (ii) Explain how transition metals such as platinum and rhodium can act as catalysts. [1]
- (iii) Draw a fully labelled diagram of the experimental set-up used to measure the standard electrode potential of the $\text{CO}_2 / \text{CH}_3\text{OH}$ half-cell. [3]
- (iv) State the relationship between the standard Gibbs free energy change, $\Delta G^\ominus$, and standard cell potential, $E^\ominus_{\text{cell}}$.


Use this relationship, the data in Table 1.1 and an energy cycle to calculate the standard electrode potential for the $\text{CO}_2 / \text{CH}_3\text{OH}$ half-cell. $\Delta G^\ominus$ can be used in the same manner as $\Delta H^\ominus$ in a Hess' law cycle but $E^\ominus$ cannot. [3]

- (v) Platinum(II) complexes are often coloured. Explain why complexes of transition metals are usually coloured. [3]


(i) The type of catalysis in the reaction used in the DMFCs is heterogeneous catalysis. This is because the catalysts, platinum and rhodium, exist in a different physical state / phase (solid state) as that of the reactants (liquid/gas).

- (ii) Some transition metals are good catalyst because of the availability of low-lying partially filled 3d orbitals / 3d and 4s electrons to form temporary bonds with reactants molecules during heterogeneous catalysis.

(iii)



(iv) $\Delta G^\circ = -nFE^\circ_{\text{cell}}$



For $\text{HCO}_2\text{H} + 3\text{H}_2\text{O} + 4\text{e}^- \rightarrow \text{CH}_3\text{OH} + 4\text{OH}^-$ $E^\circ = +0.03\text{V}$

$$\Delta G_2^\circ = -(4)(96500)(+0.03) = -11.6 \text{ kJ mol}^{-1}$$

For $2\text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g}) + 2\text{e}^- \rightarrow \text{HCO}_2\text{H} + 2\text{OH}^-$ $E^\circ = +0.61\text{V}$

$$\Delta G_3^\circ = -(2)(96500)(+0.61) = -117.7 \text{ kJ mol}^{-1}$$

By Hess's Law,

$$\Delta G_1^\circ = \Delta G_2^\circ + \Delta G_3^\circ = (-11.6) + (-117.7) = -129.3 \text{ kJ mol}^{-1}$$

Hence, $\Delta G_1^\circ = -nFE^\circ_{\text{cell}}$

$$(-129300) = -(6)(96500)E$$


$$E^\circ = +0.223\text{V}$$

(v) In the presence of ligands, the partially filled degenerate 3d orbitals are split into 2 groups of non-degenerate 3d orbitals with a small energy gap.



When exposed to visible light, a d electron from the lower energy level absorbs a particular wavelength of light and is promoted to the higher energy level. This is known as d-d* electronic transition.

The complementary colour of the wavelength absorbed is observed as the colour of the complex


- (b) Chloroacetophenone was formerly used as a tear agent and is a potent eye, throat and skin irritant. Compounds **A** and **B** are isomers of chloroacetophenone.



chloroacetophenone

**A****B**

- (i) Suggest a reason why iodoacetophenone is more reactive than chloroacetophenone towards hydrolysis. [1]



iodoacetophenone

- (ii) Arrange chloroacetophenone, **A** and **B**, in order of increasing ease of hydrolysis. Explain your answer fully. [4]

(i) The atomic size of I is larger than Cl and this results in less effective orbital overlap between C and I atom. Hence, the C-I bond length is longer than the C-Cl bond, and thus C-I bond is weaker than the C-Cl bond and easier to be broken.

(ii) In order of increasing ease of hydrolysis: B < chloroacetophenone < A
For compound A (acyl chloride), the carbonyl C atom is bonded to two electronegative atoms O and Cl. This makes the carbonyl C atom highly electron deficient. Hence, the highly electron deficient carbonyl C atom is very susceptible to hydrolysis which occurs readily.

For chloroacetophenone (chloroalkane), there is only one electronegative Cl atom bonded to the alkyl C atom. Hence, the alkyl C atom is less electron deficient and is less susceptible to hydrolysis compared to compound A.

For compound B (chloroarene), the lone pair of electrons on the Cl atom is delocalised into the benzene ring. This strengthens the C-Cl bond due to presence of partial double bond character. Hydrolysis does not occur at all.

(c) Ethanedioic acid, $\text{H}_2\text{C}_2\text{O}_4$, is the simplest dicarboxylic acid and its conjugate base, $\text{C}_2\text{O}_4^{2-}$, is a chelating agent for transition metal cations.


(i) Suggest a two-step synthesis for the formation of ethanedioic acid from ethene. Indicate clearly all reagents and conditions used in the two steps. [3]

(ii) Each ethanedioate ion $\text{C}_2\text{O}_4^{2-}$ can form two co-ordinate bonds, and under certain conditions, copper forms the complex ion $[\text{Cu}(\text{C}_2\text{O}_4)_2]^{2-}$.


Draw the structure of the complex $[\text{Cu}(\text{C}_2\text{O}_4)_2]^{2-}$.

[1]

(i)



(ii)



[Total: 20]

- 2 (a) (i) Explain what is meant by the term *entropy* of a chemical system. [2]
- (ii) Carbon monoxide, CO, has a melting point of 68 K and boiling point of 82 K.


Sketch a graph to show how the entropy of CO changes between 0 to 100 K.

Explain your reasoning. [2]

(i) Entropy of a system measures the degree of disorder. It gives a measure of the extent to which particles and energy are distributed within the chemical system.

The higher the amount of disorder, the higher the entropy of the system.

(ii)




As the temperature increases, the kinetic energy of the CO molecules increases and there are more ways of arranging the energy quanta. Hence there is an increase in disorder of the system and entropy increases.

At 68K and 82K, solid CO molecules change into liquid CO molecules and liquid CO molecules change into gaseous CO molecules respectively.

The orderly arrangement of solid is disrupted and a large increase in volume from liquid to gas, results in a sudden increase in disorder and a sharp increase in entropy.

[1 and shown on graph, indicate increase in disorder at least once]

- (b) The Fischer-Tropsch process is one method of manufacturing methane through the reaction between carbon monoxide and hydrogen.



- (i) Use the data to calculate the entropy change for this reaction at 298 K and comment on its sign with respect to the equation for the reaction. [2]
- (ii) A mixture of CO and H₂ in the molar ratio 1 : 3 is added into a sealed vessel and heated to 800 °C. At equilibrium, the total pressure is 32 atm and 40% of CO has reacted.

Write the expression for the equilibrium constant, K_p , for this reaction. Use your expression to calculate the value of K_p for this reaction. Include its units. [4]

(i) $\Delta G^\ominus = \Delta H^\ominus - T\Delta S^\ominus$

$-156 = -206 - 298 \Delta S^\ominus$

$\Delta S^\ominus = \frac{-156 + 206}{-298} = -0.1678 \text{ kJ K}^{-1} \text{ mol}^{-1} = -168 \text{ J K}^{-1} \text{ mol}^{-1}$

The sign of $\Delta S^\ominus$ is negative, indicating a decrease in entropy which is expected

as the reaction results in the decrease in number of moles of gaseous molecules

from 4 to 2, hence the degree of disorder of the system decreases.

(ii) $K_p = \frac{(p_{\text{CH}_4})(p_{\text{H}_2\text{O}})}{(p_{\text{CO}})(p_{\text{H}_2})^3}$

Let initial amount of CO and H₂ gases be 1 and 3 mol respectively

	CO	3H ₂		CH ₄	H ₂ O
Initial amt / mol	1	3		0	0
Change	-0.40	-3(0.40)		+0.40	+0.40
Equilibrium amt / mol	1 - 0.40 = 0.60	3 - 3(0.40) = 1.80		+0.40	+0.40

Total amount at equilibrium = 0.6 + 1.8 + 0.4 + 0.4 = 3.2 mol

$p_{\text{CO}} = \frac{0.6}{3.2} \times 32 = 6 \text{ atm}$

$p_{\text{H}_2} = \frac{1.8}{3.2} \times 32 = 18 \text{ atm}$

$p_{\text{CH}_4} = p_{\text{H}_2\text{O}} = \frac{0.4}{3.2} \times 32 = 4 \text{ atm}$

$K_p = \frac{(4)(4)}{(6)(18)^3} = \underline{4.57 \times 10^{-4} \text{ atm}^{-2}}$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) The Fischer-Tropsch process can be used to form many different types of hydrocarbons.

Alkynes are a series of non-cyclic hydrocarbons containing one carbon-carbon triple bond per molecule. A five-carbon alkyne has the formula, C_5H_8 , and exhibits constitutional isomerism.

- (i) Explain what is meant by *constitutional isomerism*. [1]
- (ii) Carbon-carbon triple bonds are formed between carbons that are sp hybridised. Explain how sp hybridisation of the carbon atom arises. [2]
- (iii) Draw the structures of all alkynes with the formula C_5H_8 . [2]

.....

(i) Constitutional isomers have the same molecular formula but different structural formulae.

.....

(ii) sp hybridisation is the mixing / combination of a 2s orbital with a 2p orbital to form two sp hybrid orbitals which are used to form the σ bond.

.....

The two unhybridised 2p orbitals are used to form the two π bonds between carbons.



.....

.....

.....

.....

(iii)



[Total: 15]

- 3 (a) Describe how the behaviour of beams of the ions of $^1\text{H}^+$, $^2\text{H}^+$ and the electron, e^- , differ in an electric field. [2]

(a) $^1\text{H}^+$ and $^2\text{H}^+$ will be deflected to the negative plate and electron will be deflected to the positive plate. As angle of deflection $\propto \frac{\text{charge}}{\text{mass}}$, electron will have the largest angle of deflection as it has the largest $\frac{\text{charge}}{\text{mass}}$ ratio, followed by $^1\text{H}^+$ then $^2\text{H}^+$.

- (b) Calcium is a Group 2 element. It has five stable isotopes as shown in Table 3.1.

Table 3.1

Relative isotopic mass of Ca	Percentage abundance / %
39.96	96.93
41.96	0.65
42.96	0.14
43.96	2.09
47.96	0.19

Use the data in Table 3.1 to calculate the relative atomic mass of calcium to two decimal places. [1]

(b) The relative atomic mass of calcium

$$= \frac{39.96 \times 96.93 + 41.96 \times 0.65 + 42.96 \times 0.14 + 43.96 \times 2.09 + 47.96 \times 0.19}{100}$$

$$= 40.08$$

- (c) The mineral dolomite is a double carbonate of magnesium and calcium, with the formula written as $\text{CaMg}(\text{CO}_3)_2$. When one mole of dolomite was heated at 315°C , one mole of gas was collected which produced white precipitate with limewater. When another sample of the same amount of dolomite was heated at 530°C , two moles of gas were collected.

(i) Write an equation for the thermal decomposition of a Group 2 carbonate, XCO_3 . [1]

(ii) By considering the thermal stability of Group 2 carbonates, account for the different moles of gas collected at 315°C and 530°C . [3]



(ii) Mg^{2+} has a smaller ionic radius than Ca^{2+} while their ionic charges are the same. Charge density of Mg^{2+} is higher and hence has higher polarising power than Ca^{2+} . Mg^{2+} is able to polarise / distort the electron cloud of CO_3^{2-} to larger extent, and weaken C–O bond to larger extent and easier to break.

MgCO₃ is less thermally stable / easier to decompose and decompose at 315 °C. Both carbonates will decompose at a higher temperature of 530 °C. Hence, moles of CO₂ evolved is doubled.

- (d) Iodine reacts readily with propanone, CH₃COCH₃, to form 1-iodopropanone, in the presence of an acid catalyst. The rate of this reaction is known to be zero order with respect to iodine and first order with respect to CH₃COCH₃.

(i) Define the term *order of reaction*. [1]

(ii) A series of six separate experiments is carried out, each with a different initial concentration of CH₃COCH₃, but same initial concentration of iodine and acid, and the volume of the reaction mixtures were kept the same. The amount of iodine used is much lower than the amount of CH₃COCH₃, and the time taken for the iodine to decolourise was noted for each experiment.

Describe:

- how the results can be used to calculate the relative rate for each experiment and
- how to graphically show that the reaction is first order with respect to the concentration of CH₃COCH₃. Include a sketch of the graph to be used. [3]


(iii) For a certain concentration of acid used, the rate constant, k , was found to be $2.16 \times 10^{-3} \text{ min}^{-1}$ for the simplified rate equation of $\text{rate} = k [\text{CH}_3\text{COCH}_3]$.

Use the given data to calculate the half-life for this reaction. [1]

(i) The order of reaction with respect to a given reactant is defined as the power to which the concentration of the reactant is raised in the experimentally determined rate equation.

(ii) Since a fixed amount of I₂ was used, the rate is directly proportional to 1/t, hence the relative initial rates = 1/t.

To show that the reaction is first order, a plot of rate versus [CH₃COCH₃] will give a straight-line graph through origin, showing that rate is directly proportional to [CH₃COCH₃] for a first-order reaction.



(iii) For a first order reaction,

$$t_{1/2} = \frac{\ln 2}{k} = \frac{\ln 2}{2.16 \times 10^{-3}} = 321 \text{ min}$$

(e) Nobel laureate Ronald Norrish studied the photodecomposition of propanone with UV light. He showed that the mechanism consists of three steps.

- The first step involves the homolytic fission of the C–C bond of propanone, forming a methyl radical, $\bullet\text{CH}_3$ and another free radical intermediate, **X**.
- The second step involves **X** undergoing further decomposition to produce carbon monoxide and another methyl radical, $\bullet\text{CH}_3$. This involves homolytic fission and formation of a dative bond.
- In the last step, free radicals combine to form ethane.

(i) Draw the 'dot-and-cross' diagram for carbon monoxide. [1]

(ii) Suggest the mechanism for the photodecomposition of propanone, taking note of the formation of carbon monoxide. Use curly arrows to show movement of electrons. [3]

- (iii) When Ronald Norrish performed the photodecomposition of butanone to further study the mechanism, he discovered a mixture of three different alkanes was formed.

Draw the three possible structures of these alkanes.

[1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) $\text{CH}_3\text{—CH}_2\text{CH}_3$ $\text{CH}_3\text{—CH}_3$ $\text{CH}_3\text{CH}_2\text{—CH}_2\text{CH}_3$

.....

.....

.....

.....

.....

.....

BLANK PAGE


(f) Compound **D**, $C_4H_7NO_2$, contains two different functional groups and one chiral carbon atom. Samples of compound **D** are tested separately with

- 2,4-DNPH forming an orange precipitate **E**
- Fehling's reagent in strongly alkaline NaOH solution in the presence of heat to form a red precipitate **F** and organic compound **G**, $C_4H_4O_4Na_2$
- aqueous sodium hydroxide in the presence of heat to form ammonia gas and an organic compound **H**, $C_4H_5O_3Na$.

(i) Suggest possible identities for **D**, **E**, **F**, **G**, and **H**.
For each test, state the type of reaction described. [7]


(ii) Suggest the structure of the organic product when **D** reacts with excess $LiAlH_4$. [1]

(i)




D

Reaction of **D** to give **E**: condensation



E



Reaction of **D** to give **F** and **G**: Oxidation of aliphatic aldehyde to give **F**, Cu_2O and base-catalysed hydrolysis of amide



F

G

Reaction of **D** to give **H**: base-catalysed hydrolysis of amide




[Total: 25]

Section B

Answer **one** question in this section.


- 4 1,4-dibromobutane is a halogenoalkane. It can undergo controlled hydrolysis to form 4-bromobutan-1-ol.

For this reaction, the rate equation is $\text{rate} = k[\text{CH}_2\text{BrCH}_2\text{CH}_2\text{CH}_2\text{Br}][\text{OH}^-]$.




- (a) Name and describe the mechanism involved in the above hydrolysis. [3]


(a) Nucleophilic Substitution ($\text{S}_{\text{N}}2$)





- (b) **C** and **D** can be synthesised from 4-bromobutan-1-ol using the reaction scheme proposed below.



- (i) Suggest the structures of compounds **A** and **B**. [2]
- (ii) In step 4, a carbocation intermediate is formed. Suggest the reagent used. [1]
- (iii) Suggest a simple chemical test to distinguish 4-bromobutan-1-ol from compound **D**. [2]



4-bromobutan-1-ol

**D**(i) **A****B**(ii) **I₂ in hexane and room temperature**(iii) **Test:**

Heat the compounds with aqueous NaOH followed by the addition of dilute aqueous HNO₃. Then add AgNO₃(aq)

For 4-bromobutan-1-ol, cream precipitate of AgBr will be observed.

For compound **D**, yellow precipitate of AgI will be observed.

OR

Test:

Add anhydrous PCl₅ to the compounds at r.t.p.

For 4-bromobutan-1-ol, white fumes of HCl will be observed.

For compound **D**, no white fumes observed.

- (c) Aluminum chloride, AlCl_3 , is commonly employed as a catalyst in halogenation reactions involving aromatic compounds. It is important that AlCl_3 is anhydrous when used in such reactions.

Describe the reaction of AlCl_3 with water and write equations for any reactions that occur. [2]

(c) In the presence of water, AlCl_3 undergoes both hydration and partial hydrolysis.




- (d) As compared to AlCl_3 , the oxide of aluminum, Al_2O_3 , does not react with water.

(i) Explain why Al_2O_3 is insoluble in water. [1]

- (ii) Anodising is an electrolytic process used to increase the thickness of natural oxide layer on the surface of metal parts. Aluminum can be anodised to make it more resistant to corrosion.

Draw a fully labelled diagram, indicating the direction of electron flow in the circuit, and write the equations for both the cathode and anode reactions for the anodising of aluminium. [3]

- (iii) Aluminum chloride, AlCl_3 can react with nitrogen compounds such as trimethylamine, $\text{N}(\text{CH}_3)_3$, as shown by the following equation:




Name the type of bond formed between N and Al in $(\text{CH}_3)_3\text{NAlCl}_3$ and draw the structure of the compound, $(\text{CH}_3)_3\text{NAlCl}_3$. [2]

(i) Ion-dipole attractions formed between ions and water molecules

releases insufficient energy to overcome the strong electrostatic

forces of attractions between Al^{3+} cations and O^{2-} anions and


intermolecular hydrogen bonding between water molecules.




At anode, $2\text{Al(s)} + 3\text{H}_2\text{O(l)} \rightarrow \text{Al}_2\text{O}_3\text{(s)} + 6\text{H}^+\text{(aq)} + 6\text{e}^-$

At the cathode, $2\text{H}^+\text{(aq)} + 2\text{e}^- \rightarrow \text{H}_2\text{(g)}$

(iii) The bond formed between N and Al in H_3NAlCl_3 is a dative bond.



(e) Lidocaine is commonly used as local anaesthetic for dental surgeries and minor operations.



Lidocaine

$\text{p}K_b = 6.1$

- (i) Lidocaine is a basic compound. Calculate the pH of a $0.025 \text{ mol dm}^{-3}$ solution of Lidocaine. [2]
- (ii) Hydrochloric acid is added to 1 dm^3 of the Lidocaine solution in (i) to produce a buffer solution. Determine the volume of $0.500 \text{ mol dm}^{-3} \text{ HCl}$ required to form a buffer solution of pH 7. [2]


$$(i) [\text{OH}^-] = \sqrt{K_b \times [\text{B}]}$$

$$= \sqrt{10^{-6.1} \times 0.025}$$

$$= 1.41 \times 10^{-4} \text{ mol dm}^{-3}$$

$$\text{pOH} = 3.85$$

$$\text{pH} = 10.1$$



Let x be the number of moles of HCl required, and v be the total volume.

$$pOH = pK_b + \lg \frac{[BH^+]}{[B]}$$

$$14 - 7 = 6.1 + \lg \frac{\frac{x}{v}}{(0.025 - x) \frac{1}{v}}$$


$$\frac{x}{0.025 - x} = 7.94$$

$$x = 2.23 \times 10^{-2}$$

$$\begin{aligned} \text{Volume of HCl required} &= \frac{2.23 \times 10^{-2}}{0.5} \\ &= 4.46 \times 10^{-2} \text{ dm}^3 \\ &= \underline{\underline{44.6 \text{ cm}^3}} \end{aligned}$$

[Total: 20]

- 5 (a) Hydrazine, N_2H_4 , is used in various rocket fuels. In the engine, hydrazine is passed over a suitable catalyst and decompose to its constituent elements, making hydrazine an efficient propellant. Hydrazine can be synthesised from the reaction between ammonia and hydrogen peroxide.



Relevant standard enthalpy change of formation data are shown in Table 5.1.

Table 5.1

	$\text{NH}_3(\text{g})$	$\text{H}_2\text{O}_2(\text{l})$	$\text{H}_2\text{O}(\text{l})$
$\Delta H_f^\ominus / \text{kJ mol}^{-1}$	-46.1	-187.8	-285.8

- (i) Write a balanced equation for the decomposition of hydrazine to its elements. [1]
- (ii) Calculate the standard enthalpy change for the decomposition of hydrazine to its elements and explain why this reaction makes hydrazine an efficient propellant. [3]
- (iii) Calculate the standard Gibbs free energy change, $\Delta G^\ominus$, for the reaction between ammonia and hydrogen peroxide, given the standard entropy change of the reaction is $-233.6 \text{ J mol}^{-1} \text{ K}^{-1}$. [1]
- (iv) Suggest a reason, in terms of structure and bonding, why ammonia is a gas but hydrazine, N_2H_4 , is a liquid under standard conditions. [2]



.....

(ii) $\Delta H^\ominus = \sum \Delta H_f^\ominus (\text{products}) - \sum \Delta H_f^\ominus (\text{reactants})$

.....

$$= [\Delta H_f^\ominus (\text{N}_2\text{H}_4(\text{l})) + 2\Delta H_f^\ominus (\text{H}_2\text{O}(\text{l}))] - [2\Delta H_f^\ominus (\text{NH}_3(\text{g})) + \Delta H_f^\ominus (\text{H}_2\text{O}_2(\text{l}))]$$

.....

$$-241.0 = \Delta H_f^\ominus (\text{N}_2\text{H}_4(\text{l})) + 2(-285.8) - 2(-46.1) - (-187.8)$$

.....

$$\Delta H_f^\ominus (\text{N}_2\text{H}_4(\text{l})) = +50.6 \text{ kJ mol}^{-1}$$

.....

Enthalpy change for the decomposition of hydrazine

.....

$$= -50.6 \text{ kJ mol}^{-1}$$

.....

This reaction is exothermic and releases large volume of gases /

.....

increase in moles of gases in a very short period, making hydrazine

.....

an efficient propellant.

.....

.....

.....

.....

.....

(iii) $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$

24

$$= -241.0 - [298 (-233.6 \times 10^{-3})]$$

$$= -171 \text{ kJ mol}^{-1}$$

(iv) Both ammonia and hydrazine have simple molecular structure.

More energy is required to overcome the more extensive intermolecular hydrogen bonding between hydrazine molecules than that of ammonia.

Hence, hydrazine has higher boiling point than ammonia.

(b) Hydrazine, N_2H_4 , is a weak base. It dissociates according to the equation shown.




Fig. 5.1 shows the change in pH when 0.10 mol of aqueous hydrazine is titrated with $\text{HCl}(\text{aq})$.




Fig. 5.1


- (i) Describe and explain the relative basicities of hydrazine and ammonia. [2]
- (ii) Show clearly on the graph in Fig. 5.1 the region where the mixture is acting as a buffer solution. [1]
- (iii) Identify the two major species in this buffer solution and write two equations to show how the mixture can act as a buffer solution on the addition of acid and alkali. [2]
- (iv) Suggest and explain, with the aid of an equation, whether the pH at the equivalence point is more or less than pH 7 in this titration. [2]

(b) (i) NH₂ group in hydrazine attached to another N is an electron withdrawing group. Thus the lone pair of electrons on the N atom in hydrazine is less available to accept H⁺ compared to ammonia.

Hydrazine is less basic than ammonia.

(iii) The two major species in this buffer solution are N₂H₄ and N₂H₅⁺ or N₂H₅⁺Cl⁻.

When small amount of acid is added,



When small amount of alkali is added,



(iv) N₂H₅⁺Cl⁻ is formed at the equivalence point. The pH at the equivalence point is less than 7.



N₂H₅⁺ undergoes hydrolysis to produce H₃O⁺.

The presence of excess H₃O⁺ ions or [H⁺] > [OH⁻] in the solution accounts for

pH < 7.

- (c) Hydrazine is used in Wolff-Kishner reduction, a reaction that converts carbonyl compounds to alkanes as shown in Fig. 5.2.




Fig. 5.2

In the first step of the Wolff-Kishner reduction mechanism, hydrazine reacts with a carbonyl compound to form a hydrazone as shown in Fig. 5.3.




Fig. 5.3

In the second last step of the mechanism, an azo compound formed from the hydrazone undergoes deprotonation of the N-H group to produce N_2 gas and a carbanion as shown in Fig. 5.4. The carbanion is then protonated in the last step to form an alkane product.






Fig. 5.4

- (i) On Fig. 5.4, draw three curly arrows to show the mechanism for this step. [1]
- (ii) Suggest the structure of a carbonyl compound and the derivative of hydrazine that would form the following hydrazone by a similar reaction to that shown in Fig. 5.3. [1]




(ii)




- (d) Singapore competed at the 2012 Summer Olympics in London and won two bronze medals in table tennis. Bronze medals contain copper. A redox titration is used to determine the percentage by mass of copper in the bronze medal.

- A 0.800 g sample of a bronze medal is dissolved in hot concentrated nitric acid. After cooling, an excess of potassium iodide solution was added to form iodine and the solution was made up to 250 cm³ in a volumetric flask.



- 25.0 cm³ of this solution was then titrated with aqueous sodium thiosulfate solution. 12.20 cm³ of 0.100 mol dm⁻³ sodium thiosulfate was required for complete reaction, as shown.



- (i) Calculate the percentage by mass of copper in the bronze metal. [2]
- (ii) All medals had a radius of 42.5 mm and thickness of 7 mm. The silver medal was made up of 92.5% silver and 7.5% copper, by mass. Given the density of silver is 10.49 g cm⁻³ and copper is 8.96 g cm⁻³, calculate the mass of a silver medal, assuming that the density of the alloy varies proportionally to its composition by mass.

The equation for the volume of a medal, V, is given below:

$$V = \pi \times (\text{radius})^2 \times \text{thickness}$$

[2]

$$\text{(d) (i) Amount of } \text{S}_2\text{O}_3^{2-} = 0.100 \times \frac{12.20}{1000}$$

$$= 1.22 \times 10^{-3} \text{ mol}$$

$$\text{Amount of } \text{I}_2 \text{ in } 25.0 \text{ cm}^3 = 1.22 \times 10^{-3} \div 2$$

$$= 6.10 \times 10^{-4} \text{ mol}$$

$$\text{Amount of } \text{I}_2 \text{ in } 250 \text{ cm}^3 = 6.10 \times 10^{-4} \times 10$$

$$= 6.10 \times 10^{-3} \text{ mol}$$

$$\text{Mass of } \text{Cu}^{2+} = 6.10 \times 10^{-3} \times 2 \times 63.5$$

$$= 0.7747 \text{ g}$$

$$\text{Percentage by mass of Cu in the bronze medal}$$

$$= \frac{0.7747}{0.800} \times 100$$

$$= 96.8\%$$

$$(ii) \text{ Volume of medal} = \pi \times 4.25^2 \times 0.7$$

$$= 39.72 \text{ cm}^3 \text{ or } 39722 \text{ mm}^3$$

$$\text{Density of medal} = (0.925 \times 10.49) + (0.075 \times 8.96)$$

$$= 10.38 \text{ g cm}^{-3}$$

$$\text{Mass of medal} = 39.72 \times 10.38 = 412 \text{ g}$$

[Total: 20]

Additional answer space

If you use the following pages to complete the answer to any questions, the question number must be clearly shown.

[illegible]

