Candidate Name	Form Class	Index Number



ANG MO KIO SECONDARY SCHOOL PRELIMINARY EXAMINATION 2024 SECONDARY FOUR EXPRESS

CHEMISTRYPaper 1 Multiple Choice

6092/01 27 August 2024 1 hour

Setter: Mr Vincent Voo

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Do not use staples, paper clips, glue or correction fluid.

Write your name, class and index number on the Question Paper and on the Answer Sheet in the spaces provided.

There are **forty** questions on this paper. Answer **all** questions. For each question there are four possible answers **A**, **B**, **C** and **D**.

Choose the **one** you consider correct and record your choice **in soft pencil** on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer. Any rough working should be done on this question paper.

A copy of the Periodic Table is printed on page 16.

The use of an approved scientific calculator is expected, where appropriate.

This document consists of **16** printed pages.

1	Which apparatus can be used to determine the end-point in a volumetric experiment involving
	the addition of dilute hydrochloric acid to 25.0 cm ³ of aqueous sodium hydroxide?

A electronic balance

B gas syringe

C stopwatch

D thermometer

Which of the following procedure describes the best method to obtain a pure sample of barium chloride from a mixture of barium chloride and barium sulfate salts?

	step 1	step 2	step 3	step 4
Α	dissolution	crystallisation	filtration	evaporation
В	dissolution	filtration	evaporation	crystallisation
С	evaporation	crystallisation	dissolution	filtration
D	filtration	dissolution	evaporation	crystallisation

3 100 cm³ of carbon dioxide gas diffuses through a porous pot in 50 s. 100 cm³ of gas X takes more than 50 s to diffuse through the same porous pot under the same room conditions.

Which of the following is gas X?

- **A** ammonia
- **B** hydrogen
- **C** methane
- **D** sulfur dioxide
- 4 Which statement explains why sodium chloride has a lower melting point than magnesium oxide?
 - A Sodium metal is more reactive than magnesium metal.
 - **B** The attraction between Na⁺ and C Γ is weaker than that between Mg²⁺ and O²⁻.
 - **C** The boiling point of sodium chloride is lower than that of magnesium oxide.
 - **D** The bonding in sodium chloride is covalent while it is ionic in magnesium oxide.


5 The diagram shows a molecule of a substance made up of three different elements, P, X and Y.

$$\begin{array}{c|c}
X & X & X & X \\
X & Y & X & Y \\
X & Y & X & X
\end{array}$$

If P is the element phosphorus, what elements could X and Y be?

	X	Y
Α	Н	0
В	0	CI
С	S	Na
D	Mg	Na

6 The structure of an ionic compound is as shown.



Which compound cannot have this arrangement of its ions?

- A lithium oxide
- **B** magnesium carbonate
- **C** potassium chloride
- **D** zinc sulfate

7 The formula of sodium silicate, which is used in the manufacture of silica gel, is Na₂SiO₃.

What is the formula of magnesium silicate?

- A MgSiO₃
- B Mg₂SiO₃
- \mathbf{C} Mg(SiO₃)₂
- **D** $Mg_3(SiO_3)_2$
- 8 The table shows the proton number and nucleon number of elements X and Y.

element	proton number	nucleon number
Х	9	17
Y	13	28

What is the mass of one mole of the compound formed between elements X and Y?

- **A** 22 g
- **B** 45 g
- **C** 79 g
- **D** 101 g
- 9 The mass of mixture M containing barium carbonate and barium sulfate is 8.0 g. Excess nitric acid is added to the mixture and the resulting content is then filtered. The mass of the residue and filtrate is found to be 5.2 g and 13.4 g respectively.

What is the percentage purity of barium sulfate in mixture M?

- $\frac{A}{8.0} \times 100\%$
- $\frac{\text{B}}{8.0} \times 100\%$
- $\frac{\text{C}}{13.4} \times 100\%$
- $\frac{D}{13.4} \times 100\%$

10 The complete combustion of ethane is represented by the following equation.

$$C_2H_4(g) + 3 O_2(g) \rightarrow 2 CO_2(g) + 2 H_2O(I)$$

60 cm³ of ethane reacts with 120 cm³ of oxygen in a closed container.

What is the total volume of gas collected at room temperature at the end of the reaction?


- **A** 60 cm³
- **B** 80 cm³
- **C** 100 cm³
- **D** 120 cm³
- 11 The pH of a sample of dilute hydrochloric acid is 2.

What will be the pH of the acid after the addition of 10 g of sodium chloride salt?


- **A** pH 1
- **B** pH 2
- **C** pH 7
- **D** pH 9
- Which of the following set of reactants is most appropriate to prepare a pure sample of copper(II) chloride crystals?
 - A copper and hydrochloric acid
 - B copper(II) carbonate and hydrochloric acid
 - **C** copper(II) hydroxide and ammonium chloride
 - **D** copper(II) nitrate and potassium chloride

Which graph shows the changes in pH as excess dilute hydrochloric acid is added to aqueous sodium hydroxide?







В



C



D



14	VVIIICII	of the following will react with aqueous ammonium surface to produce an alkaline gas
	Α	calcium hydroxide
	В	copper(II) oxide
	С	nitric acid
	D	sodium chloride
15	In the	Haber process, nitrogen and hydrogen react to form ammonia.
	What i	s the source of hydrogen?
	Α	cracking of oil fractions
	В	electrolysis of water
	С	fractional distillation of liquid air
	D	reaction between acid and metal
16	Which	of the following can increase the yield of ammonia in the Haber process?
	Α	increase the pressure to 450 atm
	В	increase the temperature to 1000 °C
	С	use equal volume of nitrogen and hydrogen
	D	use platinum as a catalyst
17	Which	of the following pairs of solutions will show no visible change when mixed?
	Α	barium nitrate and hydrochloric acid
	В	lead(II) nitrate and sodium sulfate
	С	sodium carbonate and magnesium nitrate
	D	zinc and copper(II) sulfate

18 A salt has the chemical formula (NH₄)₂Fe(SO₄)₂.12H₂O.

In an experiment, excess sodium hydroxide was added slowly, with shaking, to an aqueous solution of the salt until there is no further changes. The boiling tube was then left to stand.


Which of the following observations would not be made?

- A A green precipitate was produced.
- **B** A pungent gas which turned damp red litmus paper blue was produced.
- **C** On standing, a brown precipitate was produced.
- **D** The precipitate dissolved in excess aqueous sodium hydroxide.
- 19 Both aqueous sodium carbonate and sodium hydroxide solution turn red litmus paper blue.

Which one of the following substances can be used to differentiate the two solutions?

- A aqueous calcium nitrate
- B aqueous potassium chloride
- **C** carbon dioxide
- **D** dilute hydrochloric acid
- 20 In which reaction is the underlined substance behaving as an oxidising agent?
 - A $Cl_2 + 2 NaBr \rightarrow Br_2 + 2 NaCl$
 - **B** \underline{HCI} + NaOH \rightarrow NaCl + H₂O
 - C $2 \text{ FeCl}_2 + \text{Cl}_2 \rightarrow 2 \text{ FeCl}_3$
 - D $2 \text{ NaOH} + \text{FeCl}_2 \rightarrow \text{Fe(OH)}_2 + 2 \text{ NaCl}$
- 21 In which reaction does the oxidation state of iron remain unchanged?
 - A Fe + $2 \text{ FeCl}_3 \rightarrow 3 \text{ FeCl}_2$
 - **B** Fe₂O₃ + 6 HCl \rightarrow 2 FeCl₃ + 3 H₂O
 - **C** $2 \text{ Fe} + 3 \text{ Cl}_2 \rightarrow 2 \text{ FeCl}_3$
 - $D \qquad 2 \text{ FeCl}_2 + \text{Cl}_2 \rightarrow 2 \text{ FeCl}_3$

In the set-up shown, the lamp did not light up. Distilled water was then added to the beaker and the lamp lighted up.



Which statement explains these observations?

- A Electrons are free to move in the solution when sodium chloride dissolves.
- **B** Metal ions are free to move when sodium reacts with water.
- C Non-metal ions are free to move when sodium chloride dissolves.
- **D** Oppositely charged ions are free to move when sodium chloride dissolves.
- In an electrolysis experiment, the same amount of charge deposits 19.5 g of zinc and 11 g of manganese.

Which is the charge of the manganese ion? [Ar: Mn, 55; Zn, 65]

- **A** +2
- **B** +3
- **C** +4
- D +7
- 24 Many properties of an element and its compounds can be predicted from the position of the element in the Periodic Table.

What property could **not** be predicted from the position of the element in the Periodic Table?

- A the number of isotopes the element has
- **B** the formula of the oxides of the element
- **C** the acidic and basic nature of the oxides of the element
- **D** the metallic and non-metallic properties of the element

25 A student compares the properties of two Group 1 elements, lithium and potassium.

Which of the following is correct?

	higher melting point	more vigorous reaction with water
Α	lithium	lithium
В	lithium	potassium
С	potassium	lithium
D	potassium	potassium

26 Elements X and Y are in Group 17 of the Periodic Table. X is a liquid and Y is a solid at room temperature.

What statement is/are correct?

- 1 Atoms of Y have more electrons than atoms of X.
- 2 A molecule of Y has more atoms than a molecule of X.
- 3 X displaces Y from an aqueous solution of Y⁻ ions.
- A 1 only
- B 2 only
- C 1 and 3 only
- **D** 1, 2 and 3
- 27 Metal X and its compound undergo the following reactions.
 - $1 \qquad XO + H_2 \rightarrow X + H_2O$
 - 2 $X + 2 HCI \rightarrow XCI_2 + H_2$

What could metal X be?

- A calcium
- **B** iron
- **C** magnesium
- **D** potassium

- Which of the following aqueous solutions, when added to aqueous iron(II) chloride, takes part in a redox reaction?
 - A ammonia
 - **B** silver nitrate
 - C potassium hydroxide
 - **D** potassium manganate(VII)
- 29 Metals P, Q, R and S are placed in salt solutions as shown in the table.

metal	salt of P	salt of Q	salt of R	salt of S
Р	no reaction	no reaction	no reaction	S displaced
Q	P displaced	no reaction	no reaction	S displaced
R	P displaced	Q displaced	no reaction	S displaced
S	no reaction	no reaction	no reaction	no reaction

What is the order of reactivity of the metals in increasing order?

- **A** Q, P, S, R
- **B** R, Q, P, S
- **C** S, P, Q, R
- **D** S, Q, P, R

30 Ethyne (H–C≡C–H) undergoes addition of hydrogen to form ethane as shown.

$$C_2H_2 \ + \ 2 \ H_2 \ \to \ C_2H_6$$

The average bond energies of the following bonds are shown in the table.

bond	C–H	C–C	C=C	C≣C	H–H
bond energy / kJ/mol	413	347	614	839	432

What is the enthalpy change for the reaction?

- **A** −296 kJ/mol
- **B** −176 kJ/mol
- **C** +176 kJ/mol
- **D** +296 kJ/mol

31 Which of the following reactions will pressure least likely to affect the speed of reaction?

- A $C(s) + CO_2(g) \rightarrow 2CO(g)$
- $\mathbf{B} \qquad C_2H_4\left(g\right) \ + \ \mathsf{Br}_2\left(\mathsf{I}\right) \ \to \ C_2H_4\mathsf{Br}_2\left(\mathsf{I}\right)$
- **C** 2 Na (s) + $2H_2O(I) \rightarrow 2 \text{ NaOH (aq)} + H_2(g)$
- **D** $2 SO_2(g) + O_2(g) \rightarrow 2 SO_3(g)$

When excess magnesium ribbon is added to dilute hydrochloric acid, the reaction soon becomes slower and finally stops.

Which of the following statements best explains this observation?

- A The concentration of the acid is decreasing until it finally becomes zero.
- **B** The magnesium ribbon becomes smaller and finally 'dissolves' completely.
- **C** The magnesium ribbon is become covered with an insoluble layer of precipitate.
- **D** The temperature of the reaction mixture decreases during the reaction.
- Which of the following statements is true of petroleum?
 - A Bioethanol can be obtained from petroleum.
 - **B** Burning petroleum mainly produces carbon dioxide and water.
 - **C** Each fraction of petroleum has a fixed boiling point.
 - **D** Petroleum is a compound made up of carbon and hydrogen atoms.

- Why are biofuels produced from plants considered to be carbon neutral?
 - A Biofuels do not produce carbon dioxide when burnt.
 - **B** Biofuels produce carbon monoxide, which is neutral, when burnt.
 - **C** Plants give out carbon dioxide during respiration.
 - **D** Plants take in carbon dioxide during photosynthesis.
- 35 Which of the following cannot be used to distinguish between propanol and propanoic acid?
 - **A** acidified potassium manganate(VII)
 - B red litmus paper
 - **C** calcium carbonate
 - **D** zinc
- A hydrocarbon is heated strongly in the presence of a catalyst. The resulting mixture consists of butane, ethene and propene in the molar ratio of 2 : 1 : 2.

What is the molecular formula of the hydrocarbon?

- **A** C₉H₁₈
- **B** C₉H₂₀
- $C C_{16}H_{32}$
- **D** $C_{16}H_{36}$
- 1 mole of a compound X reacts completely with 2 moles of hydrogen gas in the presence of a catalyst to form 1 mole of alkane.

What compound could X be?

- A CH₂=CH-CH₂-CH₃
- B CH₂=CH-CH=CH₂
- C CH₂=CH-CH=CH-O-H
- D CH₂=CH-CH=CH-CH=CH₂

38 Polymer X has a similar linkage as nylon.

Which of the following is likely to be the monomer of polymer X.


39 Poly(methyl methacrylate) is used to make hard contact lenses. Part of its polymer chain is shown below.

$$\begin{array}{c|cccc} \mathbf{CH_3} & \mathbf{CH_3} \\ \mathbf{-CH_2} - \mathbf{C} & \mathbf{-CH_2} - \mathbf{C} & \mathbf{--} \\ \mathbf{C=O} & \mathbf{C=O} \\ \mathbf{O} & \mathbf{O} \\ \mathbf{CH_3} & \mathbf{CH_3} \end{array}$$

Which statements about poly(methyl methacrylate) are correct?

- 1 Its monomer is CH₂=CCH₃CO₂CH₃.
- 2 It can be classified as an acid.
- 3 It can be classified as an ester.
- 4 It is a condensation polymer.
- A 1 and 2 only
- B 1 and 3 only
- C 3 and 4 only
- **D** 2, 3 and 4 only

40 The diagram shows a simplified process to remove flue gas.



Which gas cannot be removed using the method shown?

- A carbon monoxide
- B nitrogen dioxide
- **C** phosphorus trioxide
- **D** sulfur dioxide

The Periodic Table of Elements

		18	무 무	helium 4	10	Ne	20 20	18	Ā	argon 40	36	궃	krypton 84	24	Xe	xenon 131	98	몺	radon	118	ő	oganesson	ı												
		17			6	ட	fluorine 19	17	C	chlorine 35.5	35	ğ	bromine 80	53	Н	iodine 127	85	¥	astatine	117	<u>r</u>	tennessine	1												
		16			8	0	oxygen 16	16	ဟ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	8	mninolod	116	ک	livermorium	ı												
		15			7	z	nitrogen 14	15	۵.	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209	115	ğ	moscovium	1												
		14			9	ပ	carbon 12	41	:S	silicon 28	32	Ge	germanium 73	20	S	119 119	82	Ъ	lead 207	114	F1	flerovium	ı												
		13			5	М	boron 11	13	ΡΊ	aluminium 27	31	Ga	gallium 70	49	Ľ	indium 115	81	<i>1</i> 1	thallium 204	113	Ę	nihonium	1												
					L			1		12	30	Zu	zinc 65	48	ප	cadmium 112	8	Ę	mercury 201	112	ర్	copernicium	1												
										1	59	ನ	copper 64	47	Ag	silver 108	79	Αu	gold 197	111	Rg	oentgenium	1												
	dr			_					<u></u>					01										Z	nickel 59	46	В	palladium 106	78	盂	platinum 195	110	Ds Ds	armstadtium	1
	Group									တ	27	ပိ	cobalt 59	45	뫈	rhodium 103	22	ä	iridium 192	109	₹	meitnerium	1												
			H hydrogen						œ	56	Pe	ion 56	44	æ	ruthenium 101	9/	SO	osmium 190	108	£	hassium	1													
				J					7	25	Ā	manganese 55	43	ည	technetium 	75	Re	rhenium 186	107	뮵	pohrium	1													
				Key	ımber		Jass			9	54	ပ်	chromium 52	42	õ	molybdenum 96	74	>	tungsten 184	106	Sg	seaborgium	1												
					Key	proton (atomic) number	nic symt	name relative atomic mass			2	\vdash			\vdash		niobium 93				\vdash			\neg											
					proton (ator	relativ			4	22	i=	titanium 48	40	ZĽ	zirconium 91	72	Ï	hafinium 178	104	弘	utherfordium	1												
					L			Į		က	21	လွ	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89-103	actinoids														
		2			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	റ്	strontium 88	56	Ba	barium 137	_		radium	1												
		-			3	:=	lithium 7	7	Ra	sodium 23	19	エ	potassium 39	37	8 2	rubidium 85	55	ပ	caesium 133	87	占	francium	1												
١		<u> </u>	I					_																											

02	Tm Yb Lu	ytterbium	173	102	2	m nobelium la	
	<u>—</u>					Ĕ	1
	운	holmium	165	66	Es	einsteinium	ı
99	٥	dysprosium	163	86	చ	californium	ı
65	<u>P</u>	terbium	159	26	益	berkelium	ı
64	g	gadolinium	157	96	S	curium	ı
63	Ш	europium	152	92	Am	americium	ı
62	Sm	samarium	150	94	Pu	plutonium	ı
61	Pa	promethium	ı	93	å	neptunium	ı
09	P	neodymium	144	92	-	uranium	238
29	ፈ	praseodymium	141	91	Ра	protactinium	231
28	රී	cerinm	140	06	드	thorium	232
22	Ľ	lanthanum	139	88	Ac	actinium	ı
	lanthanoids			actinoids	5		

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.). The Avogadro constant, $L=6.02\times10^{23}\,\text{mol}^{-1}$.

BLANK PAGE