





## H2 Mathematics (9758)

### Chapter 11 Definite Integrals

### Extra Practice Solutions

Qn 1	2009/VJC Prelim/1/10
	$\int xe^{2x} dx = \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x} dx$ $= \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + c$ $= \frac{e^{2x}}{4}(2x-1) + c \text{ (shown)}$
(a)	<p>Required area</p> $= \int_0^4 \left( \frac{1}{2}e^4 x - e^x \sqrt{x} \right) dx + \int_4^8 \left( e^x \sqrt{x} - \frac{1}{2}e^4 x \right) dx$ $= 7239.2 \text{ units}^2 \text{ (1 dp)}$
(b)	<p>Required Volume</p> $= \frac{1}{3}\pi(2e^4)(4) - \int_0^4 \pi(e^{2x}x) dx$ $= \frac{16\pi e^8}{3} - \pi \left[ \frac{e^{2x}}{4}(2x-1) \right]_0^4$ $= \frac{16\pi e^8}{3} - \pi \left( \frac{e^8}{4} \right)(7) - \frac{1}{4}\pi$ $= \pi e^8 \left( \frac{43}{12} \right) - \frac{\pi}{4} \text{ units}^3$ $A = \frac{43}{12} \text{ and } B = \frac{1}{4}$


Qn 2	2011/TJC Prelim/1/10
	 <p>Area = <math>\frac{1}{2}(\sqrt{2})(3\sqrt{2}) + \int_{\sqrt{2}}^{\sqrt{3}} \frac{6}{x} dx - \frac{1}{2}(\sqrt{3})(2\sqrt{3})</math></p> $= [6\ln x]_{\sqrt{2}}^{\sqrt{3}}$ $= 3(\ln 3 - \ln 2) \text{ units}^2$
	 <p><math>x^2 + y^2 - 4y + 3 = 0 \Rightarrow x^2 + (y-2)^2 = 1</math></p> <p>Volume = <math>2 \int_0^1 (\pi y_1^2 - \pi y_2^2) dx</math> where <math>y_1 = 2 + \sqrt{1-x^2}</math> and <math>y_2 = 2 - \sqrt{1-x^2}</math></p> $= 2\pi \int_0^1 \left[ (2 + \sqrt{1-x^2})^2 - (2 - \sqrt{1-x^2})^2 \right] dx$ $\approx 39.5 \text{ units}^3 \text{ (3 s.f.)}$


Qn3	2013/VJC Prelim/1/10
(i)	$\int_0^{\frac{1}{\sqrt{2}}} \cos^{-1} x \, dx = \left[ x \cos^{-1} x \right]_0^{\frac{1}{\sqrt{2}}} - \int_0^{\frac{1}{\sqrt{2}}} \left( -\frac{x}{\sqrt{1-x^2}} \right) dx$ $= \frac{1}{\sqrt{2}} \left( \frac{\pi}{4} \right) - \left[ \sqrt{1-x^2} \right]_0^{\frac{1}{\sqrt{2}}}$ $= \frac{\pi}{4\sqrt{2}} - \left( \sqrt{\frac{1}{2}} - 1 \right)$ $= \frac{\pi}{4\sqrt{2}} + 1 - \frac{1}{\sqrt{2}}$
(ii)	$\text{Volume} = \pi \int_0^{\frac{1}{\sqrt{2}}} y^2 \, dx - \pi \left( \frac{\sqrt{\pi}}{2} \right)^2 \left( \frac{1}{\sqrt{2}} \right)$ $= \pi \int_0^{\frac{1}{\sqrt{2}}} \cos^{-1} x \, dx - \frac{\pi^2}{4\sqrt{2}}$ $= \frac{\pi^2}{4\sqrt{2}} + \pi - \frac{\pi}{\sqrt{2}} - \frac{\pi^2}{4\sqrt{2}}$ $= \pi - \frac{\pi}{\sqrt{2}} \text{ unit}^3$
(iii)	<p>Required Equation: <math>y = \sqrt{\cos^{-1} x} - \frac{\sqrt{\pi}}{2}</math>.</p> $\text{Volume} = \pi \int_0^{\frac{1}{\sqrt{2}}} \left( \sqrt{\cos^{-1} x} - \frac{\sqrt{\pi}}{2} \right)^2 dx$ $\approx 0.116 \text{ (3 s.f.)}$

Qn 4	2013/TJC Prelim/1/9
(i)	$\int e^{2x} \cos 4x \, dx = \frac{1}{2} e^{2x} \cos 4x - \frac{1}{2} \int e^{2x} (-4 \sin 4x) \, dx$ $= \frac{1}{2} e^{2x} \cos 4x + 2 \int e^{2x} \sin 4x \, dx$ $= \frac{1}{2} e^{2x} \cos 4x + 2 \left( \frac{1}{2} e^{2x} \sin 4x - \frac{1}{2} \int e^{2x} (4 \cos 4x) \, dx \right)$ $= \frac{1}{2} e^{2x} \cos 4x + e^{2x} \sin 4x - 4 \int e^{2x} \cos 4x \, dx$ $5 \int e^{2x} \cos 4x \, dx = \frac{1}{2} e^{2x} (\cos 4x + 2 \sin 4x) + C$ $\int e^{2x} \cos 4x \, dx = \frac{1}{10} e^{2x} (\cos 4x + 2 \sin 4x) + C$

<b>(ii)</b>	<p>At the point of intersection,  <math>e^x \sin 2x = e^x \Rightarrow e^x (\sin 2x - 1) = 0</math>  either <math>e^x = 0</math>      or    <math>\sin 2x = 1</math></p> <p>(reject, as <math>e^x &gt; 0</math>)      or      <math>2x = \frac{\pi}{2}</math>  <math>x = \frac{\pi}{4}</math></p>
<b>(iii)</b>	<p>Volume of solid generated = <math>\pi \int_0^{\frac{\pi}{4}} (e^x)^2 dx - \pi \int_0^{\frac{\pi}{4}} (e^x \sin 2x)^2 dx</math></p> $= \pi \int_0^{\frac{\pi}{4}} e^{2x} dx - \pi \int_0^{\frac{\pi}{4}} e^{2x} \sin^2 2x dx$ $= \pi \int_0^{\frac{\pi}{4}} e^{2x} dx - \pi \int_0^{\frac{\pi}{4}} e^{2x} \left( \frac{1 - \cos 4x}{2} \right) dx$ $= \frac{1}{2} \pi \int_0^{\frac{\pi}{4}} (e^{2x} + e^{2x} \cos 4x) dx$ $= \frac{1}{2} \pi \left( \left[ \frac{1}{2} e^{2x} \right]_0^{\frac{\pi}{4}} + \left[ \frac{1}{10} e^{2x} (\cos 4x + 2 \sin 4x) \right]_0^{\frac{\pi}{4}} \right)$ $= \frac{1}{2} \pi \left( \frac{1}{2} \left( e^{\frac{\pi}{2}} - 1 \right) + \frac{1}{10} \left( -e^{\frac{\pi}{2}} - e^0 \right) \right) \quad \text{using (i) answer}$ $= \frac{1}{10} \pi \left( 2e^{\frac{\pi}{2}} - 3 \right) \text{ units}^3$

<b>Qn 5</b>	<b>2017/MJC Promo/4a</b>
	$\int_0^3  x^3 - 5x^2 + 4x  \, dx = \int_0^1 x^3 - 5x^2 + 4x \, dx - \int_1^3 x^3 - 5x^2 + 4x \, dx$ $= \left[ \frac{x^4}{4} - \frac{5x^3}{3} + \frac{4x^2}{2} \right]_0^1 - \left[ \frac{x^4}{4} - \frac{5x^3}{3} + \frac{4x^2}{2} \right]_1^3$ $= \frac{7}{12} - \left[ -\frac{27}{4} - \frac{7}{12} \right]$ $= \frac{95}{12}$

<b>Qn 6</b>	<b>N2018/RVHS Promo/9</b>
(i)	$\int (\ln x)^2 \, dx$ <p>Let <math>u = (\ln x)^2 \Rightarrow \frac{du}{dx} = \frac{2(\ln x)}{x}</math>      <math>v = 1 \Rightarrow \int v \, dx = x</math></p> $\int (\ln x)^2 \, dx = x(\ln x)^2 - \int \frac{2 \ln x}{x} x \, dx$ $= x(\ln x)^2 - 2 \int \ln x \, dx$ <p>Let <math>u = \ln x \Rightarrow \frac{du}{dx} = \frac{1}{x}</math>      <math>v = 1 \Rightarrow \int v \, dx = x</math></p> $\int (\ln x)^2 \, dx = x(\ln x)^2 - 2 \int \ln x \, dx$ $= x(\ln x)^2 - 2 \left[ x \ln x - \int \frac{1}{x} x \, dx \right]$ $= x(\ln x)^2 - 2x \ln x + 2x + C$
(ii)	<p>Volume</p> $= \pi(4)^2(e^4) - \pi \int_1^{e^4} x^2 \, dy$ $= 16\pi e^4 - \pi \int_1^{e^4} (\ln y)^2 \, dy$ $= 16\pi e^4 - \pi \left[ x(\ln x)^2 - 2x \ln x + 2x \right]_1^{e^4}$ $= 16\pi e^4 - \pi \left[ (e^4 \cdot 4^2 - 8e^4 + 2e^4) - 2 \right]$ $= 16\pi e^4 - \pi [10e^4 - 2]$ $= 6\pi e^4 + 2\pi \text{ units}^3$ 

Qn7	2018/YJC Promo/11
	 <p> <math>y = 3x</math>  <math>y = xe^{x^3}</math> </p> <p> <math>3x = xe^{x^3}</math>  From GC, <math>x = 1.03185, y = 3.09554</math> </p> <p> Volume = <math>\frac{1}{3}\pi(3.09554)^2(1.03186) - \pi \int_0^{1.03185} (xe^{x^3})^2 dx</math>  = 6.17 </p>

Qn 8	N2009/P1/11
(i)	
(ii)	$f(x) = xe^{-x^2}$ $f'(x) = e^{-x^2} + xe^{-x^2}(-2x) = e^{-x^2} - 2x^2e^{-x^2}$ $e^{-x^2} - 2x^2e^{-x^2} = 0$ $e^{-x^2}(1 - 2x^2) = 0$ $e^{-x^2} = 0 \quad \text{or} \quad x^2 = \frac{1}{2} \Rightarrow x = \pm \frac{1}{\sqrt{2}}$ <p>(N.A :<math>\because e^{-x^2} &gt; 0</math>)</p> <p>When <math>x = \frac{1}{\sqrt{2}}</math>, <math>y = \frac{1}{\sqrt{2}}e^{-\left(\frac{1}{\sqrt{2}}\right)^2} = \frac{1}{\sqrt{2e}}</math></p> <p>When <math>x = -\frac{1}{\sqrt{2}}</math>, <math>y = -\frac{1}{\sqrt{2}}e^{-\left(-\frac{1}{\sqrt{2}}\right)^2} = -\frac{1}{\sqrt{2e}}</math></p> <p>Coordinates of the turning points are <math>\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2e}}\right)</math> and <math>\left(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2e}}\right)</math></p>
(iii)	$\int_0^n f(x) dx = \int_0^n xe^{-x^2} dx$ $= \int_0^{n^2} \sqrt{u} e^{-u} \frac{1}{2\sqrt{u}} du$ $= \frac{1}{2} \int_0^{n^2} e^{-u} du$ $= \frac{1}{2} \left[ -e^{-u} \right]_0^{n^2}$ $= \frac{1}{2} \left( -e^{-n^2} + e^0 \right)$ $= \frac{1}{2} (1 - e^{-n^2})$ <p>Area of region between C and the positive <math>x</math>-axis</p> $= \lim_{n \rightarrow \infty} \int_0^n f(x) dx = \frac{1}{2} (1 - 0) = 0.5 \text{ units}^2$ <div style="border: 1px solid black; padding: 10px; margin-left: 20px;"> <math display="block">u = x^2 \Rightarrow x = \sqrt{u} \Rightarrow \frac{dx}{du} = \frac{1}{2\sqrt{u}}</math> <p>When <math>x = 0 \Rightarrow u = 0</math></p> <p>When <math>x = n \Rightarrow u = n^2</math></p> </div>

<b>(iv)</b>	$\int_{-2}^2  f(x)  dx = 2 \int_0^2 f(x) dx \quad (\text{by symmetry})$ $= 2 \times \frac{1}{2} (1 - e^{-2^2})$ $= 1 - e^{-4}$
<b>(v)</b>	Vol of revolution $= \pi \int_0^1 (f(x))^2 dx$ $= \pi \int_0^1 (xe^{-x^2})^2 dx$ $= 0.11570\pi$ $= 0.36349$ $\approx 0.363 \text{ units}^3$

Qn 9	2016/RVHS/Promo/11
<b>(a)</b> $\int \frac{x^2}{(x^2+9)^2} dx,$ $= \int \frac{9 \tan^2 \theta}{(9 \tan^2 \theta + 9)^2} 3 \sec^2 \theta d\theta$ $= \int \frac{9 \tan^2 \theta}{81 \sec^4 \theta} 3 \sec^2 \theta d\theta$ $= \frac{1}{3} \int \frac{\tan^2 \theta}{\sec^2 \theta} d\theta$ $= \frac{1}{3} \int \sin^2 \theta d\theta$ $= \frac{1}{6} \int (1 - \cos 2\theta) d\theta$ $= \frac{1}{6} \left[ \theta - \frac{\sin 2\theta}{2} \right] + c$ $= \frac{1}{6} \theta - \frac{\sin 2\theta}{12} + c$ $= \frac{1}{6} \theta - \frac{2 \sin \theta \cos \theta}{12} + c$ $= \frac{1}{6} \tan^{-1} \frac{x}{3} - \frac{1}{12} (2) \frac{x}{\sqrt{x^2+9}} \frac{3}{\sqrt{x^2+9}} + c$ $= \frac{1}{6} \tan^{-1} \frac{x}{3} - \frac{x}{2(x^2+9)} + c$	$x = 3 \tan \theta \Rightarrow \frac{dx}{d\theta} = 3 \sec^2 \theta,$ <div style="border: 1px solid black; padding: 10px; margin-top: 10px;"> <math display="block">x = 3 \tan \theta</math> <math display="block">\Rightarrow \tan \theta = \frac{x}{3}</math> <math display="block">\Rightarrow \sin \theta = \frac{x}{\sqrt{x^2+9}}</math> <math display="block">\Rightarrow \cos \theta = \frac{3}{\sqrt{x^2+9}}</math> </div>

<b>(b)(i)</b>	<p>When <math>a = 3</math>,  Volume generated</p> $= 2 \times \pi \int_0^3 \left( \frac{x}{x^2 + 9} \right)^2 dx$ $= 2\pi \left[ \frac{1}{6} \tan^{-1} \frac{x}{3} - \frac{x}{2(x^2 + 9)} \right]_0^3$ $= 2\pi \left[ \frac{1}{6} \cdot \frac{\pi}{4} - \frac{3}{36} \right]$ $= \frac{1}{12} \pi^2 - \frac{1}{6} \pi \text{ units}^3$
<b>(ii)</b>	<p>In the actual hourglass, the neck connecting the two glass bulbs constitutes to a volume which is not accounted for in the theoretical working.</p>
<b>(iii)</b>	$2 \times \pi \int_0^a \left( \frac{x}{x^2 + 9} \right)^2 dx$ $= 2 \times \left( \frac{1}{12} \pi^2 - \frac{1}{6} \pi \right)$ $\int_0^a \left( \frac{x}{x^2 + 9} \right)^2 dx = \frac{1}{12} \pi - \frac{1}{6}$ $\left[ \frac{1}{6} \tan^{-1} \frac{x}{3} - \frac{x}{2(x^2 + 9)} \right]_0^a = \frac{1}{12} \pi - \frac{1}{6}$ $\frac{1}{6} \tan^{-1} \frac{a}{3} - \frac{a}{2(a^2 + 9)} - \frac{1}{12} \pi + \frac{1}{6} = 0$ <p>Using GC,  <math>a = 4.8600148 \approx 4.86</math></p>

Qn 10	
<b>(a)</b> <p>Using GC, intersection between <math>(y - 2)^2 = x + 1</math> and <math>y + 2x = 6</math> occur when <math>x = 3</math> or <math>x = \frac{5}{4}</math>.</p> <p>Also, <math>(y - 2)^2 = x + 1 \Rightarrow y = 2 \pm \sqrt{x+1}</math></p> $\text{Volume generated} = \pi \int_{-1}^{\frac{5}{4}} (2 + \sqrt{x+1})^2 dx + \pi \int_{\frac{5}{4}}^3 (6 - 2x)^2 dx - \pi \int_{-1}^3 (2 - \sqrt{x+1})^2 dx$ $= 78.5725 = 78.6 \quad (3 \text{ s.f.})$	
<b>(b)</b> <p>Points of intersection of curves are <math>(-5, 9)</math> and <math>(0, 4)</math>.</p> $\text{Volume} = \pi \int_0^9 (-2 - \sqrt{y})^2 dy - \pi \int_0^4 (-2 + \sqrt{y})^2 dy - \pi \int_4^9 \left(\frac{16 - y^2}{13}\right)^2 dy$ $= 466.52 - 8.3775 - 107.66$ $= 350.48$ $= 350 \text{ (3 s.f.)}$	