eg number
** * ****
_



Chemistry

9729

27 September 2017

Section A

40 minutes

Additional Materials: OMR Sheet and Data Booklet

READ THESE INSTRUCTIONS FIRST

This booklet contains Section A of your paper.

Write your name, class and register number in the spaces provided at the top of this page.

There are twenty questions in this section. Answer all questions. For each question, there are four possible answers labelled A, B, C and D. Choose the one you consider correct and record your choice in soft pencil on the OMR answer sheet.

Read very carefully the instructions on the use of OMR answer sheet.

You are advised to fill in the OMR Answer Sheet as you go along; no additional time will be given for the transfer of answers once the examination has ended.

Use of OMR Answer Sheet

Ensure you have written your name, class register number and class on the OMR Answer Sheet.

Use a 2B pencil to shade your answers on the OMR sheet; erase any mistakes cleanly. Multiple shaded answers to a question will not be accepted.

For shading of class register number on the *OMR sheet*, please follow the given examples:

If your register number is 1, then shade <u>01</u> in the index number column. If your register number is 21, then shade <u>21</u> in the index number column.

Answer all questions in this section.

For each question there are four possible answers, A, B, C, and D. Choose the one you consider to be correct.

Methylamine, CH₃NH₂ is a colourless gas which has a strong odour similar to fish. It 1 burns in oxygen according to the following equation.

$$CH_3NH_2(g) + \frac{13}{4}O_2(g) \longrightarrow CO_2(g) + NO_2(g) + \frac{5}{2}H_2O(I)$$

A sample of 10 cm³ of methylamine was burned in 50 cm³ of oxygen. The products were cooled to room temperature and treated with an excess of aqueous sodium hydroxide.

What would be the final volume of the resultant mixture of gases?

A 17.5 cm³

B 20.0 cm³ C 27.5 cm³

D 37.5 cm³

How many atoms are present in 24 cm³ of carbon dioxide gas under room conditions? 2

 $B = \frac{3 \times 6.02 \times 10^{23}}{10^3}$

 $C = \frac{3 \times 6.02 \times 10^{23}}{24000}$

D $3 \times 24 \times 6.02 \times 10^{23}$

25.0 cm³ of 0.05 mol dm⁻³ acidified NH₄VO₃ was reduced by sulfur dioxide to form a 3 deep-blue solution containing V^{n+} ions.


The V^{n+} ions are then re-oxidised to its original oxidation state by 12.50 cm³ of 0.02 mol dm⁻³ acidified KMnO₄.

What is the oxidation state of vanadium in the deep-blue solution?

A 0

D +6

The value of pV is plotted against p at the same temperature for four gases, where p is the pressure and V is the volume of the gas.



Which of the following statements are correct?

- 1 All gases deviate most from ideal gas behaviour at high pressures.
- 2 The intermolecular forces of attraction in Gas Y is more significant than that in Gas X.
- 3 Gas W is H₂O; Gas X is HF; Gas Y is CH₃F; Gas Z is Ne.

A 1 only

B 1 and 2 only

C 1 and 3 only

D 1, 2 and 3

5 Use of the Data Booklet is relevant to this question.

The ion X4+ contains 28 electrons and 42 neutrons.

Which of the following statements about X^{4+} or X^{2+} is correct?

- A X2+ has less protons than 59Co2+.
- B X4+ has the same electronic configuration as 58Ni²⁺.
- C X4+ has the same nucleon number as 75As3+.
- D X²⁺ has a lower tendency to form covalent compounds than X⁴⁺.

The angle of deflection of a beam of ${}^{27}_{13}\text{Al}^{3+}$ in an electric field is +5.6 °. 6


What is a possible species that will give an angle of deflection of 4.2 ° when passed through the same electric field?

- B $^{24}_{12}\text{Mg}^{2+}$ C $^{32}_{16}\text{S}^{2-}$ D $^{52}_{24}\text{Cr}^{2+}$

1-chlorobutane can be synthesised by reacting butan-1-ol with thionyl chloride, SOCl2. 7

Distillation is performed to separate the 1-chlorobutane and the unreacted butan-1-ol upon completion of the reaction. A distillation set-up is shown below where the organic compound with the lower boiling point will distil out first.

Which region indicates an area where there is a higher proportion of butan-1-ol than 1-chlorobutane?



8 The melting points of two aluminium halides, A/X₃ are given below.

Compound	Formula	Melting Point/ °C
Aluminium fluoride	A/F ₃	1291
Aluminium chloride	A/C/3	178

Which pairs of factors provide the reasons to explain for the differences in the melting points?

- 1 charge density of cation
- 2 radius of anion
- 3 strength of the AI-X bond
- 4 distortion of electron cloud

A 1 and 3 only

B 1 and 4 only

C 2 and 3 only

D 2 and 4 only

9 Nickel is a by-product formed by a series of changes in which the rate-determining step is the radioactive decay of cobalt-60, a radiation source for cancer treatment. This radioactive decay is a first-order reaction with a half-life of 5.3 years.

How long would it take for a radiation cartridge of cobalt-60, originally nickel-free, to contain a molar proportion of cobalt to nickel of 1:7?

A 0.8 years

B 14.9 years

C 15.9 years

D 37.1 years

10 The reaction between nitrogen monoxide, NO and oxygen, O₂ takes place as follows:

$$2NO \longrightarrow N_2O_2$$
 (fast)
$$N_2O_2 + O_2 \longrightarrow 2NO_2$$
 (slow)

Which of the following statement is correct?

- A The overall reaction is third order.
- B The reaction is an elementary reaction.
- C The rate equation is Rate = $k [N_2O_2] [O_2]$.
- D The activation energy of the first step is higher than the second step.

11 A 0.0113 mol sample of ammonium nitrate is introduced into an evacuated vessel with a volume of 0.83 dm³. The vessel, of constant volume, is heated to a constant temperature such that the equilibrium below is established.

$$NH_4NO_3(s) \implies NH_3(g) + HNO_3(g)$$

Given that the percentage dissociation of ammonium nitrate at equilibrium is 66.8%. what is the value of the equilibrium constant, K_c ?

A 5.70×10^{-5}

B 8.27 × 10⁻⁵

C 0.0152

D 0.0183

The key stage in the manufacture of sulfuric acid is the reaction between sulfur dioxide 12 and oxygen.

$$2SO_2(g) + O_2(g) \implies 2SO_3(g)$$
 $\Delta H = -197 \text{ kJ mol}^{-1}$

Which of the following statements are correct for an increase in temperature?

- 1 The equilibrium constant increases.
- 2 The rate constant for the forward reaction increases.
- 3 The rate constant for the backward reaction increases.
- The activation energy for the forward reaction decreases.

A 1, 2, 3 and 4

B 2, 3 and 4 only

C 2 and 3 only

D 3 only

13 The dihydrogen phosphate ion, H₂PO₄⁻ can act as a base or an acid.

$$H_2PO_4^- + H_2CO_3 \implies H_3PO_4 + HCO_3^-$$
 $K_c < 1$

$$H_2PO_4^- + HCO_3^- \implies HPO_4^{2-} + H_2CO_3$$

The conjugate _____ of H₂PO₄⁻ is _____, and H₃PO₄ is a stronger acid than _____.

A acid; HPO₄²⁻; H₂CO₃

B acid; H₃PO₄; HCO₃-

C base; HPO₄²⁻; HPO₄²⁻

D base; HPO₄²⁻; H₂CO₃

14 Values for the ionic product of water, Kw, at two different temperatures are given below.

Temperature / °C	K _w / mol² dm ⁻⁶
10	2.90 x 10 ⁻¹⁵
25	1.00 x 10 ⁻¹⁴

What is correct for pure water at 10 °C?

A pH = 7

- B pH > 7
- C $[H^+] = 2.90 \times 10^{-8} \text{ mol dm}^{-3}$
- D [H+] > [OH-]

15 A mixture is made by mixing equal volumes of a solution of H₂SO₄ (aq) of pH 2 and of a solution of HC/ (aq) of pH 3.

What is the pH of the resulting solution?

- A 1.67
- B 1.96
- C 2.26
- D 2.50

16 The values of the solubility products at 25 °C of three metal sulfides are tabulated below.

Compound	Value of solubility product
CuS	8.5 x 10 ⁻⁴⁵
Ag ₂ S	1.6 x 10 ⁻⁴⁹
Bi ₂ S ₃	1.1 x 10 ⁻⁷³

The dissolution process of these metal sulfides is endothermic.

Which statements about the metal sulfides are correct?

- 1 The value of the solubility products of the metal sulfides increases with temperature.
- 2 The value of the solubility products of the metal sulfides decreases when there is a common ion effect as the solubility decreases.
- 3 The molar solubility of the metal sulfides decreases in the following order: CuS > Ag₂S > Bi₂S₃.
- 4 The molar solubility of the metal sulfides increases when pH decreases.
- A 1 and 2 only

B 1 and 4 only

C 2 and 3 only

D 3 and 4 only

17 Cyanopropyne has the following structure:

CH₃C≡CCN

Which statements about the molecule are correct?

- 1 There are 6 sigma bonds and 2 pi bonds.
- 2 All the carbon atoms lie in the same plane.
- 3 Only two of the carbon atoms are sp hybridised.
- 4 One of the terminal carbon atoms has a tetrahedral geometry.
- A 1 and 2 only

B 2 and 3 only

C 2 and 4 only

- D 3 and 4 only
- 18 How many constitutional and cis-trans isomers are there for dichloropropene, C₃H₄Cl₂?
 - A 5
- B 6
- **C** 7

D 8

19 Limonene is an oil formed in the peel of citrus fruits.

When limonene reacts with excess cold dilute KMnO₄, how many more chiral centres than limonene does the product possess?

- A 2
- B 3
- C 4

D 5

20 Which two-stage process will give the greatest yield of CH₂(Br)CH₂CH₃?

A $CH_3CH(Br)CH_2CH_3$ $\xrightarrow{ethanolic KOH, heat}$ $\xrightarrow{Br_2 \text{ in organic solvent}}$ \xrightarrow{C} $\xrightarrow{CH_3CH=CHCH_3}$ $\xrightarrow{H_2 (g), Pt}$ $\xrightarrow{Br_2 (g), uv \ light}$ $\xrightarrow{Br_2 \text{ in organic solvent}}$ $\xrightarrow{Br_2 \text{ in organic solvent}}$