

# Sec 4 Pure Chem Mock Paper Set 6 Ans


## Solutions for 2019 Chemistry EOY 6092 paper 1 and paper 2

Paper 1				
1. D	2. D	3. A	4. C	5. B
6. D	7. B	8. B	9. B	10. C
11. B	12. D	13. D	14. C	15. [A]
16. C	17. B	18. A	19. A	20. B
21. D	22. A	23. B	24. B	25. C
26. A	27. A	28. A	29. B	30. A
31. A	32. B	33. A	34. A	35. C
36. B	37. D	38. A	39. C	40. B

## Paper 2






S/No.	Answers	Marks
1a	Similarity: both have 2 protons (and or 2 electrons) Differences: Helium 3 has 1 neutron while helium 4 has 2 neutrons	1 1
1bi	Helium does not form any compounds as it is a noble gas (or has fully filled outermost(valence) shell) / duplet electronic configuration. <i>Reject: helium can form compounds (it does not show any unusual characteristic from the answer).</i>	1
1bii	At extreme high pressures, particles are close together which increases the chance of effective collisions causing the reactants to react.	1
1c	Used as insulators (for cables) The ions or electrons are held in fixed lattice structures (or localized) and cannot carry electric charges to conduct electricity. OR Include uses such as electrolyte (dissolve in water and ions can conduct electricity)	1
Total		5
2a	Ammonia it is collected by upward delivery method and dried by calcium oxide.	1 1
2b	Sodium chloride / ammonium chloride	1
2c	$\text{Na}_3\text{N} + 4\text{HCl} \rightarrow 3\text{NaCl} + \text{NH}_4\text{Cl}$	1
2d	<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;">3</div> <div style="display: flex; align-items: center;"> <div style="margin-left: 10px;"> <p>3-</p> <p>legend: X - rep. of Na O - rep. of N</p> </div> </div> </div>	2



Total		6
3a	Energy change for bond formation (making) Enthalpy change is <u>negative</u> indicates <u>less energy</u> is <u>absorbed</u> for breaking H-H and Cl-Cl than energy released for making H-Cl 6 underlined phrase 2 marks, 3 underlined phrase 1 mk. No 1/2 mark	1 1 1
3b	<p>Shape + formulae + correct direction for Ea and ΔH - 2 mks. Every 2 correct - 1 mk. No double arrows and no 1/2 mark</p>	2
3c	No. of moles of $\text{Cl}_2 = \frac{14.2}{71} = 0.2 \text{ mol}$ Energy change = $0.2 \times (-184) = -36.8 \text{ kJ}$	1 1
3d	Oxidation state of hydrogen increases from 0 in $\text{H}_2$ to +1 in HCl. It is oxidized. Oxidation state of chlorine decreases from 0 in $\text{Cl}_2$ to -1 in HCl. It is reduced. When both oxidation and reduction takes place together, it is a redox reaction. 3 points - 2 mk; 2 points - 1 mk and no 1/2 mark	2
Total		9



4e	By using powder, this increases the <u>surface area to volume ratio</u> . There are <u>more reacting surfaces</u> for $\text{H}^+$ ions to collide onto. This <u>increases the frequency of effective collision</u> and <u>increases the rates of reaction</u> . Note: reject acid molecules. 4 points 2 mks; 2 points 1 mk; no ½ mark	1
Total		11
5ai	Carbon monoxide is produced due to incomplete combustion of petrol (or carbon containing fuel). Oxides of nitrogen is produced due to high temperature in engines	1
5aii	Nitrogen and carbon dioxide Note: Accept oxygen as it is indicated in the passage.	1
5b	Petroleum is heated in a furnace and it vaporizes. The vapour rises up the fractionating column and is <u>separated according to boiling points</u> . Fractions with <u>shorter chains</u> have lower boiling points such as petrol is distilled near to the top of the fractionating column. Note: need to mention how petrol is obtained. Fractions are obtained near to the top, not by speed	1
Total		6
5ci	$\text{N}_2 + 3 \text{H}_2 \rightleftharpoons 2 \text{NH}_3$	1
5cii	From the equation, gases are entering the reactor in the ratio of 1 molecule of nitrogen to 3 molecules of hydrogen. According to Avogadro's Law, it states that equal volumes of gases at the same temperature and pressure contain equal numbers of molecules, which is the proportion demanded by the equation	1
5ciii	When ammonium nitrate dissolves in water, the forces of attraction between water molecules and ammonium nitrate is strong enough to pull ammonium nitrate from the lattice structure.	1
5civ	% of nitrogen in ammonium nitrate = $\frac{2(14)}{14+4+14+3(16)} \times 100\% = 35\%$ % of nitrogen in urea = $\frac{2(14)}{12+16+2(14+2)} \times 100\% = 33.3\%$ This shows that urea contains more mass of nitrogen. Note: need to include the justification statement.	2
Total		7



6a	C: ethene / C <sub>2</sub> H <sub>4</sub> D: dibromoethane / C <sub>2</sub> H <sub>4</sub> Br <sub>2</sub> Accept if student name it as structural formula.	E: ethane / C <sub>2</sub> H <sub>6</sub> F: chloroethene / C <sub>2</sub> H <sub>3</sub> Cl	4
6bi	C <sub>15</sub> H <sub>32</sub> → 4C <sub>2</sub> H <sub>4</sub> + C <sub>3</sub> H <sub>8</sub> + C <sub>4</sub> H <sub>8</sub> Conditions 600°C with Al <sub>2</sub> O <sub>3</sub> /SiO <sub>2</sub> as catalyst		1
6bii	 OR  Reject:   		1
Total			6
7ai	Gas G. 2H <sup>+</sup> (aq) + 2e <sup>-</sup> → H <sub>2</sub> (g) Gas H. 2Cl <sup>-</sup> (aq) → Cl <sub>2</sub> (g) + 2e <sup>-</sup> Gas I. 4OH <sup>-</sup> (aq) → O <sub>2</sub> (g) + 2H <sub>2</sub> O (l) + 4e <sup>-</sup>		1 1 1
7aii	At the beginning, there is a large concentration of Cl <sup>-</sup> . It is discharged preferentially as compared to OH <sup>-</sup> . Hence chlorine is obtained. As the electrolysis proceeds, there are lesser concentration of Cl <sup>-</sup> . OH <sup>-</sup> ions are preferentially discharged. Hence oxygen is obtained.		1 1
7aiii	Electrolysis of acid will give acidic fumes which is corrosive / irritates breathing system / harmful for health.		1
7bi	Aqueous silver nitrate / aqueous silver sulfate		1
7bii	Concentration of silver solution remains unchanged		1
7biii	Ag (s) → Ag <sup>+</sup> (aq) + e <sup>-</sup> Silver electrode dissolves/becomes smaller		1 1
Total			10

8 ai	X High melting point / high density / formed coloured compounds / have formed ions of multiple valencies Any 2 correct evidences Accept Z.	2
8 aii	Y Formed ions of charge +1 / low density / low melting point	2
8 b	2 Y + 2 HCl → 2 YCl + H <sub>2</sub>	1
8 c	Z becomes smaller / colour of solution changes from blue to colourless / reddish brown (pink) solid deposited	1
8di	 Legend • - rep e of Cl x - rep e of W	2
8dii	 Note: size of C << Fe Number of C should be less than 5 as the indicated percentage of C is "small amount". Atoms of different sizes disrupt the orderly arrangement and prevent layers from sliding.	1 1
Total		10
Either 9		
ai	Glucose reacts with yeast in air tight container at 37°C to form ethanol. The mixture of ethanol undergoes fractional distillation to obtain pure ethanol.	1 1

a ii	As a solvent / as alcoholic beverage rej: perfume / alcohol without explaining further	1
bi	$C_2H_5ONa + CH_3CH_2COCl \rightarrow NaCl + CH_3CH_2COOC_2H_5$	1
bii	Ethyl propanoate	1
biii	Terylene	1
		2
ci	Carboxylic acid	1
cii	Iron will corrode/react with the acids.	1
	<b>Total</b>	<b>10</b>

OR																		
9a	<table border="1"> <tr> <td></td> <td>Nylon</td> <td>Poly(propene)</td> </tr> <tr> <td>structure</td> <td colspan="2">macromolecule</td> </tr> <tr> <td>Types of polymerisation</td> <td>condensation</td> <td>addition</td> </tr> <tr> <td>monomers</td> <td> <math display="block">\text{H}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{[shaded box]}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{H}</math> <p>and</p> <math display="block">\text{H}-\text{N}-\text{[box]}-\text{N}-\text{H}</math> <math display="block">\text{H} \quad \text{H}</math> </td> <td> <math display="block">\begin{array}{c} \text{H} \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}</math> </td> </tr> <tr> <td>linkages</td> <td>amide</td> <td>c-c long chain</td> </tr> </table>		Nylon	Poly(propene)	structure	macromolecule		Types of polymerisation	condensation	addition	monomers	$\text{H}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{[shaded box]}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{H}$ <p>and</p> $\text{H}-\text{N}-\text{[box]}-\text{N}-\text{H}$ $\text{H} \quad \text{H}$	$\begin{array}{c} \text{H} \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$	linkages	amide	c-c long chain		
	Nylon	Poly(propene)																
structure	macromolecule																	
Types of polymerisation	condensation	addition																
monomers	$\text{H}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{[shaded box]}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{H}$ <p>and</p> $\text{H}-\text{N}-\text{[box]}-\text{N}-\text{H}$ $\text{H} \quad \text{H}$	$\begin{array}{c} \text{H} \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$																
linkages	amide	c-c long chain																
	<ul style="list-style-type: none"> <li>Both nylon and poly(propene) have macromolecular structures.</li> <li>Nylon is made by many monomers of dicarboxylic acid and diamine (accept structures) while poly(propene) is made by many monomers of propene (accept structures)</li> <li>Nylon is made by the process of condensation polymerization while poly(propene) is made by the process of addition polymerisation</li> <li>Nylon is joined by amide linkages while in poly(propene), it is joined by the opening of the double bond to form a long carbon chain.</li> </ul>		1 2  1 1  1															
	Note: accept tabulation.																	
9bi	<p>Accept condensed formula=</p> $-\text{N}(\text{H})-(\text{CH}_2)_6-\text{C}(=\text{O})-\text{N}(\text{H})-(\text{CH}_2)_6-\text{C}(=\text{O})-$		2															
9bii	Polypropene will not have any visible observation with dilute acids Nylon will have a hole/corrode/damage with dilute acids. Reject: "react", as it is used in the question.		1 1															
	Total		10															